

Bilaga 3 Exklusionslistor

Kapitel 3.1

Record nr, författare, artikel mm.

4200 *Armitage FE, Richardson DE, Li KC. Polymeric contrast agents for magnetic resonance imaging: synthesis and characterization of gadolinium diethylenetriaminepentaacetic acid conjugated to polysaccharides. *Bioconjug Chem* 1990;1:365-74.

13672 *Bailey RR, Rogers TG, Tait JJ. Measurement of glomerular filtration rate using a single injection of ⁵¹Cr-Edetic acid. *Australas Ann Med* 1970;19:255-8.

13686 *Bianchi C, Bonadio M, Donadio C, Tramonti G, Figus S. Measurement of glomerular filtration rate in man using DTPA-^{99m}Tc. *Nephron* 1979;24:174-8.

13517 *Bird NJ, Peters C, Michell AR, Peters AM. Comparison of GFR Measurements Assessed From Single Versus Multiple Samples. *American Journal of Kidney Diseases* 2009;54:278-288.

1782 *Brion LP, Fleischman AR, McCarton C, Schwartz GJ. A simple estimate of glomerular filtration rate in low birth weight infants during the first year of life: noninvasive assessment of body composition and growth. *J Pediatr* 1986;109:698-707.

13539 *Brochner Mortensen J. The glomerular filtration rate during moderate hyperglycemia in normal man. *Acta Medica Scandinavica* 1973;194:31-37.

13541 *Brochner-Mortensen J. The effect of glucose on the glomerular filtration rate in normal man. A preliminary report. *Acta medica Scandinavica* 1971;189:109-111.

13515 *Bumb A, Brechbiel MW, Choyke P. Macromolecular and dendrimer-based magnetic resonance contrast agents. *Acta Radiol.* 2010;51:751-67.

12920 *Buyschaert M, Joudi I, Wallemacq P, Hermans MP. Performance of serum cystatin-C versus serum creatinine in subjects with type 1 diabetes. *Diabetes Care.* 2003;26:1320.

1818 *Cirillo M, Anastasio P, De Santo NG. Relationship of gender, age, and body mass index to errors in predicted kidney function. *Nephrol Dial Transplant* 2005;20:1791-8.

13518 *Cittanova ML. Editorial II: Is peri-operative renal dysfunction of no consequence? *British Journal of Anaesthesia* 2001;86:164-166.

1864 *Dunea G, Freedman P. Renal clearance studies. *JAMA* 1968;205:170-1.

13533 *Epstein M. Monitoring glomerular filtration rate: Improving on creatinine-based methods: Means vary in reliability and practicality - But you needn't sacrifice either. *Journal of Critical Illness* 1999;14:166-172.

13559 *Grubb A, Bjork J, Nyman U, Pollak J, Bengzon J, Ostner G, et al. Cystatin C, a marker for successful aging and glomerular filtration rate, is not influenced by inflammation. *Scand J Clin Lab Invest* 2011.

12995 *Grubb A, Nordin G. Notable steps in obtaining improved estimates for glomerular filtration rate. *Clinical Chemistry* 2006;52:169-170.

13525 *Haycock GB. Creatinine, body size and renal function. *Pediatric Nephrology* 1989;3:22-24.

6005 *Jacobi D, Lavigne C, Halimi JM, Fierrard H, Andres C, Couet C, et al. Variability in creatinine excretion in adult diabetic, overweight men and women: consequences on creatinine-based classification of renal disease. *Diabetes Res Clin Pract* 2008;80:102-7.

2017 *Kampf D, Schurig R, Korsukewitz I, Bruckner O. Cefoxitin pharmacokinetics: relation to three different renal clearance studies in patients with various degrees of renal insufficiency. *Antimicrob Agents Chemother* 1981;20:741-6.

13530 *Kerr DNS, Davison JM. The assessment of renal function. *British Journal of Hospital Medicine* 1975;14:360-372.

4219 *Lebduskova P, Kotek J, Hermann P, Vander Elst L, Muller RN, Lukes I, et al. A gadolinium(III) complex of a carboxylic-phosphorus acid derivative of diethylenetriamine covalently bound to inulin, a potential macromolecular MRI contrast agent. *Bioconjug Chem* 2004;15:881-9.

13710 *London GM, Safar ME, Levenson JA, Simon AC, Temmar MA. Renal filtration fraction, effective vascular compliance, and partition of fluid volumes in sustained essential hypertension. *Kidney Int* 1981;20:97-103.

13397 *Maillard N, Mehdi M, Thibaudin L, Berthoux F, Alamartine E, Mariat C. Creatinine-based GFR predicting equations in renal transplantation: Reassessing the tubular secretion effect. *Nephrology Dialysis Transplantation* 2010;25:3076-3082.

13141 *Mojiminiyi OA, Al Shayji I, George S. Serum cystatin C could replace serum creatinine as an indicator of glomerular filtration rate in routine practice. *European Journal of Laboratory Medicine* 1998;6:98-101.

13535 *Moonie A, Mooney A. The estimated glomerular filtration rate in primary and secondary care. *British Journal of Hospital Medicine* 2007;68:638-639.

13702 *Morris K, Skinner R, Keir M, Palmer J, Lambert H, Pearson A, et al. Problems with the single injection method for the measurement of the glomerular filtration rate in man. *Clin Sci (Lond)* 1991;81:831-3.

13538 *Mourad A, Carney S, Gillies A, Hibberd A, Trevillian P, Nanra R. Measurement of glomerular filtration rate in renal transplant recipients: A comparison of methods. *Nephrology* 2002;7:77-82.

13145 *Mouton R, Holder K. Laboratory tests of renal function. *Anaesthesia and Intensive Care Medicine* 2006;7:240-243.

13537 *Neuberger J. Renal failure late after liver transplantation. *Liver Transplantation* 2002;8:922-924.

13721 *O'Dell RM. Determination of glomerular filtration rate by use of single injection of iodine-131 labelled sodium diatrizoate. *J Nucl Med* 1966;7:470-3.

1470 *Perico N, Gaspari F, Remuzzi G. Assessing renal function by GFR prediction equations in kidney transplantation. *Am J Transplant* 2005;5:1175-6.

7235 *Poggio ED, Batty DS, Flechner SM. Evaluation of renal function in transplantation. *Transplantation* 2007;84:131-6.

13480 *Ravnskov U. On renal handling of plasma proteins: With special reference to (alpha)(2) microglobulin, (beta)(2) microglobulin, lysozyme and albumin. *SOURCE Scandinavian Journal of Urology and Nephrology* (1973) Sup.20. Date of Publication: 1973.

13705 *Reams GP, Lau A, Hamory A, Bauer JH. Amlodipine therapy corrects renal abnormalities encountered in the hypertensive state. *Am J Kidney Dis* 1987;10:446-51.

13221 *Rodrigo E, Fernandez-Fresnedo G, Castaneda O, Arias M. Estimation of renal function in adult kidney transplant recipients by equations. *Transplantation Reviews* 2007;21:1-16.

2247 *Rosenkranz B, Fejes-Toth G, Diener U, Frolich JC. Effects of sulfipyrazone on renal function and prostaglandin formation in man. *Nephron* 1985;39:237-43.

7537 *Rule AD. Understanding estimated glomerular filtration rate: implications for identifying chronic kidney disease. *Curr Opin Nephrol Hypertens* 2007;16:242-9.

- 13252 *Sharma AP, Filler G. Monitoring kidney function and renal disease in children following transplant. *Pediatric Health* 2009;3:155-163.
- 1565 *Siebenhofer A, Meinitzer A, Brandmair J, Plank J, Pieber TR. Increased glomerular filtration rate in patients after reconstructive surgery on the abdominal aorta. *Diabet Med* 2001;18:464-8.
- 13359 *Smith WH. Measurement of the filtration rate. In: *The Kidney: Structure and function in health and disease*: Oxford University Press, USA; 1951. p 39-62.
- 13670 *Stolz A, Hoizey G, Toupance O, Lavaud S, Vitry F, Chanard J, et al. Evaluation of sample bias for measuring plasma iohexol clearance in kidney transplantation. *Transplantation*. 2010;89:440-5.
- 13534 *Van Stralen KJ, Jager KJ, Zoccali C, Dekker FW. Agreement between methods. *Kidney International* 2008;74:1116-1120.
- 13528 *Wood SM. Assessment of renal functions in hypertensive pregnancies. *Clinical Obstetrics and Gynecology* 1977;4:747-758.
- 209 *Work DF, Schwartz GJ. Estimating and measuring glomerular filtration rate in children. *Curr Opin Nephrol Hypertens* 2008;17:320-5.
- 13523 *Yeh SDJ. Recent advances in the urological application of radionuclides. *Current Opinion in Urology* 1993;3:76-85.
- 2426 *Zahran A, Qureshi M, Shoker A. Comparison between creatinine and cystatin C-based GFR equations in renal transplantation. *Nephrol Dial Transplant* 2007;22:2659-68.
- 13707 *Zech PY, Chau NP, Pozet N, Labeeuw M, Hadj-Aissa A. Ranitidine kinetics in chronic renal impairment. *Clin Pharmacol Ther* 1983;34:667-72.
- 13532 Direct measurement of glomerular filtration rate. *Nephrology* 2005;10:S168-S171.
- 34 *Aaronson IA, Mann MD. Measurement of glomerular filtration rate in children using technetium-99m diethylenetriamine penta-acetic acid. *S Afr Med J* 1985;67:507-9.
- 4272 *Abraham PA, Opsahl JA, Rachael KM, Asinger R, Halstenson CE. Renal function during erythropoietin therapy for anemia in predialysis chronic renal failure patients. *Am J Nephrol* 1990;10:128-36.
- 211 *Adefuin PY, Gur A, Siegel NJ, Spencer RP, Hayslett JP. Single subcutaneous injection of iothalamate sodium I 125 to measure glomerular filtration rate. *JAMA* 1976;235:1467-9.
- 212 *Agarwal R. Ambulatory GFR measurement with cold iothalamate in adults with chronic kidney disease. *Am J Kidney Dis* 2003;41:752-9.
- 4298 *Agarwal R. Reproducibility of renal function measurements in adult men with diabetic nephropathy: research and clinical implications. *Am J Nephrol* 2007;27:92-100.
- 1709 *Ahmed SB, Bentley-Lewis R, Hollenberg NK, Graves SW, Seely EW. A Comparison of Prediction Equations for Estimating Glomerular Filtration Rate in Pregnancy. *Hypertens Pregnancy* 2009;1-13.
- 35 *Ahrenholz P, Falkenhagen D, Roy T, Esther G, Klinkmann H. Tc-99m-DTPA--a new test substance for detoxification devices. *Int J Artif Organs* 1982;5:185-90.
- 13625 *Aissa AH, Collard M, Pozet N. Separate calculation of glomerular and tubular clearance by means of renal scintigraphy with hippural and DTPA. *Nephron* 1985;40:401-404.
- 13650 *Aitken JM, Lindsay R, Hart DM. The redistribution of body sodium in women on long term oestrogen therapy. *Clinical Science and Molecular Medicine* 1974;47:179-187.
- 12871 *Al Wakeel JS, Hammad D, Al Suwaida A, Tarif N, Chaudhary A, Isnani A, et al. Validation of predictive equations for glomerular filtration rate in the Saudi population. *Saudi J Kidney Dis Transpl*. 2009;20:1030-7.

13475 *Alatas H, Manz F, Scharer K. Cr(51) EDTA single injection clearance to measure the glomerular filtration rate in renal disease. *Annals of the Academy of Medicine Singapore* 1975;4:202-206.

13570 *Alazraki N, Verba JW, Henry JE. Determination of glomerular filtration rate utilizing X ray fluorescence of cold iothalamate levels in plasma and urine. *Journal of Nuclear Medicine* 1975;16:510.

13567 *Alazraki N, Verba JW, Henry JE, Becker R, Taylor A, Jr., Halpern SE. Noninvasive determination of glomerular filtration rate using x-ray fluorescence. *Radiology* 1977;122:183-6.

13722 *Alestig K, Hood B, Vikgren P. [Determination of glomerular infiltration with a single injection of inulin (II). Comparison with introductory kidney function tests]. *Lakartidningen* 1966;63:1659-62.

13723 *Alestig K, Hood B, Vikgren P. [Determination of glomerular filtration with a single injection of inulin (I)]. *Lakartidningen* 1966;63:1554-9.

168 *Alexander K, Del Castillo JR, Ybarra N, Morin V, Gauvin D, Authier S, et al. Single-slice dynamic computed tomographic determination of glomerular filtration rate by use of Patlak plot analysis in anesthetized pigs. *Am J Vet Res* 2007;68:297-304.

1712 *Alinei P, Guignard JP. Assessment of glomerular filtration rate in infants. Comparison of three methods used in clinical practice. *Helv Paediatr Acta* 1987;42:253-62.

2441 *Allen DB, Fogo A, el-Hayek R, Langhough R, Friedman AL. Effects of prolonged growth hormone administration in rats with chronic renal insufficiency. *Pediatr Res* 1992;31:406-10.

13521 *Almen T, Dawson P, Katzberg RW, Harnish PP, Bettmann MA, Wolf GL. Use of iohexol clearance to determine glomerular filtration rate. *Investigative Radiology* 1994;29:S2-S7.

338 *Anastasio P, Cirillo M, Spitali L, Frangiosa A, Pollastro RM, De Santo NG. Level of hydration and renal function in healthy humans. *Kidney Int* 2001;60:748-56.

1713 *Anastasio P, Spitali L, Frangiosa A, Molino D, Stellato D, Cirillo E, et al. Glomerular filtration rate in severely overweight normotensive humans. *Am J Kidney Dis* 2000;35:1144-8.

4366 *Andersen TB, Eskild-Jensen A, Frokiaer J, Brochner-Mortensen J. Measuring glomerular filtration rate in children; can cystatin C replace established methods? A review. *Pediatr Nephrol* 2009;24:929-41.

213 *Anderson CF, Jaecks DM, Ballon HS, De Palma JR, Cutler RE. Renal handling of creatinine in nephrotic and non-nephrotic patients. *Clin Sci* 1970;38:555-62.

214 *Anderson CF, Sawyer TK, Cutler RE. Iothalamate sodium I 125 vs cyanocobalamin Co 57 as a measure of glomerular filtration rate in man. *JAMA* 1968;204:653-6.

1716 *Ansquer JC, Dalton RN, Causse E, Crimet D, Le Malicot K, Foucher C. Effect of fenofibrate on kidney function: a 6-week randomized crossover trial in healthy people. *Am J Kidney Dis* 2008;51:904-13.

1719 *Aperia A, Broberger O, Elinder G, Herin P, Zetterstrom R. Postnatal development of renal function in pre-term and full-term infants. *Acta Paediatr Scand* 1981;70:183-7.

1721 *Apple FS, Benson P, Abraham PA, Rosano TG, Halstenson CE. Assessment of renal function by inulin clearance: comparison with creatinine clearance as determined by enzymatic methods. *Clin Chem* 1989;35:312-4.

985 *Arant BS, Jr. Postnatal development of renal function during the first year of life. *Pediatr Nephrol* 1987;1:308-13.

4245 *Arant BS, Jr., Edelmann CM, Jr., Spitzer A. The congruence of creatinine and inulin clearances in children: use of the Technicon AutoAnalyzer. *J Pediatr* 1972;81:559-61.

- 13519 *Argani H, Dabiri S, Taghizadeh M, Pishahang P, Rezvan NH, Emami P. Estimation of glomerular filtration rate in renal transplants based on serum creatinine level after oral cimetidine. *Transplantation Proceedings* 2000;32:545-546.
- 3183 *Armangil D, Yurdakok M, Canpolat FE, Korkmaz A, Yigit S, Tekinalp G. Determination of reference values for plasma cystatin C and comparison with creatinine in premature infants. *Pediatr Nephrol* 2008;23:2081-3.
- 13444 *Atherton JC. Renal blood flow, glomerular filtration and plasma clearance. *Anaesthesia and Intensive Care Medicine* 2009;10:271-275.
- 93 *Aurell M. Renal clearance of 51Cr-EDTA-complex. *Strahlentherapie Sonderb* 1968;67:420-4.
- 171 *Aurell M. Accurate and feasible measurements of GFR--is the iohexol clearance the answer? *Nephrol Dial Transplant* 1994;9:1222-4.
- 13583 *Bacchetta J, Cochat P, Rognant N, Ranchin B, Hadj-Aissa A, Dubourg L. Which Creatinine and Cystatin C Equations Can Be Reliably Used in Children? *Clin J Am Soc Nephrol*. 2010.
- 999 *Back SE, Ljungberg B, Nilsson-Ehle I, Borga O, Nilsson-Ehle P. Age dependence of renal function: clearance of iohexol and p-amino hippurate in healthy males. *Scand J Clin Lab Invest* 1989;49:641-6.
- 94 *Backlund L, Goransson M, Muller-Suur R, Olsen L. Evaluation of glomerular filtration rate in infants. *Acta Paediatr Scand Suppl* 1983;305:77-8.
- 13360 *Bajaj G, Alexander SR, Browne R, Sakarcin A, Seikaly MG. 125Iodine-iothalamate clearance in children. A simple method to measure glomerular filtration. *Pediatr Nephrol* 1996;10:25-8.
- 37 *Balachandran S, Toguri AG, Petrusick TW, Abbott LC. Comparative evaluation of quantitative glomerular filtration rate measured by isotopic and nonisotopic methods. *Clin Nucl Med* 1981;6:150-3.
- 2460 *Balint P, Forgacs I. CONSIDERATIONS ON THE EVALUATION OF THE CLEARANCE METHODS. *Acta Physiol Acad Sci Hung* 1964;25:203-15.
- 2461 *Balint P, Forgacs I. [RECENT VIEWS ON THE EVALUATION OF CLEARANCE METHODS.]. *Kiserl Orvostud* 1964;16:101-11.
- 1002 *Bandaranayake N, Ankrah-Tetteh T, Wijeratne S, Swaminathan R. Intra-individual variation in creatinine and cystatin C. *Clin Chem Lab Med* 2007;45:1237-9.
- 221 *Baracksky D, Jarjoura D, Cugino A, Blend D, Rutecki GW, Whittier FC. Geriatric renal function: estimating glomerular filtration in an ambulatory elderly population. *Clin Nephrol* 1997;47:222-8.
- 1004 *Barai S, Bandopadhyaya GP, Patel CD, Rathi M, Kumar R, Bhowmik D, et al. Do healthy potential kidney donors in india have an average glomerular filtration rate of 81.4 ml/min? *Nephron Physiol* 2005;101:p21-6.
- 1006 *Barai S, Gambhir S, Prasad N, Sharma RK, Ora M, Kumar A, et al. Levels of GFR and protein-induced hyperfiltration in kidney donors: a single-center experience in India. *Am J Kidney Dis* 2008;51:407-14.
- 4202 *Barbour GL, Crumb CK, Boyd CM, Reeves RD, Rastogi SP, Patterson RM. Comparison of inulin, iothalamate, and 99mTc-DTPA for measurement of glomerular filtration rate. *J Nucl Med* 1976;17:317-20.
- 13724 *Barnard HF, Bassir O, Hough JM. Fall in the inulin clearance following a single injection. *Q J Exp Physiol Cogn Med Sci* 1955;40:217-24.
- 1733 *Barri YM, Ramos EL, Balagtas RS, Peterson JC, Karlix JL. Cimetidine or ranitidine in renal transplant patients receiving cyclosporine. *Clin Transplant* 1996;10:34-8.
- 4516 *Bauer JH, Brooks CS. The long-term effect of propranolol therapy on renal function. *Am J Med* 1979;66:405-10.

4517 *Bauer JH, Brooks CS. Comparative effects of ticrynafen and hydrochlorothiazide on blood pressure, renal function, serum uric acid and electrolytes, and body fluid spaces in hypertensive man. *Nephron* 1979;23 Suppl 1:57-63.

1011 *Bauer JH, Brooks CS, Burch RN. Clinical appraisal of creatinine clearance as a measurement of glomerular filtration rate. *Am J Kidney Dis* 1982;2:337-46.

1737 *Bauer JH, Brooks CS, Burch RN. Renal function studies in man with advanced renal insufficiency. *Am J Kidney Dis* 1982;2:30-5.

1740 *Bauer R, Walter B, Zwiener U. Comparison between inulin clearance and endogenous creatinine clearance in newborn normal weight and growth restricted newborn piglets. *Exp Toxicol Pathol* 2000;52:367-72.

38 *Bauman LA, Watson NE, Jr., Scuderi PE, Peters MA. Transcutaneous renal function monitor: precision during unsteady hemodynamics. *J Clin Monit Comput* 1998;14:275-82.

1745 *Bello E, Caramelo C. [Increase of tubular secretion of creatinine by L-arginine: mechanism of practical significance in the assessment of renal function based on creatinine clearance]. *Nefrologia* 2000;20:517-22.

1753 *Berg U. Evaluation of the formula clearance as a measure of the glomerular filtration rate in cyclosporin-treated children following renal transplantation. *Transpl Int* 1991;4:72-6.

13392 *Berg UB, Back R, Celsi G, Halling SE, Homberg I, Krmar RT, et al. Comparison of Plasma Clearance of Iohexol and Urinary Clearance of Inulin for Measurement of GFR in Children. *Am J Kidney Dis* 2011; 57: 55-61.

13640 *Berg UB, Ericzon BG, Nemeth A, McDiarmid SV. Renal function in pediatric liver transplant recipients. *Transplantation* 2001;72:561-562.

2484 *Berlyne GM, Varley H, Nilwarangkur S, Hoerni M. ENDOGENOUS-CREATININE CLEARANCE AND GLOMERULAR-FILTRATION RATE. *Lancet* 1964;2:874-6.

1759 *Bertoli M, Luisetto G, Ruffatti A, Urso M, Romagnoli G. Renal function during calcitriol therapy in chronic renal failure. *Clin Nephrol* 1990;33:98-102.

4599 *Bianchi C, Donadio C, Tramonti G, Lorusso P, Norpoth M, Sibilia G, et al. A reappraisal of the bladder cumulative method as a reliable technique for the measurement of glomerular filtration rate. *Ren Fail* 1998;20:257-65.

4 *Biglarnia AR, Wadstrom J, Larsson A. Decentralized glomerular filtration rate (GFR) estimates in healthy kidney donors show poor correlation and demonstrate the need for improvement in quality and standardization of GFR measurements in Sweden. *Scand J Clin Lab Invest* 2007;67:227-35.

39 *Bird NJ, Peters C, Michell AR, Peters AM. Extracellular distribution volumes of hydrophilic solutes used to measure the glomerular filtration rate: comparison between chromium-51-EDTA and iohexol. *Physiol Meas* 2007;28:223-34.

378 *Bird NJ, Peters C, Michell AR, Peters AM. Reproducibilities and responses to food intake of GFR measured with chromium-51-EDTA and iohexol simultaneously and independently in normal subjects. *Nephrol Dial Transplant* 2008;23:1902-9.

1022 *Bird NJ, Peters C, Michell AR, Peters AM. Using the slope-only technique and estimated glomerular filtration rate for checking the reliability of slope-intercept measurement of glomerular filtration rate. *Nucl Med Commun* 2008;29:1086-92.

1023 *Bird NJ, Peters C, Michell AR, Peters AM. Suitability of a simplified technique based on iohexol for decentralized measurement of glomerular filtration rate. *Scand J Urol Nephrol* 2008;42:472-80.

13473 *Blachar Y, Eliahou EH, Iana A, Boichis H. Glomerular filtration rate (GFR) in children with vesico-urethral (V-U) reflux: Cr(51) EDTA clearance (C) versus inulin and endogenous creatinine clearance (ECC). *Pediatric Research* 1978;12:no.1052.

- 1028 *Blake GM, Gardiner N, Gnanasegaran G, Dizdarevic S. Reference ranges for ⁵¹Cr-EDTA measurements of glomerular filtration rate in children. *Nucl Med Commun* 2005;26:983-7.
- 13479 *Bochner F, Hooper WD, Sutherland JM. The renal handling of diphenylhydantoin and 5 (p hydroxyphenyl) 5 phenylhydantoin. *Clinical pharmacology and therapeutics* 1973;14:791-796.
- 1764 *Bochud M, Elston RC, Maillard M, Bovet P, Schild L, Shamlaye C, et al. Heritability of renal function in hypertensive families of African descent in the Seychelles (Indian Ocean). *Kidney Int* 2005;67:61-9.
- 1766 *Bock HA, Bachofen M, Landmann J, Thiel G. Glomerular hyperfiltration after unilateral nephrectomy in living kidney donors. *Transpl Int* 1992;5 Suppl 1:S156-9.
- 4639 *Bodem G, Grieser H, Eichelbaum M, Gugler R. Pharmacokinetics of practolol in renal failure. *Eur J Clin Pharmacol* 1974;7:249-52.
- 1767 *Bohle A, Mackensen-Haen S, von Gise H, Grund KE, Wehrmann M, Batz C, et al. The consequences of tubulo-interstitial changes for renal function in glomerulopathies. A morphometric and cytological analysis. *Pathol Res Pract* 1990;186:135-44.
- 1034 *Bokenkamp A, Dieterich C, Schumann G. Pediatric reference values for cystatin C revisited. *Pediatr Nephrol* 1999;13:367-8.
- 3231 *Bokenkamp A, Domanetzki M, Zinck R, Schumann G, Brodehl J. Reference values for cystatin C serum concentrations in children. *Pediatr Nephrol* 1998;12:125-9.
- 12907 *Bokenkamp A, Laarman CARC, Braam KI, Van Wijk JAE, Kors WA, Kool M, et al. Effect of corticosteroid therapy on low-molecular-weight protein markers of kidney function [5]. *Clinical Chemistry* 2007;53:2219-2221.
- 225 *Bollerup AC, Hesse B, Steiness E. Renal handling of iodamide and diatrizoate. Evidence of active tubular secretion of iodamide. *Eur J Clin Pharmacol* 1975;9:63-7.
- 227 *Boner G, Shelp WD, Newton M, Rieselbach RE. Factors influencing the increase in glomerular filtration rate in the remaining kidney of transplant donors. *Am J Med* 1973;55:169-74.
- 383 *Bosch JP, Saccaggi A, Lauer A, Ronco C, Belledonne M, Glabman S. Renal functional reserve in humans. Effect of protein intake on glomerular filtration rate. *Am J Med* 1983;75:943-50.
- 13638 *Boyer O, Le Bidois J, Dechaux M, Gubler MC, Niaudet P. Improvement of renal function in pediatric heart transplant recipients treated with low-dose calcineurin inhibitor and mycophenolate mofetil. *Transplantation* 2005;79:1405-1410.
- 387 *Brandle E, Melzer H, Gomez-Anson B, Flohr P, Kleinschmidt K, Sieberth HG, et al. [Can examination of spontaneous urine samples adequately replace 24-hour-urine samples for determining excretory rate of various lithogenic and inhibitory substances in metabolic evaluation of kidney calculi patients?]. *Urologe A* 1996;35:136-41.
- 1045 *Brandt JR, Wong CS, Hanrahan JD, Qualls C, McAfee N, Watkins SL. Estimating absolute glomerular filtration rate in children. *Pediatr Nephrol* 2006;21:1865-72.
- 1774 *Branten AJ, Vervoort G, Wetzels JF. Serum creatinine is a poor marker of GFR in nephrotic syndrome. *Nephrol Dial Transplant* 2005;20:707-11.
- 13624 *Brater DC. Measurement of renal function during drug development. *British Journal of Clinical Pharmacology* 2002;54:87-95.
- 13383 *Brauner L, Westling H. On the necessity of strict bed rest during determination of ⁵¹Cr-EDTA clearance. *Clinical Physiology* 1981:175-180.
- 4232 *Brien TG. Measurement of glomerular filtration-rate. *Lancet* 1968;2:1248.

- 13714 *Broberger U. Determination of glomerular filtration rate in the newborn. Comparison between results obtained by the single injection technique without collection of urine and the standard clearance technique. *Acta Paediatr Scand* 1973;62:625-9.
- 13713 *Broberger U, Aperia A. Renal function in idiopathic respiratory distress syndrome. *Acta Paediatr Scand* 1978;67:313-9.
- 13646 *Brochner-Mortensen J. Current status on assessment and measurement of glomerular filtration rate. *Clinical Physiology* 1985;5:1-17.
- 102 *Brochner-Mortensen J, Giese J, Rossing N. Renal inulin clearance versus total plasma clearance of ⁵¹Cr-EDTA. *Scand J Clin Lab Invest* 1969;23:301-5.
- 4719 *Brochner-Mortensen J, Hammerich B, Christoffersen J. Assessment of renal function from plasma urea and plasma creatinine in children. *Scand J Urol Nephrol* 1982;16:229-36.
- 1787 *Brown J, Corr L. Renal mechanisms of human alpha-atrial natriuretic peptide in man. *J Physiol* 1987;387:31-46.
- 174 *Brown SC, O'Reilly PH. Iohexol clearance for the determination of glomerular filtration rate in clinical practice: evidence for a new gold standard. *J Urol* 1991;146:675-9.
- 103 *Brucker M, Caillard S, Moulin B. [Methods of assessment of the renal function in kidney transplant patients]. *Nephrol Ther* 2008;4 Suppl 1:S40-S44.
- 2525 *Brunner H. [Effects of acetazolamide, chlorothiazide and chlormerodrin on inulin clearance, true endogenous creatinine, para-aminohippuric acid and urea. Differentiation of nephrotoxic effects of diuretics from dehydration sequelae in healthy rats.]. *Naunyn Schmiedebergs Arch Exp Pathol Pharmacol* 1959;236:559-81.
- 1790 *Bunke M, Sloan R, Brier M, Ganzel B. An improved glomerular filtration rate in cardiac transplant recipients with once-a-day cyclosporine dosing. *Transplantation* 1995;59:537-40.
- 105 *Burke JT, Brault Y, Kahan BD, Hricik DE, Grinyo JM, Chapman JR, et al. Accuracy and variability of equations to estimate glomerular filtration rates in renal transplant patients receiving sirolimus and/or calcineurin inhibitor immunosuppression. *Transpl Int* 2008;21:434-40.
- 1793 *Burkhardt H, Bojarsky G, Gretz N, Gladisch R. Creatinine clearance, Cockcroft-Gault formula and cystatin C: estimators of true glomerular filtration rate in the elderly? *Gerontology* 2002;48:140-6.
- 1797 *Burkhardt H, Hahn T, Gretz N, Gladisch R. Bedside estimation of the glomerular filtration rate in hospitalized elderly patients. *Nephron Clin Pract* 2005;101:c1-8.
- 13474 *Burland WL, Gleadle RI, Mills JG. The effect of cimetidine on renal function Cimetidine: Proceedings of the Second International Symposium on Histamine H(2)-Receptor Antagonists. SOURCE EXCERPTA MEDICA,AMSTERDAM - ICS (1977) No. 416 (67-73). Date of Publication: 1977.
- 402 *Buzio C, Mutti A, Capani F, Andrulli S, Perazzoli F, Alinovi R, et al. Circadian rhythm of proteinuria: effects of an evening meat meal. *Nephrol Dial Transplant* 1989;4:266-70.
- 403 *Buzio C, Mutti A, Perazzoli F, Alinovi R, Arisi L, Negro A. Protein-induced changes in kidney function depend on the time of administration but not on the dietary source. *Nephron* 1990;56:234-40.
- 404 *Buzio C, Quaretti P, Arisi L, Negro A, Barani R, Alquati MG. Time-dependent increase of glomerular filtration rate after oral protein load: evidence for diurnal variations in renal response. *Scand J Clin Lab Invest* 1988;48:15-9.
- 1800 *Campens D, Buntinx F. Selecting the best renal function tests. A meta-analysis of diagnostic studies. *Int J Technol Assess Health Care* 1997;13:343-56.
- 231 *Cangiano JL, Genuth SM, Renerts L, Berman LB. Simplified measurement of glomerular filtration rate. *Invest Urol* 1971;9:34-8.

- 1804 *Caregaro L, Menon F, Angeli P, Amodio P, Merkel C, Bortoluzzi A, et al. Limitations of serum creatinine level and creatinine clearance as filtration markers in cirrhosis. *Arch Intern Med* 1994;154:201-5.
- 408 *Carola F. Renal ptosis and change of glomerular filtration values in the upright and supine positions. *Clin Nucl Med* 2003;28:150-3.
- 4247 *Carrie BJ, Golbetz HV, Michaels AS, Myers BD. Creatinine: an inadequate filtration marker in glomerular diseases. *Am J Med* 1980;69:177-82.
- 411 *Castenfors J. Renal function during exercise. With special reference to exercise proteinuria and the release of renin. *Acta Physiol Scand Suppl* 1967;293:1-44.
- 412 *Castenfors J, Piscator M. Renal haemodynamics, urine flow and urinary protein excretion during exercise in supine position at different loads. *Acta Med Scand Suppl* 1967;472:231-44.
- 3265 *Cataldi L, Mussap M, Bertelli L, Ruzzante N, Fanos V, Plebani M. Cystatin C in healthy women at term pregnancy and in their infant newborns: relationship between maternal and neonatal serum levels and reference values. *Am J Perinatol* 1999;16:287-95.
- 40 *Chachati A, Meyers A, Godon JP, Rigo P. Rapid method for the measurement of differential renal function: validation. *J Nucl Med* 1987;28:829-36.
- 41 *Chachati A, Meyers A, Rigo P, Godon JP. Validation of a simple isotopic technique for the measurement of global and separated renal function. *Uremia Invest* 1985;9:177-9.
- 12925 *Chakrabarti S, Chattopadhyay I. Measuring renal function in old age. *Reviews in Clinical Gerontology* 2008;18:257-267.
- 13641 *Chan C, Maurer J, Cardella C, Cattran D, Pei Y. A randomized controlled trial of verapamil on cyclosporine nephrotoxicity in heart and lung transplant recipients. *Transplantation* 1997;63:1435-1440.
- 106 *Chantler C, Garnett ES, Parsons V, Veall N. Glomerular filtration rate measurement in man by the single injection methods using 51Cr-EDTA. *Clin Sci* 1969;37:169-80.
- 1807 *Chantrel F, Agin A, Offner M, Koehl C, Moulin B, Hannedouche T. Comparison of cystatin C versus creatinine for detection of mild renal failure. *Clin Nephrol* 2000;54:374-81.
- 12928 *Chaudhry O, Mayo MM, Mooradian AD. Undetectable plasma creatinine in a paralyzed patient infected with human immunodeficiency virus. *Laboratory Medicine* 2005;36:479-481.
- 4246 *Chennavasin P, Brater DC. Nomograms for drug use in renal disease. *Clin Pharmacokinet* 1981;6:193-214.
- 13634 *Cherney DZI, Scholey JW, Miller JA. Insights into the regulation of renal hemodynamic function in diabetic mellitus. *Current Diabetes Reviews* 2008;4:280-290.
- 42 *Chervu LR, Blafox MD. Renal radiopharmaceuticals--an update. *Semin Nucl Med* 1982;12:224-45.
- 233 *Chervu LR, Freeman LM, Blafox MD. Radiopharmaceuticals for renal studies. *Semin Nucl Med* 1974;4:3-22.
- 1813 *Chollet-Dallon E, Stoermann-Chopard C, Martin PY. [Could cystatine C replace creatinine as a marker of glomerular filtration rate?]. *Rev Med Suisse* 2006;2:582-5.
- 4911 *Cholongitas E, Shusang V, Marelli L, Nair D, Thomas M, Patch D, et al. Review article: renal function assessment in cirrhosis - difficulties and alternative measurements. *Aliment Pharmacol Ther* 2007;26:969-78.
- 1816 *Cianciaruso B, Bellizzi V, Capuano A, Bovi G, Nastasi A, Conte G, et al. Short-term effects of low protein-normal sodium diet on renal function in chronic renal failure. *Kidney Int* 1994;45:852-60.
- 234 *Cimmino CJ. Benefits of direct glomerular filtration rate (GFR) determination versus creatinine-based tests for evaluating renal function. *J Am Osteopath Assoc* 1998;98:437-44.

- 1077 *Claeys V, Nicolet G, Guignard JP. [Nephrological tests: normal values]. *Rev Med Suisse Romande* 1985;105:959-68.
- 13626 *Clarkson JC, Smidt KP. Measurement of glomerular filtration rate in neonates using the single injection clearance of (113)mIn DTPA. *Australian Paediatric Journal* 1973;9:195-199.
- 235 *Cohen ML. Radionuclide clearance techniques. *Semin Nucl Med* 1974;4:23-38.
- 236 *Cohen ML, Smith FG, Jr., Mindell RS, Vernier RL. A simple, reliable method of measuring glomerular filtration rate using single, low dose sodium iothalamate I-131. *Pediatrics* 1969;43:407-15.
- 13439 *Cohney S, Kanellis J, Howell M. Donor renal function. *Nephrology* 2010;15:S137-S145.
- 1824 *Conil JM, Georges B, Fourcade O, Seguin T, Lavit M, Samii K, et al. Assessment of renal function in clinical practice at the bedside of burn patients. *Br J Clin Pharmacol* 2007;63:583-94.
- 4243 *Coresh J, Toto RD, Kirk KA, Whelton PK, Massry S, Jones C, et al. Creatinine clearance as a measure of GFR in screenees for the African-American Study of Kidney Disease and Hypertension pilot study. *Am J Kidney Dis* 1998;32:32-42.
- 1087 *Couchoud C, Pozet N, Labeeuw M, Pouteil-Noble C. Screening early renal failure: cut-off values for serum creatinine as an indicator of renal impairment. *Kidney Int* 1999;55:1878-84.
- 13708 *Coulthard MG. Comparison of methods of measuring renal function in preterm babies using inulin. *J Pediatr* 1983;102:923-30.
- 43 *Cousins C, Gunasekera RD, Mubashar M, Mohammadtaghi S, Strong R, Myers MJ, et al. Comparative kinetics of microvascular inulin and 99mTc-labelled diethylenetriaminepenta-acetic acid exchange. *Clin Sci (Lond)* 1997;93:471-7.
- 44 *Cousins C, Mohammadtaghi S, Mubashar M, Strong R, Gunasekera RD, Myers MJ, et al. Clearance kinetics of solutes used to measure glomerular filtration rate. *Nucl Med Commun* 1999;20:1047-54.
- 45 *Cousins C, Skehan SJ, Rolph SM, Flaxman ME, Ballinger JR, Bird NJ, et al. Comparative microvascular exchange kinetics of [(77)Br]bromide and (99m)Tc-DTPA in humans. *Eur J Nucl Med Mol Imaging* 2002;29:655-62.
- 2555 *Crawford B. Depression of the exogenous creatinine/inulin or thiosulfate clearance ratios in man by diodrast and p-aminohippuric acid. *J Clin Invest* 1948;27:171-5.
- 452 *Curtis G, Fogel M. Creatinine excretion: diurnal variation and variability of whole and part-day measures. A methodologic issue in psychoendocrine research. *Psychosom Med* 1970;32:337-50.
- 13628 *Dalton RN, Chetty R, Stuart M, Iacona RB, Swaisland A. Effects of the Src inhibitor saracatinib (AZD0530) on renal function in healthy subjects. *Anticancer Research* 2010;30:2935-2942.
- 1832 *D'Amico G, Remuzzi G, Maschio G, Gentile MG, Gotti E, Oldrizzi L, et al. Effect of dietary proteins and lipids in patients with membranous nephropathy and nephrotic syndrome. *Clin Nephrol* 1991;35:237-42.
- 1833 *Daniel JP, Chantrel F, Offner M, Moulin B, Hannedouche T. Comparison of cystatin C, creatinine and creatinine clearance vs. GFR for detection of renal failure in renal transplant patients. *Ren Fail* 2004;26:253-7.
- 455 *Dann EJ, Gillis S, Burstein R. Effect of fluid intake on renal function during exercise in the cold. *Eur J Appl Physiol Occup Physiol* 1990;61:133-7.
- 13476 *Davison JM, Hytten FE. Glomerular filtration during and after pregnancy. *J.Obstet.Gynaec.Brit.Cwlth.* 1974;81:588-595.
- 13527 *Davison JM, Lindheimer MD. Renal disease in pregnant women. *Clinical Obstetrics and Gynecology* 1978;21:411-427.

- 13639 *De Rosa ML, Cardace P, Rossi M, Baiano A, De Cristofaro A. Evaluation of long-term efficacy and tolerability of irbesartan in elderly hypertensive patients with renal impairment in an open-label study. *Current Therapeutic Research - Clinical and Experimental* 2002;63:201-215.
- 1839 *De Rosa ML, de Cristofaro A, Rossi M, Baiano A, Cardace P, Albanese L, et al. Irbesartan effects on renal function in patients with renal impairment and hypertension: a drug-withdrawal study. *J Cardiovasc Pharmacol* 2001;38:482-9.
- 46 *De Santo NG, Anastasio P, Cirillo M, Santoro D, Spitali L, Mansi L, et al. Measurement of glomerular filtration rate by the ^{99m}Tc-DTPA renogram is less precise than measured and predicted creatinine clearance. *Nephron* 1999;81:136-40.
- 13643 *De Santo NG, Anastasio P, Cirillo M, Spitali L, Capazzo G, Santoro D. Sequential analysis of variation in glomerular filtration rate to calculate the haemodynamic response to a meat meal. *Nephrology Dialysis Transplantation* 1995;10:1629-1636.
- 1105 *De Santo NG, Anastasio P, Spitali L, Cirillo M, Santoro D, Pollastro RM, et al. The renal hemodynamic response to an oral protein load is normal in IgA nephropathy. *Nephron* 1997;76:406-10.
- 13454 *De Santo NG, Anastasio P, Spitali L, Del Vecchio Bianco C, Carvinelli M, Cirillo M. Creatinine clearance: An inadequate marker of renal filtration in patients with early posthepatic cirrhosis (Child A) without fluid retention and muscle wasting. *Nephron* 1995;70:421-424.
- 107 *Deckart H, Herzmann H. [Nephrographic evaluation of glomerular filtration rate using I-131-inulin, I-131-diatrizoate and Co58-EDTA respectively]. *Strahlentherapie Sonderb* 1968;67:406-13.
- 47 *Deckart H, Herzmann H, Roick H. [EDTA and DTPA labelled with Cr-51 and Co-58 in renal diagnosis]. *Med Klin* 1969;64:1858-61.
- 1847 *Dehne MG, Muhling J, Papke G, Nopens H, Kuntzsch U, Hempelmann G. Unrecognized renal damage in critically ill patients. *Ren Fail* 1999;21:695-706.
- 1112 *Delanaye P, Cavalier E, Depas G, Chapelle JP, Krzesinski JM. New data on the intraindividual variation of cystatin C. *Nephron Clin Pract* 2008;108:c246-8.
- 177 *Delanaye P, Chapelle JP, Ferir AM, Gielen J, Krzesinski JM, Rorive G. [Evaluation of glomerular filtration rate in clinical practice]. *Rev Med Liege* 2003;58:95-100.
- 13651 *Delanaye P, Scheen AJ. Effects of fibrates on cardiovascular outcomes. *The Lancet* 2010;376:1051.
- 2 *Delanghe JR. How to estimate GFR in children. *Nephrol Dial Transplant* 2009;24:714-6.
- 1854 *DeSanto NG, Anastasio P, Coppola S, Barba G, Jadanza A, Capasso G. Age-related changes in renal reserve and renal tubular function in healthy humans. *Child Nephrol Urol* 1991;11:33-40.
- 1855 *DeSanto NG, Anastasio P, Loguercio C, Spitali L, Del Vecchio Bianco C, Corvinelli M, et al. Creatinine clearance: an inadequate marker of renal filtration in patients with early posthepatic cirrhosis (Child A) without fluid retention and muscle wasting. *Nephron* 1995;70:421-4.
- 1856 *DeSanto NG, Coppola S, Anastasio P, Coscarella G, Capasso G, Bellini L, et al. Predicted creatinine clearance to assess glomerular filtration rate in chronic renal disease in humans. *Am J Nephrol* 1991;11:181-5.
- 13361 *DeSanto NG, Coppola S, Anastasio P, Coscarella G, Capasso G, Santangelo R, et al. Lithium clearance in patients with chronic renal failure. *Miner Electrolyte Metab* 1991;17:166-72.
- 13636 *Diskin CJ. Creatinine and GFR: An imperfect marriage of convenience [6]. *Nephrology Dialysis Transplantation* 2006;21:3338-3339.
- 239 *Dittrich P, Riccabona G. [Clearance examinations with ¹³¹I iothalamate]. *Wildl Dis* 1968;49:401-5.

- 110 *Ditzel J, Vestergaard P, Brinklov M. [Glomerular filtration rate determined using ⁵¹Cr-EDTA. A practical method based upon the plasma disappearance curve for 4 plasma samples]. *Nord Med* 1971;85:650-3.
- 111 *Ditzel J, Vestergaard P, Brinklov M. Glomerular filtration rate determined by ⁵¹Cr-EDTA-complex. A practical method based upon the plasma disappearance curve determined from four plasma samples. *Scand J Urol Nephrol* 1972;6:166-70.
- 468 *Dodge WF, Travis LB, Daeschner CW. Comparison of endogenous creatinine clearance with inulin clearance. *Am J Dis Child* 1967;113:683-92.
- 48 *Donadio C, Consani C, Ardini M, Caprio F, Grassi G, Lucchesi A. Prediction of glomerular filtration rate from body cell mass and plasma creatinine. *Curr Drug Discov Technol* 2004;1:221-8.
- 13540 *Donath A. The simultaneous determination in children of glomerular filtration rate and effective renal plasma flow by the single injection clearance technique. *Acta paediatrica Scandinavica* 1971;60:512-520.
- 1860 *Donckerwolcke RM, Coppes MJ. Adaptation of renal function after unilateral nephrectomy in children with renal tumors. *Pediatr Nephrol* 2001;16:568-74.
- 2583 *Doxiadis SA, Goldfinch MK. Comparison of inulin and endogenous creatinine clearance in young children. *J Physiol* 1952;118:454-60.
- 13461 *Dratwa M, Burette A, Van Gossum M, Collart F, Wens R, Charlier L, et al. No rise in glomerular filtration rate after protein loading in cirrhotics. *Kidney international* 1987;32:S-32-S-34.
- 4206 *Drukker W. Measuring GFR. *Int J Artif Organs* 1987;10:129-30.
- 50 *Durbin PW, Schmidt CT. Predicting the kinetics of chelating agents in man from animal data. *Health Phys* 1989;57 Suppl 1:165-73; discussion 174.
- 1867 *Eisenhauer T, Jungmann E, Warneboldt D, Ansorge G, Scherberich J, Talartschik J. [Behavior of the renal functional reserve in type I diabetic patients: effect of ACE-inhibition]. *Klin Wochenschr* 1990;68:750-7.
- 13627 *Eisenhauer T, Talartschik J, Scheler F. Angiotensin converting enzyme inhibition by captopril reduces amino acid-mediated increase in glomerular filtration rate. In: *Journal of Hypertension*; 1986. p S258-s260.
- 2592 *Ek J, Josephson B. On the influence of beer and a purine derivative on the renal clearance of creatinine, inulin and para-aminohippuric acid. *Acta Physiol Scand* 1953;28:347-54.
- 240 *Elwood CM, Sigman EM. The measurement of glomerular filtration rate and effective renal plasma flow in man by iothalamate ¹²⁵I and iodopyracet ¹³¹I. *Circulation* 1967;36:441-8.
- 241 *Elwood CM, Sigman EM, Treger C. The measurement of glomerular filtration rate with ¹²⁵I-sodium iothalamate (Conray). *Br J Radiol* 1967;40:581-3.
- 1872 *Emara M, Zahran A, Abd El Hady H, Shoker A. How to best define patients with moderate chronic kidney disease. *Nephron Clin Pract* 2008;110:c195-206.
- 1875 *Englund MS, Berg UB, Arfwidson K. Renal functional reserve in transplanted and native single kidneys of children and adults. *Pediatr Nephrol* 1997;11:312-7.
- 1878 *Epstein M, Parving HH, Ruilope LM. Surrogate endpoints and renal protection: focus on microalbuminuria. *Blood Press Suppl* 1997;2:52-7.
- 113 *Erd W, Fritzer W, Hofer R, Oei HJ, Pfeiffer G. [Comparative studies on renal clearance using inulin, PAH, Cr-⁵¹-EDTA and I-¹²⁵-hippuran]. *Wien Z Inn Med* 1969;50:106-11.
- 178 *Erley CM, Bader BD, Berger ED, Vochazer A, Jorzik JJ, Dietz K, et al. Plasma clearance of iodine contrast media as a measure of glomerular filtration rate in critically ill patients. *Crit Care Med* 2001;29:1544-50.

- 13642 *Erley CM, Klass M, Kramer D, Berger E, Heyne N, Braun N, et al. Randomized controlled trial of ACE-inhibitors and (beta)-blockers with and without vasodilating activity in chronic glomerulonephritis. *International journal of clinical pharmacology and therapeutics* 1996;34:504-509.
- 1158 *Esposito C, Plati A, Mazzullo T, Fasoli G, De Mauri A, Grosjean F, et al. Renal function and functional reserve in healthy elderly individuals. *J Nephrol* 2007;20:617-25.
- 13468 *Falkheden T, Ohlsson L, Sjogren B. Renal function in primary hyperparathyroidism. A follow-up study two to eleven years after surgery comprising 139 patients. *Scandinavian Journal of Urology and Nephrology* 1980;14:167-175.
- 3354 *Fanos V, Mussap M, Plebani M, Cataldi L. Cystatin C in paediatric nephrology. Present situation and prospects. *Minerva Pediatr* 1999;51:167-77.
- 1885 *Farker K, Merkel U, Schweer H, Haerting J, Madani SF, Eggers R, et al. Effects of short-term treatment with diclofenac-colestyramine on renal function and urinary prostanoid excretion in patients with type-2 diabetes. *Eur J Clin Pharmacol* 2002;58:85-91.
- 1887 *Farker K, Nassr N, Huck F, Zerle G, Rosenkranz B, Schmieder G, et al. Dipyron and diclofenac do not influence creatinine-clearance, inulin-clearance or PAH-clearance in healthy male volunteers. *Int J Clin Pharmacol Ther* 1995;33:125-30.
- 494 *Farquhar WB, Morgan AL, Zambraski EJ, Kenney WL. Effects of acetaminophen and ibuprofen on renal function in the stressed kidney. *J Appl Physiol* 1999;86:598-604.
- 13712 *Fawer CL, Torrado A, Guignard JP. Single injection clearance in the neonate. *Biol Neonate* 1979;35:321-4.
- 114 *Favre H. Critical study of the value of renal clearances measured by the single shot technic. *Contrib Nephrol* 1978;11:19-21.
- 4233 *Favre HR, Wing AJ. Simultaneous ⁵¹Cr edetic acid, inulin, and endogenous creatinine clearances in 20 patients with renal disease. *Br Med J* 1968;1:84-6.
- 1167 *Fehrman-Ekholm I, Skeppholm L. Renal function in the elderly (>70 years old) measured by means of iohexol clearance, serum creatinine, serum urea and estimated clearance. *Scand J Urol Nephrol* 2004;38:73-7.
- 12963 *Filler G. How to measure renal function in children - What is the role of cystatin C? *Current Pediatric Reviews* 2006;2:225-231.
- 51 *Filler G, Sharma AP. How to monitor renal function in pediatric solid organ transplant recipients. *Pediatr Transplant* 2008;12:393-401.
- 1170 *Filler G, Witt I, Priem F, Ehrich JH, Jung K. Are cystatin C and beta 2-microglobulin better markers than serum creatinine for prediction of a normal glomerular filtration rate in pediatric subjects? *Clin Chem* 1997;43:1077-8.
- 2620 *Finco DR, Groves C. Mechanism of renal excretion of creatinine by the pony. *Am J Vet Res* 1985;46:1625-8.
- 499 *Finkelstein J, Joshi A, Hise MK. Association of physical activity and renal function in subjects with and without metabolic syndrome: a review of the Third National Health and Nutrition Examination Survey (NHANES III). *Am J Kidney Dis* 2006;48:372-82.
- 1172 *Finney H, Newman DJ, Price CP. Adult reference ranges for serum cystatin C, creatinine and predicted creatinine clearance. *Ann Clin Biochem* 2000;37 (Pt 1):49-59.
- 1173 *Finney H, Newman DJ, Thakkar H, Fell JM, Price CP. Reference ranges for plasma cystatin C and creatinine measurements in premature infants, neonates, and older children. *Arch Dis Child* 2000;82:71-5.

- 3369 *Fischbach M, Graff V, Terzic J, Bergere V, Oudet M, Hamel G. Impact of age on reference values for serum concentration of cystatin C in children. *Pediatr Nephrol* 2002;17:104-6.
- 1897 *Fliser D, Bischoff I, Hanses A, Block S, Joest M, Ritz E, et al. Renal handling of drugs in the healthy elderly. Creatinine clearance underestimates renal function and pharmacokinetics remain virtually unchanged. *Eur J Clin Pharmacol* 1999;55:205-11.
- 13701 *Florijn KW, Barendregt JN, Lentjes EG, van Dam W, Prodjosudjadi W, van Saase JL, et al. Glomerular filtration rate measurement by "single-shot" injection of inulin. *Kidney Int* 1994;46:252-9.
- 116 *Fogh-Andersen N. Binding of the chromium-ethylenediaminetetraacetic acid complex (CrEDTA) to human albumin. *Scand J Clin Lab Invest* 1980;40:805-8.
- 52 *Fotopoulos A, Bokharhli JA, Tsiouris S, Katsaraki A, Papadopoulos A, Tsironi M, et al. [Comparison of six radionuclidic and non-radionuclidic methods for the assessment of glomerular filtration rate in patients with chronic renal failure]. *Hell J Nucl Med* 2006;9:133-40.
- 13438 *Francoz C, Glotz D, Moreau R, Durand F. The evaluation of renal function and disease in patients with cirrhosis. *Journal of Hepatology* 2010;52:605-613.
- 13631 *Frank H, Graf J, Amann-Gassner U, Bratke R, Daniel H, Heemann U, et al. Effect of short-term high-protein compared with normal-protein diets on renal hemodynamics and associated variables in healthy young men. *American Journal of Clinical Nutrition* 2009;90:1509-1516.
- 53 *Frennby B, Sterner G. Contrast media as markers of GFR. *Eur Radiol* 2002;12:475-84.
- 181 *Frennby B, Sterner G, Almen T, Chai CM, Jonsson BA, Mansson S. Extrarenal plasma clearance of iohexol, chromium-51-ethylenediaminetetraacetic acid, and inulin in anephric pigs. *Acad Radiol* 1996;3:145-53.
- 13522 *Frennby B, Sterner G, Almen T, Hagstam KE, Jacobsson L. Determination of low glomerular filtration rate using iohexol clearance. *Investigative Radiology* 1994;29:S234-S235.
- 1910 *Freysz M, Lafleur P, Dupont G, Guillard JC, d'Athis P, Escousse A, et al. Comparison of creatinine and inulin clearances in multiple trauma. *Biomed Pharmacother* 1990;44:175-80.
- 54 *Funck-Brentano JL, Leski M. [Measurement of the indirect clearance of DTPA 140 La without taking urine in edematous patients. Preliminary note]. *J Urol Nephrol (Paris)* 1969;75:933-9.
- 3391 *Galteau MM, Guyon M, Gueguen R, Siest G. Determination of serum cystatin C: biological variation and reference values. *Clin Chem Lab Med* 2001;39:850-7.
- 526 *Gambaro G, Verlato F, Budakovic A, Casara D, Saladini G, Del Prete D, et al. Renal impairment in chronic cigarette smokers. *J Am Soc Nephrol* 1998;9:562-7.
- 4234 *Garnett ES, Parsons V, Veall N. Measurement of glomerular filtration-rate in man using a 51Cr-edetic-acid complex. *Lancet* 1967;1:818-9.
- 249 *Gaspar Hernandez J, Vazquez Torres G, Herrera Acosta J, Pena JC. [Studies on the mechanism of 131I-iodothalamate renal excretion and its use to measure glomerular filtration (author's transl)]. *Rev Invest Clin* 1980;32:487-95.
- 182 *Gaspari F, Guerini E, Perico N, Mosconi L, Ruggenti P, Remuzzi G. Glomerular filtration rate determined from a single plasma sample after intravenous iohexol injection: is it reliable? *J Am Soc Nephrol* 1996;7:2689-93.
- 183 *Gaspari F, Perico N, Matalone M, Signorini O, Azzollini N, Mister M, et al. Precision of plasma clearance of iohexol for estimation of GFR in patients with renal disease. *J Am Soc Nephrol* 1998;9:310-3.
- 55 *Gaspari F, Perico N, Remuzzi G. Measurement of glomerular filtration rate. *Kidney Int Suppl* 1997;63:S151-4.
- 185 *Gaspari F, Perico N, Remuzzi G. Application of newer clearance techniques for the determination of glomerular filtration rate. *Curr Opin Nephrol Hypertens* 1998;7:675-80.

- 186 *Gaspari F, Perico N, Ruggenenti P, Mosconi L, Amuchastegui CS, Guerini E, et al. Plasma clearance of nonradioactive iohexol as a measure of glomerular filtration rate. *J Am Soc Nephrol* 1995;6:257-63.
- 119 *Gibb DM, Dalton NR, Barratt MT. Measurement of glomerular filtration rate in children with insulin-dependent diabetes mellitus. *Clin Chim Acta* 1989;182:131-9.
- 1919 *Giovannetti S, Barsotti G. In defense of creatinine clearance. *Nephron* 1991;59:11-4.
- 13633 *Glasscock RJ, Winearls CG. eGFR: Readjusting its rating. *Clinical Journal of the American Society of Nephrology* 2009;4:867-869.
- 121 *Granerus G, Aurell M. Reference values for 51Cr-EDTA clearance as a measure of glomerular filtration rate. *Scand J Clin Lab Invest* 1981;41:611-6.
- 122 *Grebe SF, Weise M, Berghauer H, Fangewisch GL, Temme H. [Comparison of conventional renal clearance values with differently evaluated results of radioisotope nephrography (author's transl)]. *Radiologe* 1981;21:553-61.
- 1209 *Grewal GS, Blake GM. Reference data for 51Cr-EDTA measurements of the glomerular filtration rate derived from live kidney donors. *Nucl Med Commun* 2005;26:61-5.
- 1211 *Gross JL, Friedman R, Azevedo MJ, Silveiro SP, Pecis M. Effect of age and sex on glomerular filtration rate measured by 51Cr-EDTA. *Braz J Med Biol Res* 1992;25:129-34.
- 254 *Groth T, Tengstrom B. A simple method for the estimation of glomerular filtration rate. *Scand J Clin Lab Invest* 1977;37:39-47.
- 255 *Groth T, Tengstrom B. The usefulness of 125I-sodium iothalamate as a GFR-indicator in single intravenous injection tests. *Ups J Med Sci* 1978;83:53-63.
- 1931 *Grzeszczak W, Zukowska-Szczechowska E, Zembala M, Religa Z, Pokrzywnicki W, Kochanska-Dziurawicz A. Serum beta-2-microglobulin (beta-2-M) levels in heart transplant patients (HTP). *J Cardiovasc Surg (Torino)* 1996;37:105-7.
- 256 *Guedon J, Gluckmann JC, Chapiro E, Semat P. [Inadequacy of intravenous urography for evaluation of individual kidney function in unilateral renal disease]. *Presse Med* 1970;78:1263-8.
- 13465 *Guignard JP, Tabin R, Vienny H, Genton N. Effect of trimethoprim-sulfamethoxazole on renal function. *Current Therapeutic Research - Clinical and Experimental* 1983;34:801-806.
- 1218 *Guignard JP, Torrado A, Feldman H, Gautier E. Assessment of glomerular filtration rate in children. *Helv Paediatr Acta* 1980;35:437-47.
- 56 *Gunasekera RD, Allison DJ, Peters AM. Glomerular filtration rate in relation to extracellular fluid volume: similarity between 99mTc-DTPA and inulin. *Eur J Nucl Med* 1996;23:49-54.
- 57 *Hadj Aissa A, Collard M, Pozet N, Zech P, Traeger J. Separate calculation of glomerular and tubular clearance by means of renal scintigraphy with hippuran and DTPA. *Nephron* 1985;40:401-4.
- 544 *Hadj-Aissa A, Bankir L, Fraysse M, Bichet DG, Laville M, Zech P, et al. Influence of the level of hydration on the renal response to a protein meal. *Kidney Int* 1992;42:1207-16.
- 1940 *Hadj-Aissa A, Cochat P, Dubourg L, Wright C, Pozet N. [Measurement of renal function in children]. *Arch Pediatr* 1994;1:273-80.
- 4248 *Hagstam KE, Nordenfelt I, Svensson L, Svensson SE. Comparison of different methods for determination of glomerular filtration rate in renal disease. *Scand J Clin Lab Invest* 1974;34:31-6.
- 553 *Halimi JM, Philippon C, Mimran A. Contrasting renal effects of nicotine in smokers and non-smokers. *Nephrol Dial Transplant* 1998;13:940-4.

- 13529 *Hall JE, Gutyon AC. Single injection clearances of 125 I iothalamate. Federation Proceedings 1976;35:No.1857.
- 187 *Haller M, Rohner K, Muller W, Reutter F, Binder H, Estelberger W, et al. Single-injection inulin clearance for routine measurement of glomerular filtration rate in cats. J Feline Med Surg 2003;5:175-81.
- 124 *Hamilton D, Riley P, Miola U, Mousa D, Popovich W, al Khader A. Total plasma clearance of 51Cr-EDTA: variation with age and sex in normal adults. Nucl Med Commun 2000;21:187-92.
- 259 *Harries JD, Mildenerger RR, Malowany AS, Drummond KN. A computerized cumulative integral method for the precise measurement of the glomerular filtration rate. 1. Proc Soc Exp Biol Med 1972;140:1148-55.
- 260 *Harris KM, Koike TI. Apparent volumes of distribution of 125-I-iothalamate and inulin in chickens. Proc Soc Exp Biol Med 1975;149:242-7.
- 1948 *Hatano M. [Clinical study of tubular creatinine secretion in renal dysfunction]. Nippon Jinzo Gakkai Shi 1991;33:1097-104.
- 13526 *Haycock GB. Old and new tests of renal function. Journal of Clinical Pathology 1981;34:1276-1281.
- 1232 *Heaf JG. The origin of the 1 x 73-m² body surface area normalization: problems and implications. Clin Physiol Funct Imaging 2007;27:135-7.
- 4235 *Heath DA, Knapp MS, Walker WH. Comparison between inulin and 51Cr-labelled edetic acid for the measurement of glomerular filtration-rate. Lancet 1968;2:1110-2.
- 13704 *Hedman A, Adan-Abdi Y, Alvan G, Strandvik B, Arvidsson A. Influence of the glomerular filtration rate on renal clearance of ceftazidime in cystic fibrosis. Clin Pharmacokinet 1988;15:57-65.
- 1954 *Heering P, Kutkuhn B, Kreuzpaintner G, Reinhard T, Sundmacher R, Grabensee B. [Kidney function of patients with healthy kidneys during cyclosporin treatment]. Klin Wochenschr 1991;69:880-6.
- 1957 *Hellerstein S, Alon U, Warady BA. Creatinine for estimation of glomerular filtration rate. Pediatr Nephrol 1992;6:507-11.
- 1958 *Hellerstein S, Berenbom M, Alon US, Warady BA. Creatinine clearance following cimetidine for estimation of glomerular filtration rate. Pediatr Nephrol 1998;12:49-54.
- 1961 *Henry ML, Elkhammas EA, Davies EA, Ferguson RM. A clinical trial of cyclosporine G in cadaveric renal transplantation. Pediatr Nephrol 1995;9 Suppl:S49-51.
- 1962 *Herbst CP, Dunn M, Viviers AC, van Zyl M, van Aswegen A, Otto AC. [A comparison of three different techniques for measuring glomerular filtration rate]. S Afr Med J 1990;78:266-7.
- 1963 *Herrera J, Rodriguez-Iturbe B. Stimulation of tubular secretion of creatinine in health and in conditions associated with reduced nephron mass. Evidence for a tubular functional reserve. Nephrol Dial Transplant 1998;13:623-9.
- 125 *Hilbrands LB, Artz MA, Wetzels JF, Koene RA. Cimetidine improves the reliability of creatinine as a marker of glomerular filtration. Kidney Int 1991;40:1171-6.
- 1969 *Hilbrands LB, Wetzels JF, Hoitsma AJ, Koene RA. Cyclosporin does not inhibit the tubular secretion of creatinine. Nephrol Dial Transplant 1996;11:833-6.
- 13536 *Hilson AJW, Moore AEB, Blake GM, Fogelman I. Overestimation of true renal clearance by conventional measurements of GFR using (51)Cr-EDTA (multiple letters). European Journal of Nuclear Medicine and Molecular Imaging 2003;30:934-935.
- 13457 *Hirata-Dulas CAT, Halstenson CE, Kasiske BL. Improvement in the accuracy and precision of creatinine clearance as a measure of glomerular filtration rate with oral cimetidine in renal transplant recipients. Clinical Transplantation 1993;7:552-558.

- 1244 *Hladunewich MA, Derby GC, Lafayette RA, Blouch KL, Druzin ML, Myers BD. Effect of L-arginine therapy on the glomerular injury of preeclampsia: a randomized controlled trial. *Obstet Gynecol* 2006;107:886-95.
- 261 *Holliday MA, Heilbron D, al-Uzri A, Hidayat J, Uauy R, Conley S, et al. Serial measurements of GFR in infants using the continuous iohalamate infusion technique. Southwest Pediatric Nephrology Study Group (SPNSG). *Kidney Int* 1993;43:893-8.
- 262 *Holweger K, Bokemeyer C, Lipp HP. Accurate measurement of individual glomerular filtration rate in cancer patients: an ongoing challenge. *J Cancer Res Clin Oncol* 2005;131:559-67.
- 263 *Hor G. Advantages and limitations of radioisotope clearance methods from a clinician's point of view. *J Nucl Biol Med* 1972;16:167-76.
- 1986 *Horackova M, Schuck O, Komers R, Charvat J, Teplan V, Kvapil M. Effect of rofecoxib on the glomerular filtration rate, proteinuria and the renin-angiotensin-aldosterone system in elderly subjects with chronic renal impairment. *Int J Clin Pharmacol Ther* 2005;43:413-9.
- 1988 *Horakova M, Schuck O, Komers R, Teplan V, Charvat J, Kvapil M. [Effects of specific cyclooxygenase-2 inhibition on the renal functions of elderly patients with renal function impairment]. *Cas Lek Cesk* 2006;145:777-81.
- 13697 *Horio M, Yasuda Y, Takahara S, Imai E, Watanabe T, Matsuo S. Comparison of a simple and a standard method for inulin renal clearance. *Clin Exp Nephrol* 2010;14:427-30.
- 58 *Hosain F, Wagner HN, Jr. Measurement of extracellular fluid volume with ¹⁶⁹Yb-diethylenetriaminepentaacetate. *J Lab Clin Med* 1971;77:699-704.
- 1990 *Hottelart C, el Esper N, Achard JM, Pruna A, Fournier A. [Fenofibrate increases blood creatinine, but does not change the glomerular filtration rate in patients with mild renal insufficiency]. *Nephrologie* 1999;20:41-4.
- 60 *Huttunen K, Huttunen NP, Koivula A, Ahonen A, Puukka R. ^{99m}Tc-DTPA--a useful clinical tool for the measurement of glomerular filtration rate. *Scand J Urol Nephrol* 1982;16:237-41.
- 61 *Ibrahim S, Rashid L, Darai M. Modification of Diet in Renal Disease equation underestimates glomerular filtration rate in Egyptian kidney donors. *Exp Clin Transplant* 2008;6:144-8.
- 13520 *Idee JM, Bonnemain B. Reliability of experimental models of iodinated contrast media-induced acute renal failure: From methodological considerations to pathophysiology. *Investigative Radiology* 1996;31:230-241.
- 2000 *Imai E. [Equation for estimating GFR from creatinine in Japan]. *Nippon Rinsho* 2008;66:1725-9.
- 2002 *Imai E, Horio M, Nitta K, Yamagata K, Iseki K, Hara S, et al. Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. *Clin Exp Nephrol* 2007;11:41-50.
- 1266 *Imai E, Horio M, Nitta K, Yamagata K, Iseki K, Tsukamoto Y, et al. Modification of the Modification of Diet in Renal Disease (MDRD) Study equation for Japan. *Am J Kidney Dis* 2007;50:927-37.
- 3473 *Imperiale A, Olianti C, Comis G, La Cava G. Evaluation of (123)I-orthoiodohippurate single kidney clearance rate by renal sequential scintigraphy in a large cohort of likely normal subjects aged between 0 and 18 years. *Eur J Nucl Med Mol Imaging* 2006;33:1483-90.
- 583 *Irving RA, Noakes TD, Burger SC, Myburgh KH, Querido D, van Zyl Smit R. Plasma volume and renal function during and after ultramarathon running. *Med Sci Sports Exerc* 1990;22:581-7.
- 13355 *Isaka Y, Fujiwara Y, Yamamoto S, Ochi S, Shin S, Inoue T, et al. Modified plasma clearance technique using nonradioactive iohalamate for measuring GFR. *Kidney Int* 1992;42:1006-11.
- 192 *Isaka Y, Yamamoto S, Fujiwara Y, Shin SH, Ochi S, Kamada T, et al. [A new analytical method for simultaneous measurement of iohalamate and iohexol]. *Rinsho Byori* 1992;40:703-7.

- 266 *Israelit AH, Long DL, White MG, Hull AR. Measurement of glomerular filtration rate utilizing a single subcutaneous injection of 125I-iothalamate. *Kidney Int* 1973;4:346-9.
- 126 *Jagenburg R, Attman PO, Aurell M, Bucht H. Determination of glomerular filtration rate in advanced renal insufficiency. *Scand J Urol Nephrol* 1978;12:133-7.
- 62 *James AE, Jr., Hosain F, DeLand FH, Reba RC, Wagner HN, Jr. 169 Ytterbium diethylenetriaminepentaacetic acid (169 YB DTPA)--a versatile radiopharmaceutical. *J Can Assoc Radiol* 1971;22:136-43.
- 13652 *Janssen A, Dumoulin A, De Waele JJ, Vanholder R, Waterloos MA, Hoste EA. Assessment of increased glomerular filtration rate in ICU patients. *SOURCE Intensive Care Medicine* (2010) 36 SUPPL. 2 (S402). Date of Publication: September 2010 CONFERENCE NAME 23rd Annual Congress of the European Society of Intensive Care Medicine, ESICM CONFERENCE LOCATION Barcelona, Spain CONFERENCE DATE 2010-10-09 to 2010-10-13.
- 13711 *Jarnberg PO, Ekstrand J, Irestedt L. Renal fluoride excretion and plasma fluoride levels during and after enflurane anesthesia are dependent on urinary pH. *Anesthesiology* 1981;54:48-52.
- 1287 *Jose PA, Fildes RD. Postnatal development of renal function. *Mead Johnson Symp Perinat Dev Med* 1988:71-6.
- 13703 *Jung K, Henke W, Schulze BD, Sydow K, Precht K, Klotzek S. Single-injection inulin clearance using only one blood sample as a suitable procedure to measure glomerular filtration rate. *Nephron* 1991;59:694-5.
- 4214 *Jung K, Henke W, Schulze BD, Sydow K, Precht K, Klotzek S. Practical approach for determining glomerular filtration rate by single-injection inulin clearance. *Clin Chem* 1992;38:403-7.
- 1290 *Kabat-Koperska J, Motyl W, Domanski L, Safranow K, Golembiewska E, Kedzierska K, et al. Methods of GFR determination--creatinine clearance after cimetidine administration in clinical practice. *Acta Med Austriaca* 2004;31:51-5.
- 602 *Kachadorian WA, Johnson RE. Renal responses to various rates of exercise. *J Appl Physiol* 1970;28:748-52.
- 13441 *Kakuta Y, Okumi M, Ichimaru N, Abe T, Nonomura N, Okuyama A, et al. Utility of the Japanese GFR estimation equation for evaluating potential donor kidney function. *Clinical and experimental nephrology* 2010;14:63-67.
- 13635 *Kamper AL. The importance of a correct evaluation of progression in studies on chronic kidney disease. *Nephrology Dialysis Transplantation* 2007;22:3-5.
- 2018 *Kampmann JP, Hansen JM. Glomerular filtration rate and creatinine clearance. *Br J Clin Pharmacol* 1981;12:7-14.
- 2020 *Karlsen SJ, Berg KJ. Acute changes in renal function following extracorporeal shock wave lithotripsy in patients with a solitary functioning kidney. *J Urol* 1991;145:253-6.
- 267 *Kasmer RJ, Cutler RE, Munger MA, Jarvis RC, Hricik D, Nara AR, et al. Single-dose effects of ibopamine hydrochloride on renal function in patients with congestive heart failure. *Br J Clin Pharmacol* 1990;30:485-9.
- 4216 *Katzberg RW, Pabico RC, Morris TW, Hayakawa K, McKenna BA, Ventura JA, et al. A comparison of clearance and arteriovenous extraction techniques for measurements of renal hemodynamic functions. *Invest Radiol* 1986;21:910-6.
- 1302 *Keevil BG, Kilpatrick ES, Nichols SP, Maylor PW. Biological variation of cystatin C: implications for the assessment of glomerular filtration rate. *Clin Chem* 1998;44:1535-9.
- 13629 *Keller F, Hartmann B. An easy to calculate equation to estimate GFR based on inulin clearance. *Nephrology Dialysis Transplantation* 2010;25:1345-1346.

2028 *Kim KE, Onesti G, Ramirez O, Brest AN, Swartz C. Creatinine clearance in renal disease. A reappraisal. *Br Med J* 1969;4:11-4.

1308 *King AJ, Levey AS. Dietary protein and renal function. *J Am Soc Nephrol* 1993;3:1723-37.

13455 *Kinowski JM, Bressolle F, Rodier M, Augey V, Fabre D, Richard JL, et al. A limited sampling model with Bayesian estimation to determine inulin pharmacokinetics using the population data modelling program P-PHARM. *Clinical Drug Investigation* 1995;9:260-269.

6214 *Kitt TM, Park GD, Spector R, Lawton W, Tsalikian E. Renal clearances of oxypurinol and inulin on an isocaloric, low-protein diet. *Clin Pharmacol Ther* 1988;43:681-7.

1310 *Klassen DK, Weir MR, Buddemeyer EU. Simultaneous measurements of glomerular filtration rate by two radioisotopic methods in patients without renal impairment. *J Am Soc Nephrol* 1992;3:108-12.

2752 *Klaus D. [Endogenous creatinine and inulin clearance.]. *Z Gesamte Inn Med* 1953;8:954-6.

63 *Kleinert J, Lorenz M, Hauser AC, Becherer A, Staudenherz A, Fodinger M, et al. Measurement of renal function in patients with Fabry disease. *Acta Paediatr Suppl* 2005;94:19-23; discussion 9-10.

64 *Klopper JF, Hauser W, Atkins HL, Eckelman WC, Richards P. Evaluation of 99m Tc-DTPA for the measurement of glomerular filtration rate. *J Nucl Med* 1972;13:107-10.

13573 *Knapp MS, Heath DA. Measurement of G.F.R. in renal disease. *Lancet* 1970;1:362-3.

2035 *Koch M, Beien A, Fusha Ller A, Zitta S, Haastert B, Trapp R. Impact of age, body mass index, insulin resistance and proteinuria on the kidney function in obese patients with Type 2 diabetes and renal insufficiency. *Clin Nephrol* 2008;69:10-7.

13653 *Koetje PM, Spaan JJ, Kooman JP, Spaanderman ME, Peeters LL. Pregnancy Reduces the Accuracy of the Estimated Glomerular Filtration Rate Based on Cockcroft-Gault and MDRD Formulas. *Reprod Sci*. 2010.

13442 *Kooman JP. Estimation of renal function in patients with chronic kidney disease. *Journal of Magnetic Resonance Imaging* 2009;30:1341-1346.

623 *Koopman MG, Koomen GC, Krediet RT, de Moor EA, Hoek FJ, Arisz L. Circadian rhythm of glomerular filtration rate in normal individuals. *Clin Sci (Lond)* 1989;77:105-11.

625 *Koopman MG, Koomen GC, van Acker BA, Arisz L. Circadian rhythm in glomerular transport of macromolecules through large pores and shunt pathway. *Kidney Int* 1996;49:1242-9.

626 *Koopman MG, Krediet RT, Arisz L. Circadian rhythms and the kidney. A review. *Neth J Med* 1985;28:416-23.

13456 *Kover G, Tost H. The renal actions of adenosine. *Acta Physiologica Hungarica* 1994;82:215-228.

2041 *Kuan Y, Hossain M, Surman J, El Nahas AM, Haylor J. GFR prediction using the MDRD and Cockcroft and Gault equations in patients with end-stage renal disease. *Nephrol Dial Transplant* 2005;20:2394-401.

13507 *Kwong YT, Stevens LA, Selvin E, Zhang YL, Greene T, Van Lente F, et al. Imprecision of urinary iothalamate clearance as a gold-standard measure of GFR decreases the diagnostic accuracy of kidney function estimating equations. In: *American journal of kidney diseases : the official journal of the National Kidney Foundation*; 2010. p 39-49.

129 *La Batide-Alanore A, Azizi M, Froissart M, Raynaud A, Plouin PF. Split renal function outcome after renal angioplasty in patients with unilateral renal artery stenosis. *J Am Soc Nephrol* 2001;12:1235-41.

13717 *Ladegaard-Pedersen HJ. Measurement of extracellular volume and renal clearance by a single injection of inulin. *Scand J Clin Lab Invest* 1972;29:145-53.

130 *Ladegaard-Pedersen HJ, Engell HC. A comparison of the distribution volumes of inulin and (51 Cr)EDTA in man and nephrectomized dogs. *Scand J Clin Lab Invest* 1972;30:267-70.

131 *Ladegaard-Pedersen HJ, Engell HC. A comparison between the changes in the distribution volumes of inulin and [51Cr]EDTA after major surgery. *Scand J Clin Lab Invest* 1975;35:109-13.

13080 *Lamb E. Assessment of kidney function in adults. *Medicine* 2007;35:359-364.

644 *Larsson A, Akerstedt T, Hansson LO, Axelsson J. Circadian variability of cystatin C, creatinine, and glomerular filtration rate (GFR) in healthy men during normal sleep and after an acute shift of sleep. *Chronobiol Int* 2008;25:1047-61.

1339 *Larsson M, Jagenburg R, Landahl S. Renal function in an elderly population. A study of S-creatinine, 51Cr-EDTA clearance, endogenous creatinine clearance and maximal tubular water reabsorption. *Scand J Clin Lab Invest* 1986;46:593-8.

4236 *Lavender S, Hilton PJ, Jones NF. The measurement of glomerular filtration-rate in renal disease. *Lancet* 1969;2:1216-8.

2054 *Leblond V, Sutton L, Jacquiaud C, Item C, Sadoun R, Jaudon MC, et al. Evaluation of renal function in 60 long-term survivors of bone marrow transplantation. *J Am Soc Nephrol* 1995;6:1661-5.

13654 *Lee CS, Cha RH, Lim YH, Kim H, Song KH, Gu N, et al. Ethnic coefficients for glomerular filtration rate estimation by the Modification of Diet in Renal Disease study equations in the Korean population. *J Korean Med Sci*. 2010;25:1616-25.

2060 *Lema G, Meneses G, Urzua J, Jalil R, Canessa R, Moran S, et al. Effects of extracorporeal circulation on renal function in coronary surgical patients. *Anesth Analg* 1995;81:446-51.

2061 *Lema G, Urzua J, Jalil R, Canessa R, Moran S, Sacco C, et al. Renal protection in patients undergoing cardiopulmonary bypass with preoperative abnormal renal function. *Anesth Analg* 1998;86:3-8.

2064 *Lema G, Vogel A, Canessa R, Jalil R, Carvajal C, Becker P, et al. Renal function and cardiopulmonary bypass in pediatric cardiac surgical patients. *Pediatr Nephrol* 2006;21:1446-51.

271 *Lemann J, Bidani AK, Bain RP, Lewis EJ, Rohde RD. Use of the serum creatinine to estimate glomerular filtration rate in health and early diabetic nephropathy. Collaborative Study Group of Angiotensin Converting Enzyme Inhibition in Diabetic Nephropathy. *Am J Kidney Dis* 1990;16:236-43.

13460 *Lemann Jr J, Bidani AK, Bain RP, Lewis EJ, Rohde RD. Use of the serum creatinine to estimate glomerular filtration rate in health and early diabetic nephropathy. *American Journal of Kidney Diseases* 1990;16:236-243.

13645 *Levey AS. Use of glomerular filtration rate measurements to assess the progression of renal disease. *Seminars in Nephrology* 1989;9:370-379.

13362 *Levey AS, Berg RL, Gassman JJ, Hall PM, Walker WG. Creatinine filtration, secretion and excretion during progressive renal disease. Modification of Diet in Renal Disease (MDRD) Study Group. *Kidney Int Suppl* 1989;27:S73-80.

13363 *Levey AS, Greene T, Schluchter MD, Cleary PA, Teschan PE, Lorenz RA, et al. Glomerular filtration rate measurements in clinical trials. Modification of Diet in Renal Disease Study Group and the Diabetes Control and Complications Trial Research Group. *J Am Soc Nephrol* 1993;4:1159-71.

65 *Lewis R, Kerr N, Van Buren C, Lowry P, Sandler C, Frazier OH, et al. Comparative evaluation of urographic contrast media, inulin, and 99mTc-DTPA clearance methods for determination of glomerular filtration rate in clinical transplantation. *Transplantation* 1989;48:790-6.

1357 *Lin J, Knight EL, Hogan ML, Singh AK. A comparison of prediction equations for estimating glomerular filtration rate in adults without kidney disease. *J Am Soc Nephrol* 2003;14:2573-80.

13524 *Lindberg JS, Copley JB. Evaluation of glomerular filtration rate in intensive care unit patients. *Journal of Intensive Care Medicine* 1992;7:283-292.

- 197 *Lindblad HG, Berg UB. Comparative evaluation of iohexol and inulin clearance for glomerular filtration rate determinations. *Acta Paediatr* 1994;83:418-22.
- 13644 *Lloveras JJ, Durand D, Ader JL, Tran Van T, Tack I, Rostaing L, et al. Comparison between glomerular filtration rate and renal plasma flow between kidney and heart transplant recipients maintained on a similar cyclosporine dosage. *Transplantation Proceedings* 1994;26:248.
- 13630 *Lorenzen J, Kramer R, Kliem V, Bode-Boeger SM, Veldink H, Haller H, et al. Circulating levels of osteopontin are closely related to glomerular filtration rate and cardiovascular risk markers in patients with chronic kidney disease. *European Journal of Clinical Investigation* 2010;40:294-300.
- 274 *Ludescher E, Hohenwallner W, Riccabona J. [Single injection clearance with sodium iothalamate I-131 in children (author's transl)]. *Pediatr Padol* 1971;6:357-64.
- 668 *Luft FC, Fineberg NS, Miller JZ, Rankin LI, Grim CE, Weinberger MH. The effects of age, race and heredity on glomerular filtration rate following volume expansion and contraction in normal man. *Am J Med Sci* 1980;279:15-24.
- 2083 *Luke DR, Halstenson CE, Opsahl JA, Matzke GR. Validity of creatinine clearance estimates in the assessment of renal function. *Clin Pharmacol Ther* 1990;48:503-8.
- 13395 *Macdonald JH, Marcora SM, Jibani M, Roberts G, Kumwenda MJ, Glover R, et al. Bioelectrical impedance can be used to predict muscle mass and hence improve estimation of glomerular filtration rate in non-diabetic patients with chronic kidney disease. *Nephrology Dialysis Transplantation* 2006;21:3481-3487.
- 13396 *Macdonald JH, Marcora SM, Kumwenda MJ, Jibani M, Roberts G, Glover R, et al. The relationship between estimated glomerular filtration rate, demographic and anthropometric variables is mediated by muscle mass in non-diabetic patients with chronic kidney disease. *Nephrology Dialysis Transplantation* 2006;21:3488-3494.
- 669 *Macisaac RJ, Jerums G. Albuminuric and non-albuminuric pathways to renal impairment in diabetes. *Minerva Endocrinol* 2005;30:161-77.
- 1374 *Mahajan S, Mukhiya GK, Singh R, Tiwari SC, Kalra V, Bhowmik DM, et al. Assessing glomerular filtration rate in healthy Indian adults: a comparison of various prediction equations. *J Nephrol* 2005;18:257-61.
- 277 *Maher FT, Nolan NG, Elveback LR. Comparison of simultaneous clearances of 125-I-labeled sodium iothalamate (Glofil) and of inulin. *Mayo Clin Proc* 1971;46:690-1.
- 278 *Maher FT, Tauxe WN. Renal clearance in man of pharmaceuticals containing radioactive iodine. *JAMA* 1969;207:97-104.
- 2087 *Maillard N, Mariat C, Bonneau C, Mehdi M, Thibaudin L, Laporte S, et al. Cystatin C-based equations in renal transplantation: moving toward a better glomerular filtration rate prediction? *Transplantation* 2008;85:1855-8.
- 279 *Mak RH, Al Dahhan J, Azzopardi D, Bosque M, Chantler C, Haycock GB. Measurement of glomerular filtration rate in children after renal transplantation. *Kidney Int* 1983;23:410-3.
- 280 *Malamos B, Dontas AS, Koutras DA, Marketos S, Sfontouris J, Papanicolaou N. 125I-sodium iothalamate in the determination of the glomerular filtration rate. *Nucl Med (Stuttg)* 1967;6:304-10.
- 137 *Manz F, Alatas H, Kochen W, Lutz P, Rebien W, Scharer K. Determination of glomerular function in advanced renal failure. *Arch Dis Child* 1977;52:721-4.
- 13623 *Manzar S, Nair AK, Pai MG, Al-Khusaiby SM. A simplified bedside method for estimating glomerular filtration rate in term neonates. *Saudi Medical Journal* 2004;25:1501-1502.
- 13637 *Mariat C, Alamartine E, Afiani A, Thibaudin L, Laurent B, Berthoux P, et al. Predicting glomerular filtration rate in kidney transplantation: Are the K/DOQI guidelines applicable? *American Journal of Transplantation* 2005;5:2698-2703.

- 2095 *Mariat C, Alamartine E, Barthelemy JC, De Filippis JP, Thibaudin D, Berthoux P, et al. Assessing renal graft function in clinical trials: can tests predicting glomerular filtration rate substitute for a reference method? *Kidney Int* 2004;65:289-97.
- 2099 *Martini S, Prevot A, Mosig D, Werner D, van Melle G, Guignard JP. Glomerular filtration rate: measure creatinine and height rather than cystatin C! *Acta Paediatr* 2003;92:1052-7.
- 281 *Materson BJ. Measurement of glomerular filtration rate. *CRC Crit Rev Clin Lab Sci* 1971;2:1-43.
- 4241 *Mazzachi BC, Peake MJ, Ehrhardt V. Reference range and method comparison studies for enzymatic and Jaffe creatinine assays in plasma and serum and early morning urine. *Clin Lab* 2000;46:53-5.
- 2110 *McDougal WS, Koch MO. Accurate determination of renal function in patients with intestinal urinary diversions. *J Urol* 1986;135:1175-8.
- 138 *McElduff A, Shuter B, Cooper R, Davies L, Fulcher G, Hoschl R, et al. Measuring renal function in patients with diabetes mellitus. *J Diabetes Complications* 1997;11:225-9.
- 2827 *McGonigal MD, Lucas CE, Ledgerwood AM. The effects of treatment of renal trauma on renal function. *J Trauma* 1987;27:471-6.
- 139 *Medeiros FS, Sapienza MT, Prado ES, Agena F, Shimizu MH, Lemos FB, et al. Validation of plasma clearance of 51Cr-EDTA in adult renal transplant recipients: comparison with inulin renal clearance. *Transpl Int* 2009;22:323-31.
- 2829 *Mertz DP, Sarre H, Cremer Z. [On the diagnostic value of semiquantitative kidney function tests. III. Correlations between the plasma concentrations of rest nitrogen and "true" endogenous creatinine and inulin clearance.]. *Klin Wochenschr* 1962;40:889-93.
- 2830 *Mertz DP, Sarre H, Cremer Z. [On the diagnostic value of semi-quantitative kidney function tests. I. Correlation of plasma concentration of "genuine" endogenous creatinine and inulin clearance.]. *Klin Wochenschr* 1962;40:687-92.
- 67 *Milutinovic J, Cutler RE, Hoover P, Meijssen B, Scribner BH. Measurement of residual glomerular filtration rate in the patient receiving repetitive hemodialysis. *Kidney Int* 1975;8:185-90.
- 13399 *Mingat N, Allard J, Villard E, Guillotreau J, Paparel P, Roche JB, et al. Which is the most accurate biomarker to assess renal functions in neurogenic patients? *European Urology, Supplements* 2010;9:180-181.
- 285 *Mirouze J, Monnier L, Olivier C. Study of renal functions by repeated constant infusion of radiotracers before and after initiation of therapy in hypertension or in diabetes mellitus. *Pathol Biol (Paris)* 1975;23:599-603.
- 286 *Mogensen CE. Glomerular filtration rate and renal plasma flow in short-term and long-term juvenile diabetes mellitus. *Scand J Clin Lab Invest* 1971;28:91-100.
- 140 *Monteiro MC, Alonso G, Ajzen H, Pereira AB. Assessment of glomerular filtration rate utilizing subcutaneously injected 51Cr-EDTA. *Braz J Med Biol Res* 1994;27:2557-64.
- 68 *Moran SM, Myers BD. Pathophysiology of protracted acute renal failure in man. *J Clin Invest* 1985;76:1440-8.
- 69 *Morton KA, Pisani DE, Whiting JH, Jr., Cheung AK, Arias JM, Valdivia S. Determination of glomerular filtration rate using technetium-99m-DTPA with differing degrees of renal function. *J Nucl Med Technol* 1997;25:110-4.
- 142 *Motwani JG, Fenwick MK, Struthers AD. Comparison of three methods of glomerular filtration rate measurement with and without captopril pretreatment in groups of patients with left ventricular dysfunction. *Eur Heart J* 1992;13:1195-200.
- 1408 *Motyl W. [The effect of H2 receptor blockers on renal function. An attempt to accurately measure glomerular filtration]. *Ann Acad Med Stetin* 2004;50:97-105.

- 2131 *Mpio I, Laville M, Hadj-Aissa A, Fauvel JP. Predicted creatinine clearance to evaluate glomerular filtration rate in black Caribbean subjects. *Nephrol Dial Transplant* 2003;18:1307-10.
- 2133 *Mujais SK, Quintanilla A, Zahid M, Koch K, Shaw W, Gibson T. Renal handling of enalaprilat. *Am J Kidney Dis* 1992;19:121-5.
- 143 *Muller-Suur R, Goransson M, Olsen L, Backlund G, Backlund L. Inulin single injection clearance. Microsample technique useful in children for determination of glomerular filtration rate. *Clin Physiol* 1983;3:19-27.
- 2134 *Munar MY, Elzinga L, Brummett R, Golper TA, Bennett WM. The effect of tobramycin on the renal handling of vancomycin. *J Clin Pharmacol* 1991;31:618-23.
- 2851 *Murphy GP, Gagnon JA. THE ALTERATIONS IN GLOMERULAR FILTRATION RATE, OSMOLAR, AND FREE WATER CLEARANCE DURING MANNITOL OSMOTIC DIURESIS. *J Urol* 1964;92:17-24.
- 2136 *Murray MD, Black PK, Kuzmik DD, Haag KM, Manatunga AK, Mullin MA, et al. Acute and chronic effects of nonsteroidal antiinflammatory drugs on glomerular filtration rate in elderly patients. *Am J Med Sci* 1995;310:188-97.
- 2137 *Murray MD, Greene PK, Brater DC, Manatunga AK, Hall SD. Effects of flurbiprofen on renal function in patients with moderate renal insufficiency. *Br J Clin Pharmacol* 1992;33:385-93.
- 6897 *Muther RS, Bennett WM. Effects of aspirin on glomerular filtration rate in normal humans. *Ann Intern Med* 1980;92:386-7.
- 1417 *Myers GL, Miller WG, Coresh J, Fleming J, Greenberg N, Greene T, et al. Recommendations for improving serum creatinine measurement: a report from the Laboratory Working Group of the National Kidney Disease Education Program. *Clin Chem* 2006;52:5-18.
- 3696 *Nakai K, Kikuchi M, Koyamaa T, Saito K, Suwabe A. [Evaluation of glomerular filtration rate in children--serum markers and prediction equations]. *Rinsho Byori* 2009;57:338-44.
- 13706 *Nielsen OM, Engell HC. Increased glomerular filtration rate in patients after reconstructive surgery on the abdominal aorta. *Br J Surg* 1986;73:34-7.
- 2144 *Nitta K, Hayashi T, Uchida K, Honda K, Tsukada M, Sekine S, et al. Serum cystatin C concentration as a marker of glomerular filtration rate in patients with various renal diseases. *Intern Med* 2002;41:931-5.
- 70 *Notghi A, Merrick MV, Ferrington C, Anderton JL. A comparison of simplified and standard methods for the measurement of glomerular filtration rate and renal tubular function. *Br J Radiol* 1986;59:35-9.
- 2149 *Nowack R, Hofner MC, Reichel H, Schmidt-Gayk H, Ritz E. Subacute effects of thiazide administration on renal hemodynamics and calcium metabolism. *Clin Investig* 1992;70:686-91.
- 145 *Oberhausen E. [Investigation of renal clearance in nuclear medicine (author's transl)]. *Radiologe* 1981;21:548-52.
- 13453 *Oda A, Iwatsubo E, Haraoka M, Segawa J, Kitada S, Osada Y. Clinical significance of serum (beta)(2)-microglobulin as a predictor of renal function in patients with spinal cord injury. *Nishinihon Journal of Urology* 1996;58:5-8.
- 146 *Odlind B, Hallgren R, Sohtell M, Lindstrom B. Is 125I iothalamate an ideal marker for glomerular filtration? *Kidney Int* 1985;27:9-16.
- 2155 *Opravil M, Keusch G, Luthy R. Pyrimethamine inhibits renal secretion of creatinine. *Antimicrob Agents Chemother* 1993;37:1056-60.
- 2157 *Orita Y, Gejyo F, Sakatsume M, Shiigai T, Maeda Y, Imai E, et al. [Estimation of glomerular filtration rate by inulin clearance: comparison with creatinine clearance]. *Nippon Jinzo Gakkai Shi* 2005;47:804-12.

- 2159 *Orita Y, Okada M, Harada S, Horio M. Skim soy protein enhances GFR as much as beefsteak protein in healthy human subjects. *Clin Exp Nephrol* 2004;8:103-8.
- 2161 *Orlando R, Floreani M, Napoli E, Padrini R, Palatini P. Renal clearance of N(1)-methylnicotinamide: a sensitive marker of the severity of liver dysfunction in cirrhosis. *Nephron* 2000;84:32-9.
- 13700 *Orlando R, Floreani M, Padrini R, Palatini P. Determination of inulin clearance by bolus intravenous injection in healthy subjects and ascitic patients: equivalence of systemic and renal clearances as glomerular filtration markers. *Br J Clin Pharmacol* 1998;46:605-9.
- 2163 *Orlando R, Floreani M, Padrini R, Palatini P. Evaluation of measured and calculated creatinine clearances as glomerular filtration markers in different stages of liver cirrhosis. *Clin Nephrol* 1999;51:341-7.
- 1451 *Orlando R, Mussap M, Plebani M, Piccoli P, De Martin S, Floreani M, et al. Diagnostic value of plasma cystatin C as a glomerular filtration marker in decompensated liver cirrhosis. *Clin Chem* 2002;48:850-8.
- 751 *Orth SR, Schroeder T, Ritz E, Ferrari P. Effects of smoking on renal function in patients with type 1 and type 2 diabetes mellitus. *Nephrol Dial Transplant* 2005;20:2414-9.
- 13632 *Ott C, Ritt M, Titze SI, Schaufele T, Schmieder RE. Rosuvastatin does not affect intrarenal hemodynamics in patients with hypercholesterolemia. *Journal of Nephrology* 2009;22:675-681.
- 13516 *Ott NT. A simple technique for estimating glomerular filtration rate with subcutaneous injection of (125I)iothalamate. *Mayo Clin Proc.* 1975;50:664-8.
- 13514 *Ovcharenko E, Hansen MK, Goltsman I, Abu-Saleh N, Abassi Z, Walsh K, et al. Peroxisome proliferator-activated receptor alpha and Alpha/Gamma agonists do not cause impairment in renal function in the rat. *Nephron - Physiology* 2010;115:p21-p30.
- 754 *Paaby P, Brochner-Mortensen J, Fjeldborg P, Raffn K, Larsen CE, Moller-Petersen J. Endogenous overnight creatinine clearance compared with 51Cr-EDTA clearance during the menstrual cycle. *Acta Med Scand* 1987;222:281-4.
- 13464 *Pathe M, Schwartz J. Renal hemodynamics and function during oral administration of betaxolol
ORIGINAL (NON-ENGLISH) TITLE MODIFICATIONS DE L'HEMODYNAMIQUE ET DE LA FONCTION RENALE DURANT L'ADMINISTRATION ORALE DE BETAXOLOL. *Therapie* 1985;40:209-212.
- 1465 *Payne RB. Creatinine clearance: a redundant clinical investigation. *Ann Clin Biochem* 1986;23 (Pt 3):243-50.
- 771 *Pena JC, Herrera Acosta J. [The circadian rhythms of the kidney]. *Bol Estud Med Biol* 1976;29:47-60.
- 2175 *Perez A, Raab R, Chen TC, Turner A, Holick MF. Safety and efficacy of oral calcitriol (1,25-dihydroxyvitamin D3) for the treatment of psoriasis. *Br J Dermatol* 1996;134:1070-8.
- 4242 *Perrone RD, Madias NE, Levey AS. Serum creatinine as an index of renal function: new insights into old concepts. *Clin Chem* 1992;38:1933-53.
- 71 *Perrone RD, Steinman TI, Beck GJ, Skibinski CI, Royal HD, Lawlor M, et al. Utility of radioisotopic filtration markers in chronic renal insufficiency: simultaneous comparison of 125I-iothalamate, 169Yb-DTPA, 99mTc-DTPA, and inulin. *The Modification of Diet in Renal Disease Study. Am J Kidney Dis* 1990;16:224-35.
- 72 *Peters AM, Henderson BL, Lui D, Blunkett M, Cosgriff PS, Myers MJ. Appropriate corrections to glomerular filtration rate and volume of distribution based on the bolus injection and single-compartment technique. *Physiol Meas* 1999;20:313-27.
- 73 *Petri M, Bockenstedt L, Colman J, Whiting-O'Keefe Q, Fitz G, Sebastian A, et al. Serial assessment of glomerular filtration rate in lupus nephropathy. *Kidney Int* 1988;34:832-9.
- 200 *Piepsz A. Iohexol and inulin clearance. *Acta Paediatr* 1994;83:1160.

- 74 *Piepsz A, Dobbeleir A, Erbsmann F. Measurement of separate kidney clearance by means of ^{99m}Tc-DTPA complex and a scintillation camera. *Eur J Nucl Med* 1977;2:173-7.
- 75 *Piepsz A, Erbsmann F, Vaincel M, Cantraine F. Determination of glomerular filtration rate in children by external measurement of the disappearance of ^{113m}indium-DTPA in plasma. *Biomedicine* 1974;20:192-7.
- 2178 *Piepsz A, Ham H. [On what is renal clearance based]. *Arch Pediatr* 2000;7:213-4.
- 148 *Piepsz A, Pintelon H, Ham HR. Estimation of normal chromium-51 ethylene diamine tetra-acetic acid clearance in children. *Eur J Nucl Med* 1994;21:12-6.
- 1480 *Piepsz A, Tondeur M, Ham H. Revisiting normal (⁵¹Cr)-ethylenediaminetetraacetic acid clearance values in children. *Eur J Nucl Med Mol Imaging* 2006;33:1477-82.
- 1481 *Piepsz A, Tondeur M, Ham H. Escaping the correction for body surface area when calculating glomerular filtration rate in children. *Eur J Nucl Med Mol Imaging* 2008;35:1669-72.
- 2181 *Pierrat A, Gravier E, Saunders C, Caira MV, Ait-Djafer Z, Legras B, et al. Predicting GFR in children and adults: a comparison of the Cockcroft-Gault, Schwartz, and modification of diet in renal disease formulas. *Kidney Int* 2003;64:1425-36.
- 296 *Pixberg HU, Hilger P, Zielske F. [Determination of glomerular filtration rate with ¹²⁵I iothalamate]. *Fortschr Geb Rontgenstr Nuklearmed* 1969;111:108-11.
- 2186 *Plebani M, Dall'Amico R, Mussap M, Montini G, Ruzzante N, Marsilio R, et al. Is serum cystatin C a sensitive marker of glomerular filtration rate (GFR)? A preliminary study on renal transplant patients. *Ren Fail* 1998;20:303-9.
- 7226 *Podracka L, Feber J, Lepage N, Filler G. Intra-individual variation of cystatin C and creatinine in pediatric solid organ transplant recipients. *Pediatr Transplant* 2005;9:28-32.
- 1485 *Poggio ED, Rule AD. Can we do better than a single estimated GFR threshold when screening for chronic kidney disease? *Kidney Int* 2007;72:534-6.
- 3768 *Poggio ED, Rule AD, Tanchanco R, Arrigain S, Butler RS, Srinivas T, et al. Demographic and clinical characteristics associated with glomerular filtration rates in living kidney donors. *Kidney Int* 2009;75:1079-87.
- 2192 *Poinsot O, Romand JA, Favre H, Suter PM. Fenoldopam improves renal hemodynamics impaired by positive end-expiratory pressure. *Anesthesiology* 1993;79:680-4.
- 13352 *Ponteil-Noble C. Is there any relationship between the early glomerular filtration rate and the late graft dysfunction in renal transplantation? In: Touraine JL, editor. *Late graft loss*: Kluwer Academic Publishers Group; 1997. p 31-37.
- 781 *Poortmans JR. Exercise and renal function. *Exerc Sport Sci Rev* 1977;5:255-94.
- 782 *Poortmans JR. Exercise and renal function. *Sports Med* 1984;1:125-53.
- 784 *Poortmans JR, Jourdain M, Heyters C, Reardon FD. Postexercise proteinuria in rowers. *Can J Sport Sci* 1990;15:126-30.
- 785 *Poortmans JR, Ouchinsky M. Glomerular filtration rate and albumin excretion after maximal exercise in aging sedentary and active men. *J Gerontol A Biol Sci Med Sci* 2006;61:1181-5.
- 13715 *Poulsen HL, Jensen HA, Parving HH. Extracellular fluid volume determined by a single injection of inulin in men with untreated essential hypertension. *Scand J Clin Lab Invest* 1977;37:691-6.
- 7270 *Pouteil-Noble C, Chapuis F, Berra N, Hadj-Aissa A, Lacavalerie B, Lefrancois N, et al. Misoprostol in renal transplant recipients: a prospective, randomized, controlled study on the prevention of acute rejection episodes and cyclosporin A nephrotoxicity. *Nephrol Dial Transplant* 1994;9:552-5.

- 297 *Pouteil-Noble C, Maiza H, Remontet L. Post-transplant glomerular filtration rate as a marker for long-term outcome. *Ann Transplant* 2000;5:29-36.
- 2200 *Prescott LF, Freestone S, McAuslane JA. Reassessment of the single intravenous injection method with inulin for measurement of the glomerular filtration rate in man. *Clin Sci (Lond)* 1991;80:167-76.
- 13470 *Preuss LE, Michaels RS, Bolin FP. Evaluation of a simplified measurement for low glomerular filtration rates with indium-111 DTPA. *Henry Ford Hospital Medical Journal* 1979;27:122-126.
- 201 *Prevot A, Martini S, Guignard JP. [Glomerular filtration markers in pediatrics]. *Rev Med Suisse Romande* 2002;122:625-30.
- 1491 *Price DA, Fisher ND, Osei SY, Lansang MC, Hollenberg NK. Renal perfusion and function in healthy African Americans. *Kidney Int* 2001;59:1037-43.
- 13649 *Price M. Inulin clearance as a screening test for kidney function. *Archives of Physical Medicine and Rehabilitation* 1974;55:522-523.
- 2209 *Proulx NL, Akbari A, Garg AX, Rostom A, Jaffey J, Clark HD. Measured creatinine clearance from timed urine collections substantially overestimates glomerular filtration rate in patients with liver cirrhosis: a systematic review and individual patient meta-analysis. *Nephrol Dial Transplant* 2005;20:1617-22.
- 298 *Prueksaritanont T, Lui CY, Lee MG, Chiou WL. Renal and non-renal clearances of iothalamate. *Biopharm Drug Dispos* 1986;7:347-55.
- 2215 *Rahman SN, Conger JD. Glomerular and tubular factors in urine flow rates of acute renal failure patients. *Am J Kidney Dis* 1994;23:788-93.
- 7329 *Rahman SN, Kim GE, Mathew AS, Goldberg CA, Allgren R, Schrier RW, et al. Effects of atrial natriuretic peptide in clinical acute renal failure. *Kidney Int* 1994;45:1731-8.
- 202 *Rahn KH, Heidenreich S, Bruckner D. How to assess glomerular function and damage in humans. *J Hypertens* 1999;17:309-17.
- 2218 *Rapoport A, Husdan H. Endogenous creatinine clearance and serum creatinine in the clinical assessment of kidney function. *Can Med Assoc J* 1968;99:149-56.
- 4227 *Reba RC, Hosain F, Wagner HN, Jr. Indium-113m diethylenetriaminepentaacetic acid (DTPA): a new radiopharmaceutical for study of the kidneys. *Radiology* 1968;90:147-9.
- 76 *Rehling M, Moller ML, Jensen JJ, Thamdrup B, Lund JO, Trap-Jensen J. Reliability of single kidney glomerular filtration rate measured by a 99mTc-DTPA gamma camera technique. *Uremia Invest* 1985;9:171-6.
- 77 *Rehling M, Moller ML, Thamdrup B, Lund JO, Trap-Jensen J. Simultaneous measurement of renal clearance and plasma clearance of 99mTc-labelled diethylenetriaminepenta-acetate, 51Cr-labelled ethylenediaminetetraacetate and inulin in man. *Clin Sci (Lond)* 1984;66:613-9.
- 78 *Rehling M, Moller ML, Thamdrup B, Lund JO, Trap-Jensen J. Reliability of a 99mTc-DTPA gamma camera technique for determination of single kidney glomerular filtration rate. A comparison to plasma clearance of 51Cr-EDTA in one-kidney patients, using the renal clearance of inulin as a reference. *Scand J Urol Nephrol* 1986;20:57-62.
- 13463 *Reubi FC, Franz KA, Horber F. Hypertension as related to renal function in diabetes mellitus. *Hypertension* 1985;7:21-28.
- 151 *Riccabona G, Dittrich P, Knapp E, Bauer H. [Comparative studies on various methods of determining the glomerular filtration rate with 51 Cr-EDTA]. *Nucl Med (Stuttg)* 1970;9:242-7.
- 813 *Ritz E, Benck U, Franek E, Keller C, Seyfarth M, Clorius J. Effects of smoking on renal hemodynamics in healthy volunteers and in patients with glomerular disease. *J Am Soc Nephrol* 1998;9:1798-804.

- 2226 *Robert S, Zarowitz BJ. Is there a reliable index of glomerular filtration rate in critically ill patients? DICP 1991;25:169-78.
- 2227 *Robert S, Zarowitz BJ, Peterson EL, Dumler F. Predictability of creatinine clearance estimates in critically ill patients. Crit Care Med 1993;21:1487-95.
- 203 *Rodrigo E, de Francisco AL, Escallada R, Ruiz JC, Fresnedo GF, Pinera C, et al. Measurement of renal function in pre-ESRD patients. Kidney Int Suppl 2002:11-7.
- 13648 *Rosenbaum R, Hruska K, Anderson C. Inulin, an inadequate marker of glomerular filtration (GF) in the transplanted kidney. Clinical Research 1977;25:596 A.
- 301 *Rosenbaum RW, Hruska KA, Anderson C, Robson AM, Slatopolsky E, Klahr S. Inulin: an inadequate marker of glomerular filtration rate in kidney donors and transplant recipients? Kidney Int 1979;16:179-86.
- 79 *Roubenoff R, Drew H, Moyer M, Petri M, Whiting-O'Keefe Q, Hellmann DB. Oral cimetidine improves the accuracy and precision of creatinine clearance in lupus nephritis. Ann Intern Med 1990;113:501-6.
- 2251 *Roy L, Legault L, Pomier-Layrargues G. Glomerular filtration rate measurement in cirrhotic patients with renal failure. Clin Nephrol 1998;50:342-6.
- 1523 *Rule AD, Gussak HM, Pond GR, Bergstralh EJ, Stegall MD, Cosio FG, et al. Measured and estimated GFR in healthy potential kidney donors. Am J Kidney Dis 2004;43:112-9.
- 1526 *Rutland MD. A comprehensive analysis of renal DTPA studies. I. Theory and normal values. Nucl Med Commun 1985;6:11-20.
- 1527 *Rydstrom M, Tengstrom B, Cederquist I, Ahlmen J. Measurement of glomerular filtration rate by single-injection, single-sample techniques, using ⁵¹Cr-EDTA or iohexol. Scand J Urol Nephrol 1995;29:135-9.
- 2256 *Saile P, Fiedler R, Markau S, Kuhn C, Osten B. [Determination of renal function in clinical routine: which is the best method?]. Dtsch Med Wochenschr 2007;132:1093-7.
- 13231 *Sambasivan AS, Lepage N, Filler G. Cystatin C inpatient variability in children with chronic kidney disease is less than serum creatinine. Clin Chem. 2005;51:2215-6.
- 13364 *Sambataro M, Thomaseth K, Pacini G, Robaudo C, Carraro A, Bruseghin M, et al. Plasma clearance rate of ⁵¹Cr-EDTA provides a precise and convenient technique for measurement of glomerular filtration rate in diabetic humans. J Am Soc Nephrol 1996;7:118-27.
- 2258 *Sansoe G, Ferrari A, Castellana CN, Bonardi L, Villa E, Manenti F. Cimetidine administration and tubular creatinine secretion in patients with compensated cirrhosis. Clin Sci (Lond) 2002;102:91-8.
- 153 *Saudan P, Muller H, Feraille E, Martin PY, Mach F. Renin-angiotensin system blockade and contrast-induced renal toxicity. J Nephrol 2008;21:681-5.
- 1537 *Schaefer RM, Seufert H, Schafferhans K, Gilge U, Schulz HU, Pooth R, et al. [Determination of glomerular filtration rate in man using single shot piracetam clearance]. Internist (Berl) 1992;33 Suppl 1:S23-6.
- 1540 *Schirmeister J. [Proceedings: Normal values and samples of findings in kidney diseases]. Med Klin 1975;70:1004-5.
- 154 *Schroder H, Schutte H. [Comparative determination of glomerular filtration with inulin and with Cr⁵¹-EDTA in hypertensive patients and diabetics]. Fortschr Geb Rontgenstr Nuklearmed 1969;111:573-6.
- 1548 *Schuck O, Gottfriedova H, Maly J, Jabor A, Stolova M, Bruzkova I, et al. Glomerular filtration rate assessment in individuals after orthotopic liver transplantation based on serum cystatin C levels. Liver Transpl 2002;8:594-9.
- 1551 *Schuck O, Teplan V, Jabor A, Stolova M, Skibova J. Glomerular filtration rate estimation in patients with advanced chronic renal insufficiency based on serum cystatin C levels. Nephron Clin Pract 2003;93:c146-51.

- 2282 *Schuck O, Teplan V, Nadvornikova H, Skibova J. [Relation between creatinine clearance and glomerular filtration in various stages of chronic renal disease]. *Vnitr Lek* 1992;38:1181-6.
- 13450 *Schuck O, Teplan V, Skibova J, Stolova M. Predicting the GFR in Patients with CRF from the Arithmetic Mean of Creatinine and Urea Clearances: A Simple Mathematical Explanation of an Assumed Coincidence. *Dialysis and Transplantation* 2003;32:666-674.
- 80 *Schulze BD, Sydow K, Precht K, Strangfeld D, Schurer M, Buchali K, et al. [Studies and results of the use of nuclear medicine methods in kidney function tests]. *Z Urol Nephrol* 1984;77:647-54.
- 1552 *Schumann L, Wustenberg PW, Hortian B, Kuhnle HF. Determination of glomerular filtration rate (GFR) on two consecutive days using inulin in a single-sample plasma clearance method. *Clin Nephrol* 1993;39:65-9.
- 155 *Schwartz GJ, Furth SL. Glomerular filtration rate measurement and estimation in chronic kidney disease. *Pediatr Nephrol* 2007;22:1839-48.
- 16 *Schwartz GJ, Haycock GB, Edelmann CM, Jr., Spitzer A. A simple estimate of glomerular filtration rate in children derived from body length and plasma creatinine. *Pediatrics* 1976;58:259-63.
- 13245 *Schwartz GJ, Work DF. Measurement and estimation of GFR in children and adolescents. *Clin J Am Soc Nephrol*. 2009;4:1832-43.
- 304 *Seki T, Orita Y, Yamamoto S, Ueda N, Yanagihara Y, Noguchi K. [A method for the simultaneous determination of iothalamate and creatinine clearance by coupled-column liquid chromatography]. *Rinsho Byori* 1992;40:697-702.
- 81 *Shemesh O, Golbetz H, Kriss JP, Myers BD. Limitations of creatinine as a filtration marker in glomerulopathic patients. *Kidney Int* 1985;28:830-8.
- 13655 *Shimokata T, Ando Y, Yasuda Y, Hamada A, Kawada K, Saito H, et al. Prospective evaluation of pharmacokinetically guided dosing of carboplatin in Japanese patients with cancer. *Cancer Sci*. 2010;101:2601-5.
- 13445 *Shimokata T, Yasuda Y, Hamada A, Kawada K, Hasagawa Y, Ando Y. Evaluation of renal function and carboplatin clearance in Japanese cancer patients. *Journal of Clinical Oncology* 2009;27:2556.
- 2300 *Shirota T. [Studies on the renal handling of urea nitrogen, creatinine, water and electrolytes in hyperthyroid patients with Graves' disease]. *Nippon Naibunpi Gakkai Zasshi* 1991;67:611-21.
- 1562 *Shirota T, Shinoda T, Yamada T, Aizawa T. Alteration of renal function in hyperthyroidism: increased tubular secretion of creatinine and decreased distal tubule delivery of chloride. *Metabolism* 1992;41:402-5.
- 13354 *Sigman EM, Elwood C, Reagan ME, Morris AM, Catanzaro A. The Renal Clearance of I-131 Labelled Sodium Iothalamate in Man. *Invest Urol* 1965;2:432-8.
- 305 *Sigman EM, Elwood CM, Knox F. The measurement of glomerular filtration rate in man with sodium iothalamate 131-I (Conray). *J Nucl Med* 1966;7:60-8.
- 306 *Silkals GI, Jeck D, Earon J, Edelmann CM, Jr., Chervu LR, Blaufox MD, et al. Simultaneous measurement of glomerular filtration rate and renal plasma flow using plasma disappearance curves. *J Pediatr* 1973;83:749-57.
- 862 *Sims EA. Renal function in normal pregnancy. *Clin Obstet Gynecol* 1968;11:461-72.
- 863 *Singh NP, Ingle GK, Saini VK, Jami A, Beniwal P, Lal M, et al. Prevalence of low glomerular filtration rate, proteinuria and associated risk factors in North India using Cockcroft-Gault and Modification of Diet in Renal Disease equation: an observational, cross-sectional study. *BMC Nephrol* 2009;10:4.
- 13262 *Sirwal IA, Bandy KA, Reshi AR, Bhat MA, Wani MM. Estimation of glomerular filtration rate (GFR). *JK Science* 2004;6:121-123.
- 307 *Skov PE. Glomerular filtration rate in patients with severe and very severe renal insufficiency. Determined by simultaneous inulin, creatinine and 125 iothalamate clearance. *Acta Med Scand* 1970;187:419-28.

- 308 *Skov PE, Hansen HE. Glomerular filtration rate, renal plasma flow and filtration fraction in living donors before and after nephrectomy. *Acta Med Scand* 1974;195:97-103.
- 4013 *Slack TK, Wilson DM. Normal renal function: CIN and CPAH in healthy donors before and after nephrectomy. *Mayo Clin Proc* 1976;51:296-300.
- 2306 *Smith CL, Hampton EM. Using estimated creatinine clearance for individualizing drug therapy: a reassessment. *DICP* 1990;24:1185-90.
- 13272 *Soulsby N, Greville H, Coulthard K, Doecke C. Renal dysfunction in cystic fibrosis: Is there cause for concern? *Pediatric Pulmonology* 2009;44:947-953.
- 157 *Speck B. Determination of renal clearances with radioactive diatrizoate, EDTA and o-l-hippuric acid using the single injection technique. *Helv Med Acta* 1969;34:486-97.
- 1577 *Spigt MG, Knottnerus JA, Westerterp KR, Olde Rikkert MG, Schayck CP. The effects of 6 months of increased water intake on blood sodium, glomerular filtration rate, blood pressure, and quality of life in elderly (aged 55-75) men. *J Am Geriatr Soc* 2006;54:438-43.
- 4237 *Stamp TC. 51Cr-EDTA clearance and G.F.R. *Lancet* 1968;2:1348.
- 13353 *Stamp TC, Stacey TE, Rose GA. Comparison of glomerular filtration rate measurements using inulin, 51CrEDTA, and a phosphate infusion technique. *Clin Chim Acta* 1970;30:351-8.
- 3025 *Starlinger H, Berghoff A. [Studies on the usefulness of endogenous creatinine clearance in comparison to inulin clearance in experiments during rest and exertion on healthy subjects.]. *Int Z Angew Physiol* 1962;19:194-200.
- 310 *Steinhoff H, Pabst HW, Heinze HG, Hor G, Hadid D. [Chloriodopropyl-inulin-131-I for the quantitative determination of the glomerular filtrate]. *Nucl Med (Stuttg)* 1967;321-8.
- 1582 *Steinstrasser A, Berberich R, Wilhelm H, Curschmann J. [Age dependence of kidney function]. *Urologe A* 1989;28:106-11.
- 2312 *Stejskal J, Novakova J, Reitschlagrova V. [Validity of routinely used methods of testing glomerular filtration in pediatric practice]. *Cesk Pediatr* 1990;45:662-6.
- 1585 *Stenvinkel P, Saggarr-Malik AK, Alvestrand A. Renal haemodynamics and tubular sodium handling following volume expansion with sodium chloride (NaCl) and glucose in healthy humans. *Scand J Clin Lab Invest* 1992;52:837-46.
- 206 *Stern G, Frennby B, Mansson S, Nyman U, Van Westen D, Almen T. Determining 'true' glomerular filtration rate in healthy adults using infusion of inulin and comparing it with values obtained using other clearance techniques or prediction equations. *Scand J Urol Nephrol* 2008;42:278-85.
- 13385 *Stevens LA, Levey AS. Measured GFR as a confirmatory test for estimated GFR. *J Am Soc Nephrol* 2009;20:2305-13.
- 82 *Stribrna J, Oppelt A, Woller P, Franke WG, Jirickova E, Janata V, et al. [Glomerular filtration measured by plasma clearance of 169Yb-DTPA in kidney recipients. Comparison with the clearance of inulin, polyfructosan S and creatinine]. *Cas Lek Cesk* 1986;125:1070-2.
- 13462 *Stribrna J, Skodova Z, Kotik L. Inulin and creatinine clearance as a measure of glomerular filtration after operation with extracorporeal circulation. *Casopis Lekarů Ceskych* 1985;124:1622-1624.
- 13709 *Swabb EA, Sugerman AA, Frantz M, Platt TB, Stern M. Renal handling of the monobactam aztreonam in healthy subjects. *Clin Pharmacol Ther* 1983;33:609-14.
- 13452 *Swan SK, Rudy DW, Lasseter KC, Ryan CF, Buechel KL, Lambrecht LJ, et al. Effect of cyclooxygenase-2 inhibition on renal function in elderly persons receiving a low-salt diet. A randomized, controlled trial. *Annals of internal medicine* 2000;133:1-9.

2327 *Svenningsen NW. Single injection polyfructosan clearance in normal and asphyxiated neonates. *Acta Paediatr Scand* 1975;64:87-95.

13699 *Swinkels DW, Hendriks JC, Nauta J, de Jong MC. Glomerular filtration rate by single-injection inulin clearance: definition of a workable protocol for children. *Ann Clin Biochem* 2000;37 (Pt 1):60-6.

83 *Sziklas JJ, Hosain F, Reba RC, Wagner HN, Jr. Comparison of ¹⁶⁹Yb-DTPA, ^{113m}In-DTPA, ¹⁴C-insulin and endogenous creatinine to estimate glomerular filtration. *J Nucl Biol Med* 1971;15:122-5.

2333 *Takahira R, Yonemura K, Yonekawa O, Iwahara K, Kanno T, Fujise Y, et al. Tryptophan glycoconjugate as a novel marker of renal function. *Am J Med* 2001;110:192-7.

84 *Tauxe WN. Determination of glomerular filtration rate by single-plasma sampling technique following injection of radioiodinated diatrizoate. *J Nucl Med* 1986;27:45-50.

2338 *Teplan V, Schuck O, Horackova M, Skibova J, Holecek M. Effect of a keto acid-amino acid supplement on the metabolism and renal elimination of branched-chain amino acids in patients with chronic renal insufficiency on a low protein diet. *Wien Klin Wochenschr* 2000;112:876-81.

13716 *Tilstone WJ, Semple PF, Lawson DH, Boyle JA. Effects of furosemide on glomerular filtration rate and clearance of practolol, digoxin, cephaloridine, and gentamicin. *Clin Pharmacol Ther* 1977;22:389-94.

1616 *Toffaletti JG, McDonnell EH. Variation of serum creatinine, cystatin C, and creatinine clearance tests in persons with normal renal function. *Clin Chim Acta* 2008;395:115-9.

85 *Tomlanovich S, Golbetz H, Perlroth M, Stinson E, Myers BD. Limitations of creatinine in quantifying the severity of cyclosporine-induced chronic nephropathy. *Am J Kidney Dis* 1986;8:332-7.

2353 *Toto R, Shultz P, Raj L, Mitchell H, Shaw W, Ramjit D, et al. Efficacy and tolerability of losartan in hypertensive patients with renal impairment. Collaborative Group. *Hypertension* 1998;31:684-91.

2355 *Toto RD. Conventional measurement of renal function utilizing serum creatinine, creatinine clearance, inulin and para-aminohippuric acid clearance. *Curr Opin Nephrol Hypertens* 1995;4:505-9; discussion 503-4.

13297 *Toto RD. Conventional measurement of renal function utilizing serum creatine, creatine clearance, inulin and para-aminohippuric acid clearance. *Current Opinion in Nephrology and Hypertension* 1995;4:505-509.

160 *Traub YM, Sameul R, Lubin E, Lewitus Z, Rosenfeld JB. A comparison between the clearances of inulin, endogenous creatinine and ⁵¹Cr-EDTA. *Isr J Med Sci* 1973;9:487-9.

314 *Traynor J, Mactier R, Geddes CC, Fox JG. How to measure renal function in clinical practice. *BMJ* 2006;333:733-7.

13531 *Truniger B, Donath A, Kappeler M. Simplified clearance techniques. The single injection method and its modifications. *Helvetica medica acta* 1968;34:116-129.

4244 *Tsubaki T, Goodin S, Leader WG, Chandler MH. Estimation of creatinine clearance in patients with gynecologic cancer. *Clin Pharm* 1993;12:685-90.

2365 *Tufro A, Arrizurieta EE, Repetto H. Renal functional reserve in children with a previous episode of haemolytic-uraemic syndrome. *Pediatr Nephrol* 1991;5:184-8.

161 *Tvedegaard E, Madsen S. Alphacalcidol and renal function in normal subjects. *Miner Electrolyte Metab* 1988;14:158-62.

13440 *Uegaki A, Sasaki M, Nakajima K, Tobari M, Ushio Y, Kino K. Cystatin C-based assessment of renal function is more reliable compared to creatine-based assessment in cirrhotic patients. *Hepatology International* 2010;4:259.

2368 *Uemura O, Yamada T, Nagai T, Fujita N. [Chronic kidney disease in children]. *Nippon Rinsho* 2008;66:1814-20.

- 2371 *Walczyk MH, Pulliam J, Bennett WM. Effects of hyperglycemia and mannitol infusions on renal hemodynamics in normal subjects. *Am J Med Sci* 1990;300:218-24.
- 2376 *van Acker BA, Koomen GC, Koopman MG, de Waart DR, Arisz L. Creatinine clearance during cimetidine administration for measurement of glomerular filtration rate. *Lancet* 1992;340:1326-9.
- 919 *van Acker BA, Koomen GC, Koopman MG, Krediet RT, Arisz L. Discrepancy between circadian rhythms of inulin and creatinine clearance. *J Lab Clin Med* 1992;120:400-10.
- 2378 *van den Anker JN, de Groot R, Broerse HM, Sauer PJ, van der Heijden BJ, Hop WC, et al. Assessment of glomerular filtration rate in preterm infants by serum creatinine: comparison with inulin clearance. *Pediatrics* 1995;96:1156-8.
- 2379 *van den Berg JG, Koopman MG, Arisz L. Ranitidine has no influence on tubular creatinine secretion. *Nephron* 1996;74:705-8.
- 1642 *van der Heijden AJ, Frohn-Mulder IM. [Glomerular filtration in premature infants]. *Tijdschr Kindergeneesk* 1985;53:181-5.
- 2381 *van Olden RW, Krediet RT, Struijk DG, Arisz L. Measurement of residual renal function in patients treated with continuous ambulatory peritoneal dialysis. *J Am Soc Nephrol* 1996;7:745-50.
- 2382 *van Olden RW, van Acker BA, Koomen GC, Krediet RT, Arisz L. Time course of inulin and creatinine clearance in the interval between two haemodialysis treatments. *Nephrol Dial Transplant* 1995;10:2274-80.
- 2383 *van Olden RW, van Acker BA, Koomen GC, Krediet RT, Arisz L. Contribution of tubular anion and cation secretion to residual renal function in chronic dialysis patients. *Clin Nephrol* 1998;49:167-72.
- 13433 *van Pottelbergh G, van Heden L, Mathei C, Degryse J. Methods to evaluate renal function in elderly patients: A systematic literature review. *Age and Ageing* 2010;39:542-548.
- 13698 *van Rossum LK, Cransberg K, de Rijke YB, Zietse R, Lindemans J, Vulto AG. Determination of inulin clearance by single injection or infusion in children. *Pediatr Nephrol* 2005;20:777-81.
- 1644 *van Rossum LK, Mathot RA, Cransberg K, Vulto AG. Optimal sampling strategies to assess inulin clearance in children by the inulin single-injection method. *Clin Chem* 2003;49:1170-9.
- 207 *van Westen D, Almen T, Chai CM, Frennby B, Mansson S, Sterner G. Biliary and total extrarenal clearance of inulin and iohexol in pigs. A source of error when determining gfr as body clearance. *Nephron* 2002;91:300-7.
- 2388 *Vargo DL, Brater DC, Rudy DW, Swan SK. Dopamine does not enhance furosemide-induced natriuresis in patients with congestive heart failure. *J Am Soc Nephrol* 1996;7:1032-7.
- 2392 *Weekers L, Scheen AJ, Godon JP. [How I examine...diabetic nephropathy. Second part: assessment of glomerular filtration]. *Rev Med Liege* 1998;53:571-5.
- 3109 *Vere DW, Walduck A. ENDOGENOUS-CREATININE CLEARANCE AND GLOMERULAR FILTRATION-RATE. *Lancet* 1964;2:1299.
- 1655 *Vervoort G, Willems HL, Wetzels JF. Assessment of glomerular filtration rate in healthy subjects and normoalbuminuric diabetic patients: validity of a new (MDRD) prediction equation. *Nephrol Dial Transplant* 2002;17:1909-13.
- 13371 *Wesson L. Renal hemodynamics in physiological states. In: *Physiology of the Human Kidney*. New York & London: Grune & Stratton; 1969.
- 317 *Westenfelder SR, Naber KG, Madsen PO. Pharmacokinetics of a new cephalosporin, cephacetrile, in patients with normal and impaired renal function. *Infection* 1973;1:157-62.
- 2397 *Wetzels JF, Hoitsma AJ, Berden JH, Koene RA. Renal hemodynamic effects of a short-term high protein and low protein diet in patients with renal disease. *Clin Nephrol* 1988;30:42-7.

- 86 *Wharton WW, 3rd, Sondeen JL, McBiles M, Gradwohl SE, Wade CE, Ciceri DP, et al. Measurement of glomerular filtration rate in ICU patients using 99mTc-DTPA and inulin. *Kidney Int* 1992;42:174-8.
- 1661 *White AJ, Strydom WJ. Normalisation of glomerular filtration rate measurements. *Eur J Nucl Med* 1991;18:385-90.
- 2401 *White CA, Huang D, Akbari A, Garland J, Knoll GA. Performance of creatinine-based estimates of GFR in kidney transplant recipients: a systematic review. *Am J Kidney Dis* 2008;51:1005-15.
- 13656 *White MT, Diebel LN, Ledgerwood AM, Lucas CE. The significance of a serum creatinine in defining renal function in seriously injured and septic patients. *J Trauma*. 2011;70:421-7.
- 8348 *Wibell L, Evrin PE, Berggard I. Serum 2-microglobulin in renal disease. *Nephron* 1973;10:320-31.
- 87 *Wickre CG, Bennett WM. Renal cyst epithelial transport in non-uremic polycystic kidney disease. *Kidney Int* 1983;23:514-8.
- 8354 *Widstam-Attorps U, Berg U. Urinary protein excretion and renal function in young people with diabetes mellitus. *Nephrol Dial Transplant* 1992;7:487-92.
- 938 *Wiegmann TB, Zlomke AM, MacDougall ML, Kipp DE. Controlled changes in chronic dietary protein intake do not change glomerular filtration rate. *Am J Kidney Dis* 1990;15:147-54.
- 2408 *Wilkins BH. A reappraisal of the measurement of glomerular filtration rate in pre-term infants. *Pediatr Nephrol* 1992;6:323-7.
- 941 *Wilkinson J, Fleming JS, Waller DG. Effect of food and activity on the reproducibility of isotopic GFR estimation. *Nucl Med Commun* 1990;11:697-700.
- 2409 *Willems HL, Hilbrands LB, van de Calseyde JF, Monnens LA, Swinkels DW. Is serum cystatin C the marker of choice to predict glomerular filtration rate in paediatric patients? *Ann Clin Biochem* 2003;40:60-4.
- 13647 *Vincent C, Pozet N, Revillard JP. Plasma (beta)(2) microglobulin turnover in renal insufficiency. *Acta Clinica Belgica* 1980;35:2-13.
- 2411 *Windstetter D, Schaefer F, Scharer K, Reiter K, Eife R, Harms HK, et al. Renal function and renotropic effects of secretin in cystic fibrosis. *Eur J Med Res* 1997;2:431-6.
- 162 *Winterborn MH, Beetham R, White RH. Comparison of plasma disappearance and standard clearance techniques for measuring glomerular filtration rate in children with and without vesico-ureteric reflux. *Clin Nephrol* 1977;7:262-70.
- 13719 *Vogeli B, Riedwyl H, Donath A, Oetliker O. Comparison of glomerular filtration rate and effective renal plasma flow determinations obtained by a single injection technique and by means of a standard clearance technique in children. *Acta Paediatr Scand* 1971;60:528-32.
- 1671 *Woitas RP, Stoffel-Wagner B, Flommersfeld S, Poege U, Schiedermaier P, Klehr HU, et al. Correlation of serum concentrations of cystatin C and creatinine to inulin clearance in liver cirrhosis. *Clin Chem* 2000;46:712-5.
- 1674 *Vora JP, Burch A, Owens DR, Peters JR. Simultaneous determination of glomerular filtration rate and effective renal plasma flow. *Clin Phys Physiol Meas* 1991;12:269-77.
- 166 *Vorburger C, Riedwyl H, Reubi F. [Comparative studies on the renal clearance of Na-Cr²Cr⁵¹ ethylenediaminetetraacetate, inulin and sodium thiosulfate in humans]. *Klin Wochenschr* 1969;47:415-20.
- 89 *Wustenberg PW, Roman W, Bottcher M, Neumann J, Esther G, Unterspann S. [Comparative estimation of monocompartmental methods of analyzing 99m-Tc-diethylenetriamine pentaacetic acid clearance by declining plasma levels with special reference to long term studies]. *Biomed Biochim Acta* 1983;42:547-59.

8438 *Wuyts B, Bernard D, Van den Noortgate N, Van de Walle J, Van Vlem B, De Smet R, et al. Reevaluation of formulas for predicting creatinine clearance in adults and children, using compensated creatinine methods. *Clin Chem* 2003;49:1011-4.

2428 *Zarowitz BJ, Robert S, Peterson EL. Prediction of glomerular filtration rate using aminoglycoside clearance in critically ill medical patients. *Ann Pharmacother* 1992;26:1205-10.

13365 *Zdolsek JH, Lisander B, Hahn RG. Measuring the size of the extracellular fluid space using bromide, iohexol, and sodium dilution. *Anesth Analg* 2005;101:1770-7.

2429 *Zeiler M, Bock A, Zurcher R, Gratwohl A, Thiel G. Plasma creatinine for the evaluation of recovery of glomerular filtration rate after discontinuation of cyclosporine A in bone marrow transplantation: results of a prospective study. *Transplant Proc* 1994;26:2590-3.

3815 *Zimmermann R. [Reference values in pregnancy]. *Praxis (Bern 1994)* 2003;92:331-6.

210 *Zurth C. Mechanism of renal excretion of various X-ray contrast materials in rabbits. *Invest Radiol* 1984;19:110-5.

Kapitel 3.2

Record nr, artikel, författare mm

14104 K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002;39:S1-266.

4265 *Abbink FC, Laarman CA, Braam KI, van Wijk JA, Kors WA, Bouman AA, et al. Beta-trace protein is not superior to cystatin C for the estimation of GFR in patients receiving corticosteroids. *Clin Biochem* 2008;41:299-305.

12864 *Abrahamsson K, Jodal U, Sixt R, Olsson I, Sillen U. Estimation of renal function in children and adolescents with spinal dysraphism. *J Urol.* 2008;179:2407-9.

4274 *Accetta NA, Gladstone EH, DiSogra C, Wright EC, Briggs M, Narva AS. Prevalence of estimated GFR reporting among US clinical laboratories. *Am J Kidney Dis* 2008;52:778-87.

3148 *Adebisi SA, Adekunle BA, Etu AK. Creatinine clearance: alternative approach to traditional 24-hour urine collection in normal individuals. *Afr J Med Med Sci* 2001;30:27-30.

4296 *Agarwal R. Estimating GFR from serum creatinine concentration: pitfalls of GFR-estimating equations. *Am J Kidney Dis* 2005;45:610-3.

1709 *Ahmed SB, Bentley-Lewis R, Hollenberg NK, Graves SW, Seely EW. A Comparison of Prediction Equations for Estimating Glomerular Filtration Rate in Pregnancy. *Hypertens Pregnancy* 2009:1-13.

4317 *Aksun SA, Ozmen D, Ozmen B, Parildar Z, Mutaf I, Turgan N, et al. Beta2-microglobulin and cystatin C in type 2 diabetes: assessment of diabetic nephropathy. *Exp Clin Endocrinol Diabetes* 2004;112:195-200.

971 *Al-Khader AA, Tamim H, Sulaiman MH, Jondeby MS, Taher S, Hejaili FF, et al. What is the most appropriate formula to use in estimating glomerular filtration rate in adult Arabs without kidney disease? *Ren Fail* 2008;30:205-8.

13683 *Alper AB, Yi Y, Rahman M, Webber LS, Magee L, von Dadelszen P, et al. Performance of Estimated Glomerular Filtration Rate Prediction Equations in Preeclamptic Patients. *Am J Perinatol* 2010.

1713 *Anastasio P, Spitali L, Frangiosa A, Molino D, Stellato D, Cirillo E, et al. Glomerular filtration rate in severely overweight normotensive humans. *Am J Kidney Dis* 2000;35:1144-8.

4364 *Anders HJ, Rihl M, Vielhauer V, Schattenkirchner M. Assessment of renal function in rheumatoid arthritis: validity of a new prediction method. *J Clin Rheumatol* 2002;8:130-3.

4365 *Andersen S, van Nieuwenhoven FA, Tarnow L, Rossing P, Rossing K, Wieten L, et al. Reduction of urinary connective tissue growth factor by Losartan in type 1 patients with diabetic nephropathy. *Kidney Int* 2005;67:2325-9.

12878 *Andersen TB, Erlandsen EJ, Frokiaer J, Eskild-Jensen A, Brochner-Mortensen J. Comparison of within- and between-subject variation of serum cystatin C and serum creatinine in children aged 2-13 years. *Scand J Clin Lab Invest.* 2010.

4366 *Andersen TB, Eskild-Jensen A, Frokiaer J, Brochner-Mortensen J. Measuring glomerular filtration rate in children; can cystatin C replace established methods? A review. *Pediatr Nephrol* 2009;24:929-41.

14023 *Andersen TB, Jodal L, Boegsted M, Erlandsen EJ, Morsing A, Frokiaer J, et al. GFR Prediction From Cystatin C and Creatinine in Children: Effect of Including Body Cell Mass. *American Journal of Kidney Diseases* 2011.

14024 *Anderson J, Glynn LG. Definition of chronic kidney disease and measurement of kidney function in original research papers: A review of the literature. *Nephrology Dialysis Transplantation* 2011;26:2793-2798.

- 1716 *Ansquer JC, Dalton RN, Causse E, Crimet D, Le Malicot K, Foucher C. Effect of fenofibrate on kidney function: a 6-week randomized crossover trial in healthy people. *Am J Kidney Dis* 2008;51:904-13.
- 13481 *Apakkan Aksun S, Ozmen D, Ozmen B, Parildar Z, Mutaf I, Turgan N, et al. (beta)2-microglobulin and cystatin C in type 2 diabetes: Assessment of diabetic nephropathy. *Experimental and Clinical Endocrinology and Diabetes* 2004;112:195-200.
- 982 *Apperloo JJ, Gerlag PG, Beerenhout CH, Vader HL. [Estimating renal function based on creatinine clearance: application of different formulas and correction for obese patients]. *Ned Tijdschr Geneesk* 2007;151:1016-23.
- 170 *Artunc FH, Fischer IU, Risler T, Erley CM. Improved estimation of GFR by serum cystatin C in patients undergoing cardiac catheterization. *Int J Cardiol* 2005;102:173-8.
- 4419 *Assadi M, Eftekhari M, Hozhabrosadati M, Saghari M, Ebrahimi A, Nabipour I, et al. Comparison of methods for determination of glomerular filtration rate: low and high-dose Tc-99m-DTPA renography, predicted creatinine clearance method, and plasma sample method. *Int Urol Nephrol* 2008;40:1059-65.
- 1724 *Assan R, Blanchet F, Feutren G, Timsit J, Larger E, Boitard C, et al. Normal renal function 8 to 13 years after cyclosporin A therapy in 285 diabetic patients. *Diabetes Metab Res Rev* 2002;18:464-72.
- 13673 *Aucella F, Guida CC, Lauriola V, Vergura M. How to assess renal function in the geriatric population. *J Nephrol*. 2010;23 Suppl 15:S46-54.
- 994 *Aydn F, Gungor F, Cengiz AK, Tuncer M, Mahsereci E, Ozdem S, et al. Comparison of glomerular filtration rate measurements with the two plasma sample and single plasma sample, gamma camera Gates, creatinine clearance, and prediction equation methods in potential kidney donors with normal renal function. *Nucl Med Commun* 2008;29:157-65.
- 12885 *Aydn F, Tezcan G, Gungor O, Cengiz AK, Hazar V, Akman S, et al. Can serum cystatin C reflect the glomerular filtration rate accurately in pediatric patients under chemotherapeutic treatment? A comparative study with Tc-99m DTPA two-plasma sample method. *Nucl Med Commun*. 2010.
- 13482 *Aygün B, Mortier NA, Smeltzer MP, Hankins JS, Ware RE. Glomerular hyperfiltration and microalbuminuria in children with sickle cell anemia. *SOURCE Blood* (2009) 114:22. Date of Publication: 20 Nov 2009 CONFERENCE NAME 51st Annual Meeting of the American Society of Hematology, ASH CONFERENCE LOCATION New Orleans, LA CONFERENCE DATE 2009-12-05 to 2009-12-08.
- 13583 *Bacchetta J, Cochat P, Rognant N, Ranchin B, Hadj-Aissa A, Dubourg L. Which Creatinine and Cystatin C Equations Can Be Reliably Used in Children? *Clin J Am Soc Nephrol*. 2011;6:552-560.
- 4464 *Bakoush O, Grubb A, Rippe B. Inaccuracy of GFR predictions by plasma cystatin C in patients without kidney dysfunction and in advanced kidney disease. *Clin Nephrol* 2008;69:331-8.
- 14025 *Ballester-Asensio E, Navarro-Ruiz A, Martinez-Pons L, Vallecillo-Hernandez J, Del Valle-Perez R, Codoner-Franch P. Cystatin C in children with severe obesity: Relationship with metabolic risk factors and body composition. *International Journal of Obesity* 2011;35:S148.
- 355 *Banfi G, Del Fabbro M, Lippi G. Serum creatinine concentration and creatinine-based estimation of glomerular filtration rate in athletes. *Sports Med* 2009;39:331-7.
- 13584 *Bargnoux AS, Cavalier E, Cristol JP, Simon N, Dupuy AM, Garrigue V, et al. Cystatin C is a reliable marker for estimation of glomerular filtration rate in renal transplantation: validation of a new turbidimetric assay using monospecific sheep antibodies. *Clin Chem Lab Med*. 2011;49:265-270.
- 13832 *Bargnoux AS, Perrin M, Garrigue V, Badiou S, Dupuy AM, Mourad G, et al. Analytical performances of cystatin C turbidimetric assay: which impact on accuracy of glomerular filtration rate estimation in renal transplantation? *Clin Chem Lab Med* 2011.

14026 *Bargnoux AS, Simon N, Garrigue V, Dupuy AM, Badiou S, Mourad G, et al. Accuracy of GFR predictive equations using ID-MS traceable creatinine: Compensated Jaffe and enzymatic. *Clinical Laboratory* 2011;57:357-361.

13483 *Barlow AD, Taylor AH, Elwell R, Buttress AS, Moorhouse J, Nicholson ML. The performance of three estimates of glomerular filtration rate before and after live donor nephrectomy. *Transplant International* 2010;23:417-423.

4492 *Barracough LH, Field C, Wieringa G, Swindell R, Livsey JE, Davidson SE. Estimation of renal function -- what is appropriate in cancer patients? *Clin Oncol (R Coll Radiol)* 2008;20:721-6.

13484 *Barri Y, Parker T, Daoud Y, Glasscock R. Age, proteinuria, and low iothalamate GFR (iGFR) before donation predict stage 3 CKD post-kidney donation. SOURCE American Journal of Transplantation (2009) 9 (372) CONFERENCE NAME American Transplant Congress 2009 CONFERENCE LOCATION Boston, MA CONFERENCE DATE 2009-05-30 to 2009-06-03.

13485 *Barri YM, Parker T, Daoud Y, Glasscock RJ. Definition of chronic kidney disease after uninephrectomy in living donors: What are the implications? *Transplantation* 2010;90:575-580.

4497 *Barroso S, Martinez JM, Martin MV, Rayo I, Caravaca F. [Accuracy of indirect estimates of renal function in advanced chronic renal failure patients]. *Nefrologia* 2006;26:344-50.

4529 *Beauvieux MC, Le Moigne F, Lasseur C, Raffaitin C, Perlemoine C, Barthe N, et al. New predictive equations improve monitoring of kidney function in patients with diabetes. *Diabetes Care* 2007;30:1988-94.

4535 *Beddhu S, Samore MH, Roberts MS, Stoddard GJ, Pappas LM, Cheung AK. Creatinine production, nutrition, and glomerular filtration rate estimation. *J Am Soc Nephrol* 2003;14:1000-5.

1745 *Bello E, Caramelo C. [Increase of tubular secretion of creatinine by L-arginine: mechanism of practical significance in the assessment of renal function based on creatinine clearance]. *Nefrologia* 2000;20:517-22.

1750 *Benohr P, Grenz A, Hartmann JT, Muller GA, Blaschke S. Cystatin C--a marker for assessment of the glomerular filtration rate in patients with cisplatin chemotherapy. *Kidney Blood Press Res* 2006;29:32-5.

13486 *Benöhr P, Grenz A, Hartmann JT, Müller GA, Blaschke S. Cystatin C--a marker for assessment of the glomerular filtration rate in patients with cisplatin chemotherapy. In: *Kidney & blood pressure research*; 2006. p 32-5.

13392 *Berg UB, Back R, Celsi G, Halling SE, Homberg I, Krmár RT, et al. Comparison of Plasma Clearance of Iohexol and Urinary Clearance of Inulin for Measurement of GFR in Children. *Am J Kidney Dis* 2010;57:55-61.

1754 *Berg UB, Ericzon BG, Nemeth A. Renal function before and long after liver transplantation in children. *Transplantation* 2001;72:631-7.

4578 *Beringer PM, Hidayat L, Heed A, Zheng L, Owens H, Benitez D, et al. GFR estimates using cystatin C are superior to serum creatinine in adult patients with cystic fibrosis. *J Cyst Fibros* 2009;8:19-25.

13586 *Beringer PM, Owens H, Nguyen A, Mordwinkin N, Louie S, Mak M, et al. Estimation of glomerular filtration rate by using serum cystatin C and serum creatinine concentrations in patients with human immunodeficiency virus. *Pharmacotherapy*. 2010;30:1004-10.

14027 *Bevc S, Hojs R, Ekart R. Creatinine-based formula (MDRD) versus cystatin c-based formula (simple cystatin C formula) for estimation of GFR in patients with chronic kidney disease. *European Journal of Internal Medicine* 2011;22:S12.

13844 *Bevc S, Hojs R, Ekart R, Gorenjak M, Puklavec L. Simple cystatin C formula compared to sophisticated CKD-EPI formulas for estimation of glomerular filtration rate in the elderly. *Ther Apher Dial* 2011;15:261-8.

13487 *Bevc S, Završnik M, Hojs R, Ekart R, Gorenjak M, Puklavec L. Cystatin C formula compared to creatinine formulas for estimation of glomerular filtration rate in overweight patients with diabetes mellitus type 2 and chronic kidney disease. SOURCE *Diabetes* (2009) 58 CONFERENCE NAME 69th Annual Meeting of the American

Diabetes Association CONFERENCE LOCATION New Orleans, LA CONFERENCE DATE 2009-06-05 to 2009-06-09.

13845 *Bhavsar NA, Appel LJ, Kusek JW, Contreras G, Bakris G, Coresh J, et al. Comparison of Measured GFR, Serum Creatinine, Cystatin C, and Beta-Trace Protein to Predict ESRD in African Americans With Hypertensive CKD. *Am J Kidney Dis* 2011.

4598 *Bianchi C, Donadio C, Tramonti G, Consani C, Lorusso P, Rossi G. Reappraisal of serum beta2-microglobulin as marker of GFR. *Ren Fail* 2001;23:419-29.

4601 *Biancofiore G, Pucci L, Cerutti E, Penno G, Pardini E, Esposito M, et al. Cystatin C as a marker of renal function immediately after liver transplantation. *Liver Transpl* 2006;12:285-91.

4 *Biglarnia AR, Wadstrom J, Larsson A. Decentralized glomerular filtration rate (GFR) estimates in healthy kidney donors show poor correlation and demonstrate the need for improvement in quality and standardization of GFR measurements in Sweden. *Scand J Clin Lab Invest* 2007;67:227-35.

4619 *Bird NJ, Peters C, Michell AR, Peters AM. Comparison between slope-intercept and slope-only techniques for measuring glomerular filtration rate: use of two independent markers and an independent arbiter. *Nucl Med Commun* 2007;28:711-8.

1022 *Bird NJ, Peters C, Michell AR, Peters AM. Using the slope-only technique and estimated glomerular filtration rate for checking the reliability of slope-intercept measurement of glomerular filtration rate. *Nucl Med Commun* 2008;29:1086-92.

4621 *Bird NJ, Peters C, Michell AR, Peters AM. Reliability of the MDRD method for estimating glomerular filtration rate in relation to gender, body mass index and extracellular fluid volume. *Eur J Clin Invest* 2008;38:486-93.

4622 *Bird NJ, Peters C, Michell AR, Peters AM. Association between glomerular filtration rate and extracellular fluid volume in normal subjects and patients with renal impairment. *Scand J Clin Lab Invest* 2008;68:39-49.

33 *Björk J, Bäck SE, Sterner G, Carlson J, Lindström V, Bakoush O, et al. Prediction of relative glomerular filtration rate in adults: new improved equations based on Swedish Caucasians and standardized plasma-creatinine assays. *Scand J Clin Lab Invest* 2007;67:678-95.

4628 *Björk J, Grubb A, Nyman U. Variability in diagnostic accuracy can be estimated using simple population weighting. *J Clin Epidemiol* 2009;62:54-7.

13671 *Björk J, Grubb A, Sterner G, Nyman U. Revised equations for estimating glomerular filtration rate based on the Lund-Malmö Study cohort. *Scand J Clin Lab Invest* 2011;71:232-239.

13846 *Björk J, Grubb A, Sterner G, Nyman U. Revised equations for estimating glomerular filtration rate based on the Lund-Malmö Study cohort. *Scand J Clin Lab Invest* 2011;71:232-9.

13831 *Björk J, Jones I, Nyman U, Sjöström P. Validation of the Lund-Malmö, CKD-EPI and MDRD equations to estimate GFR in a large Swedish clinical population. *Scand J Urol Nephrol* 2011.

13847 *Blufpand HN, Tromp J, Abbink FC, Stoffel-Wagner B, Bouman AA, Schouten-van Meeteren AY, et al. Cystatin C more accurately detects mildly impaired renal function than creatinine in children receiving treatment for malignancy. *Pediatr Blood Cancer* 2011;57:262-7.

1764 *Bochud M, Elston RC, Maillard M, Bovet P, Schild L, Shamlaye C, et al. Heritability of renal function in hypertensive families of African descent in the Seychelles (Indian Ocean). *Kidney Int* 2005;67:61-9.

13488 *Boer DP, De Rijke YB, Hop WC, Cransberg K, Dorresteijn EM. Reference values for serum creatinine in children younger than 1 year of age. *Pediatric Nephrology* 2010;25:2107-2113.

4650 *Bokarewa M, Abrahamson M, Levshin N, Egesten A, Grubb A, Dahlberg L, et al. Cystatin C binds serum amyloid A, downregulating its cytokine-generating properties. *J Rheumatol* 2007;34:1293-301.

1768 *Bokenkamp A, Domanetzki M, Zinck R, Schumann G, Byrd D, Brodehl J. Cystatin C--a new marker of glomerular filtration rate in children independent of age and height. *Pediatrics* 1998;101:875-81.

4652 *Bokenkamp A, Domanetzki M, Zinck R, Schumann G, Byrd D, Brodehl J. Cystatin C serum concentrations underestimate glomerular filtration rate in renal transplant recipients. *Clin Chem* 1999;45:1866-8.

13674 *Bonjoch A, Bayes B, Riba J, Puig J, Estany C, Perez-Alvarez N, et al. Validation of estimated renal function measurements compared with the isotopic glomerular filtration rate in an HIV-infected cohort. *Antiviral Res.* 2010;88:347-54.

4665 *Bosma RJ, Doorenbos CR, Stegeman CA, van der Heide JJ, Navis G. Predictive performance of renal function equations in renal transplant recipients: an analysis of patient factors in bias. *Am J Transplant* 2005;5:2193-203.

4667 *Bostom AG, Kronenberg F, Ritz E. Predictive performance of renal function equations for patients with chronic kidney disease and normal serum creatinine levels. *J Am Soc Nephrol* 2002;13:2140-4.

1772 *Botev R, Mallie JP, Couchoud C, Schuck O, Fauvel JP, Wetzels JF, et al. Estimating glomerular filtration rate: Cockcroft-Gault and Modification of Diet in Renal Disease formulas compared to renal inulin clearance. *Clin J Am Soc Nephrol* 2009;4:899-906.

13833 *Botev R, Mallie JP, Wetzels JF, Couchoud C, Schuck O. The clinician and estimation of glomerular filtration rate by creatinine-based formulas: current limitations and quo vadis. *Clin J Am Soc Nephrol* 2011;6:937-50.

4674 *Boudville N, Salama M, Jeffrey GP, Ferrari P. The inaccuracy of cystatin C and creatinine-based equations in predicting GFR in orthotopic liver transplant recipients. *Nephrol Dial Transplant* 2009;24:2926-30.

4680 *Bouvet Y, Bouissou F, Coulais Y, Seronie-Vivien S, Tafani M, Decramer S, et al. GFR is better estimated by considering both serum cystatin C and creatinine levels. *Pediatr Nephrol* 2006;21:1299-306.

1044 *Brandt JR, Wong C, Jones DR, Qualls C, McAfee N, Brewer E, et al. Glomerular filtration rate in children with solid tumors: normative values and a new method for estimation. *Pediatr Hematol Oncol* 2003;20:309-18.

1045 *Brandt JR, Wong CS, Hanrahan JD, Qualls C, McAfee N, Watkins SL. Estimating absolute glomerular filtration rate in children. *Pediatr Nephrol* 2006;21:1865-72.

1774 *Branten AJ, Vervoort G, Wetzels JF. Serum creatinine is a poor marker of GFR in nephrotic syndrome. *Nephrol Dial Transplant* 2005;20:707-11.

4729 *Broekroelofs J, Stegeman CA, Navis GJ, de Haan J, van der Bij W, de Boer WJ, et al. Creatinine-based estimation of rate of long term renal function loss in lung transplant recipients. Which method is preferable? *J Heart Lung Transplant* 2000;19:256-62.

13587 *Bronden B, Eyjolfsson A, Blomquist S, Dardashti A, Ederoth P, Bjursten H. Evaluation of cystatin C with iohexol clearance in cardiac surgery. *Acta Anaesthesiol Scand.* 2011;55:196-202.

13834 *Brown MA, Pirabhahar S, Kelly JJ, Mangos GJ, Mackenzie C, McConachie P, et al. Inaccuracies in estimated glomerular filtration rate in one Australian renal centre. *Nephrology (Carlton)* 2011;16:486-94.

14028 *Budisavljevic MN, Nietert PJ, Zhai Y, Dooley MJ, Rajagopalan PR. Long-term effects of kidney donation on renal function and blood pressure in african americans. *Clinical Journal of the American Society of Nephrology* 2011;6:1474-1480.

1791 *Burkhardt H, Bojarsky G, Gladisch R. Diagnostic efficiency of cystatin C and serum creatinine as markers of reduced glomerular filtration rate in the elderly. *Clin Chem Lab Med* 2002;40:1135-8.

1793 *Burkhardt H, Bojarsky G, Gretz N, Gladisch R. Creatinine clearance, Cockcroft-Gault formula and cystatin C: estimators of true glomerular filtration rate in the elderly? *Gerontology* 2002;48:140-6.

- 1795 *Burkhardt H, Hahn T, Gladisch R. Is kidney size a useful predictor of renal function in the elderly? *Clin Nephrol* 2003;59:415-22.
- 1797 *Burkhardt H, Hahn T, Gretz N, Gladisch R. Bedside estimation of the glomerular filtration rate in hospitalized elderly patients. *Nephron Clin Pract* 2005;101:c1-8.
- 14029 *Buron F, Hadj-Aissa A, Dubourg L, Morelon E, Steghens JP, Ducher M, et al. Estimating glomerular filtration rate in kidney transplant recipients: Performance over time of four creatinine-based formulas. *Transplantation* 2011;92:1005-1011.
- 12921 *Cabarkapa V, Stosic Z, Deric M, Vucurevic-Ristic L, Zeravica R, Ilincic B. Serum cystatin C in estimating glomerular filtration rate. *Journal of Medical Biochemistry* 2008;27:46-51.
- 14030 *Cachapuz I, Oliveira E, Cruz E, Alves V. Children with chronic kidney disease: Schwartz formula for estimating glomerular filtration rate is replaced by cystatin c formula. *Clinical Biochemistry* 2011;44:526.
- 13675 *Camargo EG, Soares AA, Detanico AB, Weinert LS, Veronese FV, Gomes EC, et al. The Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation is less accurate in patients with Type 2 diabetes when compared with healthy individuals. *Diabet Med.* 2011;28:90-5.
- 1800 *Campens D, Buntinx F. Selecting the best renal function tests. A meta-analysis of diagnostic studies. *Int J Technol Assess Health Care* 1997;13:343-56.
- 13489 *Carnevale V, Pastore L, Camaioni M, Mellozzi M, Sabatini M, Arietti E, et al. Estimate of renal function in oldest old inpatients by MDRD study equation, Mayo Clinic equation and creatinine clearance. *Journal of nephrology* 2010;23:306-313.
- 13848 *Carter JL, Lane CE, Fan SL, Lamb EJ. Estimation of residual glomerular filtration rate in peritoneal dialysis patients using cystatin C: comparison with 51Cr-EDTA clearance. *Nephrol Dial Transplant* 2011;26:3729-3732.
- 13835 *Carter JL, Stevens PE, Irving JE, Lamb EJ. Estimating glomerular filtration rate: comparison of the CKD-EPI and MDRD equations in a large UK cohort with particular emphasis on the effect of age. *QJM* 2011;104:839-47.
- 13589 *Cha RH, Lee CS, Lim YH, Kim H, Lee SH, Yu KS, et al. Clinical usefulness of serum cystatin C and the pertinent estimation of glomerular filtration rate based on cystatin C. *Nephrology (Carlton)*. 2010;15:768-76.
- 12925 *Chakrabarti S, Chattopadhyay I. Measuring renal function in old age. *Reviews in Clinical Gerontology* 2008;18:257-267.
- 13490 *Chang TI, Chertow GM. GFR estimating equations, CKD prevalence and the public health. *Journal of internal medicine* 2010;267:354-356.
- 1807 *Chantrel F, Agin A, Offner M, Koehl C, Moulin B, Hannedouche T. Comparison of cystatin C versus creatinine for detection of mild renal failure. *Clin Nephrol* 2000;54:374-81.
- 13491 *Chattopadhyay I. Measuring renal function in old age. *Reviews in Clinical Gerontology* 2003;13:297-302.
- 13676 *Cherney DZ, Sochett EB, Dekker MG, Perkins BA. Ability of cystatin C to detect acute changes in glomerular filtration rate provoked by hyperglycaemia in uncomplicated Type 1 diabetes. *Diabet Med.* 2010;27:1358-65.
- 4884 *Cherry RA, Eachempati SR, Hydo L, Barie PS. Accuracy of short-duration creatinine clearance determinations in predicting 24-hour creatinine clearance in critically ill and injured patients. *J Trauma* 2002;53:267-71.
- 14031 *Cheung KL, Soman S, Tamura MK. Special considerations in the management of chronic kidney disease in the elderly. *Dialysis and Transplantation* 2011;40:241-243.

- 4911 *Cholongitas E, Shusang V, Marelli L, Nair D, Thomas M, Patch D, et al. Review article: renal function assessment in cirrhosis - difficulties and alternative measurements. *Aliment Pharmacol Ther* 2007;26:969-78.
- 4918 *Christensson A, Ekberg J, Grubb A, Ekberg H, Lindstrom V, Lilja H. Serum cystatin C is a more sensitive and more accurate marker of glomerular filtration rate than enzymatic measurements of creatinine in renal transplantation. *Nephron Physiol* 2003;94:p19-27.
- 4919 *Christensson AG, Grubb AO, Nilsson JA, Norrgren K, Sterner G, Sundkvist G. Serum cystatin C advantageous compared with serum creatinine in the detection of mild but not severe diabetic nephropathy. *J Intern Med* 2004;256:510-8.
- 4923 *Chudleigh RA, Ollerton RL, Dunseath G, Peter R, Harvey JN, Luzio S, et al. Performance of the revised '175' Modification of Diet in Renal Disease equation in patients with type 2 diabetes. *Diabetologia* 2008;51:1714-8.
- 4924 *Chudleigh RA, Ollerton RL, Dunseath G, Peter R, Harvey JN, Luzio S, et al. Use of cystatin C-based estimations of glomerular filtration rate in patients with type 2 diabetes. *Diabetologia* 2009;52:1274-8.
- 1818 *Cirillo M, Anastasio P, De Santo NG. Relationship of gender, age, and body mass index to errors in predicted kidney function. *Nephrol Dial Transplant* 2005;20:1791-8.
- 4957 *Cole M, Price L, Parry A, Keir MJ, Pearson AD, Boddy AV, et al. Estimation of glomerular filtration rate in paediatric cancer patients using 51CR-EDTA population pharmacokinetics. *Br J Cancer* 2004;90:60-4.
- 1081 *Coll E, Botey A, Alvarez L, Poch E, Quinto L, Saurina A, et al. Serum cystatin C as a new marker for noninvasive estimation of glomerular filtration rate and as a marker for early renal impairment. *Am J Kidney Dis* 2000;36:29-34.
- 14032 *Comak E, Dogan CS, Uslu Gokceoglu A, Akbas H, Dinckan A, Gurkan A, et al. Estimation of GFR in pediatric renal transplant recipients: Which formula is best? *Pediatric Nephrology* 2011;26:1631-1632.
- 4980 *Coresh J, Astor BC, McQuillan G, Kusek J, Greene T, Van Lente F, et al. Calibration and random variation of the serum creatinine assay as critical elements of using equations to estimate glomerular filtration rate. *Am J Kidney Dis* 2002;39:920-9.
- 4981 *Coresh J, Auguste P. Reliability of GFR formulas based on serum creatinine, with special reference to the MDRD Study equation. *Scand J Clin Lab Invest Suppl* 2008;241:30-8.
- 4984 *Coresh J, Stevens LA. Kidney function estimating equations: where do we stand? *Curr Opin Nephrol Hypertens* 2006;15:276-84.
- 4988 *Corrao AM, Lisi G, Di Pasqua G, Guizzardi M, Marino N, Ballone E, et al. Serum cystatin C as a reliable marker of changes in glomerular filtration rate in children with urinary tract malformations. *J Urol* 2006;175:303-9.
- 14033 *Craig AJ, Britten A, Heenan SD, Irwin AG. Significant differences when using MDRD for GFR estimation compared to radionuclide measured clearance. *European Radiology* 2011;21:2211-2217.
- 13628 *Dalton RN, Chetty R, Stuart M, Iacona RB, Swaisland A. Effects of the Src inhibitor saracatinib (AZD0530) on renal function in healthy subjects. *Anticancer Research* 2010;30:2935-2942.
- 14034 *Damman K, van der Harst P, Smilde TDJ, Voors AA, Navis G, van Veldhuisen DJ, et al. Use of cystatin C levels in estimating renal function and prognosis in patients with chronic systolic heart failure. *Heart* 2011.
- 1833 *Daniel JP, Chantrel F, Offner M, Moulin B, Hannedouche T. Comparison of cystatin C, creatinine and creatinine clearance vs. GFR for detection of renal failure in renal transplant patients. *Ren Fail* 2004;26:253-7.
- 13593 *De Francisco ALM, Fernandez E. The importance of glomerular filtration rate (GFR) estimating equations in the care of hospitalized patients. *European Journal of Internal Medicine* 2011;22:120-121.

5066 *de Lemos ML, Hsieh T, Hamata L, Levin A, Swenerton K, Djurdjev O, et al. Evaluation of predictive formulae for glomerular filtration rate for carboplatin dosing in gynecological malignancies. *Gynecol Oncol* 2006;103:1063-9.

13492 *de Lusignan S, Tomson C, Harris K, van Vlymen J, Gallagher H. Creatinine Fluctuation Has a Greater Effect than the Formula to Estimate Glomerular Filtration Rate on the Prevalence of Chronic Kidney Disease. *Nephron - Clinical Practice* 2010;117:c213-c224.

14035 *De Souza V, Selistre L, Rabilloud M, Kassai B, Hadj-Aissa A, Ranchin B, et al. Can we accurately use the nullnewnull Schwartz-formula to estimate glomerular filtration rate in pediatric renal transplant recipients? *Pediatric Nephrology* 2011;26:1583.

14036 *De Souza V, Selistre L, Rabilloud M, Kassai B, Hadj-Aissa A, Ranchin B, et al. Estimating glomerular filtration rate in children and young adults. *Pediatric Nephrology* 2011;26:1635-1636.

13493 *Deacon A. Limitations of estimating kidney function in adults using formulae [3]. *Annals of Clinical Biochemistry* 2006;43:85.

13494 *Delanaye P, Cavalier E, Chapelle JP, Krzesinski JM, Froissart M. Renal extraction of cystatin C [1]. *Nephrology Dialysis Transplantation* 2006;21:3333.

1112 *Delanaye P, Cavalier E, Depas G, Chapelle JP, Krzesinski JM. New data on the intraindividual variation of cystatin C. *Nephron Clin Pract* 2008;108:c246-8.

13495 *Delanaye P, Cavalier E, Froissart M, Krzesinski JM. Reproducibility of GFR measured by chromium-51-EDTA and iohexol. *Nephrology Dialysis Transplantation* 2008;23:4077-4078.

5097 *Delanaye P, Cavalier E, Krzesinski JM, Chapelle JP. Why the MDRD equation should not be used in patients with normal renal function (and normal creatinine values)? *Clin Nephrol* 2006;66:147-8.

5098 *Delanaye P, Cavalier E, Krzesinski JM, Mariat C. Cystatin C-based equations: don't repeat the same errors with analytical considerations. *Nephrol Dial Transplant* 2008;23:1065; author reply 1065-6.

13496 *Delanaye P, Cavalier E, Mariat C, Maillard N, Krzesinski JM. MDRD or CKD-EPI study equations for estimating prevalence of stage 3 CKD in epidemiological studies: which difference? Is this difference relevant? *SOURCE BMC nephrology* (2010) 11 (8).

5101 *Delanaye P, Cavalier E, Radermecker RP, Paquot N, Depas G, Chapelle JP, et al. Cystatin C or creatinine for detection of stage 3 chronic kidney disease in anorexia nervosa. *Nephron Clin Pract* 2008;110:c158-63.

5102 *Delanaye P, Cavalier E, Radermecker RP, Paquot N, Depas G, Chapelle JP, et al. Estimation of GFR by different creatinine- and cystatin-C-based equations in anorexia nervosa. *Clin Nephrol* 2009;71:482-91.

1114 *Delanaye P, Krzesinski JM. The new Mayo Clinic equation for estimating glomerular filtration rate. *Ann Intern Med* 2005;142:679-80; author reply 681.

13836 *Delanaye P, Krzesinski JM. Indexing of Renal Function Parameters by Body Surface Area: Intelligence or Folly? *Nephron Clin Pract* 2011;119:c289-c292.

13849 *Delanaye P, Mariat C, Cavalier E, Maillard N, Krzesinski JM, White CA. Trimethoprim, Creatinine and Creatinine-Based Equations. *Nephron Clin Pract* 2011;119:c187-c194.

13850 *Delanaye P, Mariat C, Maillard N, Krzesinski JM, Cavalier E. Are the creatinine-based equations accurate to estimate glomerular filtration rate in African American populations? *Clin J Am Soc Nephrol* 2011;6:906-12.

5109 *Delanaye P, Nellessen E, Cavalier E, Depas G, Grosch S, Defraigne JO, et al. Is cystatin C useful for the detection and the estimation of low glomerular filtration rate in heart transplant patients? *Transplantation* 2007;83:641-4.

109 *Delanaye P, Nellessen E, Grosch S, Depas G, Cavalier E, Defraigne JO, et al. Creatinine-based formulae for the estimation of glomerular filtration rate in heart transplant recipients. *Clin Transplant* 2006;20:596-603.

5111 *Delanaye P, Pieroni L, Abshoff C, Lutteri L, Chapelle JP, Krzesinski JM, et al. Analytical study of three cystatin C assays and their impact on cystatin C-based GFR-prediction equations. *Clin Chim Acta* 2008;398:118-24.

2 *Delanghe JR. How to estimate GFR in children. *Nephrol Dial Transplant* 2009;24:714-6.

5116 *Demirovic JA, Pai AB, Pai MP. Estimation of creatinine clearance in morbidly obese patients. *Am J Health Syst Pharm* 2009;66:642-8.

5117 *Demirtas S, Bozbas A, Akbay A, Yavuz Y, Karaca L. Diagnostic value of serum cystatin C for evaluation of hepatorenal syndrome. *Clin Chim Acta* 2001;311:81-9.

12948 *Dharnidharka VR, Kwon C, Stevens G. Serum cystatin C is superior to serum creatinine as a marker of kidney function: a meta-analysis. *Am J Kidney Dis.* 2002;40:221-6.

13497 *Didangelos T, Ntemka A, Iliadis F, Makedou A, Margaritidis C, Papanastasiou E, et al. Estimation of glomerular filtration rate in patients with diabetes mellitus type 2: Comparison of CKD EPI equation and cystatin C-based formula. *SOURCE Diabetologia* (2010) 53 (S478) CONFERENCE NAME 46th Annual Meeting of the European Association for the Study of Diabetes, EASD 2010 CONFERENCE LOCATION Stockholm CONFERENCE DATE 2010-09-20 to 2010-09-24 2011.

48 *Donadio C, Consani C, Ardini M, Caprio F, Grassi G, Lucchesi A. Prediction of glomerular filtration rate from body cell mass and plasma creatinine. *Curr Drug Discov Technol* 2004;1:221-8.

5165 *Donadio C, Lucchesi A, Ardini M, Cosio S, Fanucchi A, Gadducci A. Dose individualization can minimize nephrotoxicity due to carboplatin therapy in patients with ovarian cancer. *Ther Drug Monit* 2009;31:63-9.

5166 *Donadio C, Lucchesi A, Ardini M, Cosio S, Gadducci A. Renal impairment in patients with ovarian cancer. *Eur J Obstet Gynecol Reprod Biol* 2003;106:198-202.

5173 *Donahue A, McCune JS, Faucette S, Gillenwater HH, Kowalski RJ, Socinski MA, et al. Measured versus estimated glomerular filtration rate in the Calvert equation: influence on carboplatin dosing. *Cancer Chemother Pharmacol* 2001;47:373-9.

1860 *Donckerwolcke RM, Coppes MJ. Adaptation of renal function after unilateral nephrectomy in children with renal tumors. *Pediatr Nephrol* 2001;16:568-74.

14037 *Dooley M, Poole S, Rischin D. Chemotherapy dose adjustment in cancer patients: Clinical utility of MDRD, CKD-EPI, Cockcroft and Gault and other formulae for renal function estimation. *Journal of Oncology Pharmacy Practice* 2011;17:11.

5177 *Dooley MJ, Singh S, Rischin D. Rounding of low serum creatinine levels and consequent impact on accuracy of bedside estimates of renal function in cancer patients. *Br J Cancer* 2004;90:991-5.

1134 *Douville P, Martel AR, Talbot J, Desmeules S, Langlois S, Agharazii M. Impact of age on glomerular filtration estimates. *Nephrol Dial Transplant* 2009;24:97-103.

13498 *El-Minshawy O, Saber RA, Osman A. 24-hour creatinine clearance reliability for estimation of glomerular filtration rate in different stages of chronic kidney disease. *Saudi journal of kidney diseases and transplantation : an official publication of the Saudi Center for Organ Transplantation, Saudi Arabia* 2010;21:686-693.

1872 *Emara M, Zahran A, Abd El Hady H, Shoker A. How to best define patients with moderate chronic kidney disease. *Nephron Clin Pract* 2008;110:c195-206.

13499 *Eriksen BO, Mathisen UD, Melsom T, Ingebretsen OC, Jenssen TG, Njolstad I, et al. Cystatin C is not a better estimator of GFR than plasma creatinine in the general population. *Kidney Int* 2010;78:1305-11.

13851 *Eriksen BO, Mathisen UD, Melsom T, Ingebretsen OC, Jenssen TG, Njolstad I, et al. The Role of Cystatin C in Improving GFR Estimation in the General Population. *Am J Kidney Dis* 2011.

- 5276 *Erlandsen EJ, Randers E, Kristensen JH. Evaluation of the Dade Behring N Latex Cystatin C assay on the Dade Behring Nephelometer II System. *Scand J Clin Lab Invest* 1999;59:1-8.
- 178 *Erley CM, Bader BD, Berger ED, Vochazer A, Jorzik JJ, Dietz K, et al. Plasma clearance of iodine contrast media as a measure of glomerular filtration rate in critically ill patients. *Crit Care Med* 2001;29:1544-50.
- 5283 *Esnault VL, Brown EA, Apetrei E, Bagon J, Calvo C, DeChatel R, et al. The effects of amlodipine and enalapril on renal function in adults with hypertension and nondiabetic nephropathies: a 3-year, randomized, multicenter, double-blind, placebo-controlled study. *Clin Ther* 2008;30:482-98.
- 5285 *Estorch M, Tembl A, Antonijuan R, Hernandez A, Mari C, Flotats A, et al. Evaluation of renal graft haemodynamia by ⁵¹Cr-EDTA and o-[¹³¹I]iodohippurate: its use in the early diagnosis of glomerular hyperfiltration. *Nucl Med Commun* 2003;24:679-82.
- 13500 *Estrella MM, Astor BC, Kottgen A, Selvin E, Coresh J, Parekh RS. Prevalence of kidney disease in anaemia differs by GFR-estimating method: The Third National Health and Nutrition Examination Survey (1988-94). *Nephrology Dialysis Transplantation* 2010;25:2542-2548.
- 13852 *Fabbian F, Pala M, Scanelli G, Manzato E, Longhini C, Portaluppi F. Estimation of renal function in patients with eating disorders. *Int J Eat Disord* 2011;44:233-7.
- 14038 *Fadrowski JJ, Furth SL. GFR estimation in children: Questions and answers (and Questions). *Clinical Journal of the American Society of Nephrology* 2011;6:1810-1812.
- 13853 *Fadrowski JJ, Neu AM, Schwartz GJ, Furth SL. Pediatric GFR estimating equations applied to adolescents in the general population. *Clin J Am Soc Nephrol* 2011;6:1427-35.
- 5297 *Fadrowski JJ, Pierce CB, Cole SR, Moxey-Mims M, Warady BA, Furth SL. Hemoglobin decline in children with chronic kidney disease: baseline results from the chronic kidney disease in children prospective cohort study. *Clin J Am Soc Nephrol* 2008;3:457-62.
- 13678 *Faluyi OO, Masinghe SP, Hayward RL, Clive S. Accuracy of GFR estimation by the Cockcroft and Gault, MDRD, and Wright equations in Oncology patients with renal impairment. *Med Oncol*. 2011.
- 1885 *Farker K, Merkel U, Schweer H, Haerting J, Madani SF, Eggers R, et al. Effects of short-term treatment with diclofenac-colestyramine on renal function and urinary prostanoid excretion in patients with type-2 diabetes. *Eur J Clin Pharmacol* 2002;58:85-91.
- 5324 *Fehrman-Ekholm I, Duner F, Brink B, Tyden G, Elinder CG. No evidence of accelerated loss of kidney function in living kidney donors: results from a cross-sectional follow-up. *Transplantation* 2001;72:444-9.
- 5325 *Fehrman-Ekholm I, Seeberger A, Bjork J, Sterner G. Serum cystatin C: a useful marker of kidney function in very old people. *Scand J Clin Lab Invest* 2009;69:606-11.
- 1167 *Fehrman-Ekholm I, Skeppholm L. Renal function in the elderly (>70 years old) measured by means of iohexol clearance, serum creatinine, serum urea and estimated clearance. *Scand J Urol Nephrol* 2004;38:73-7.
- 12963 *Filler G. How to measure renal function in children - What is the role of cystatin C? *Current Pediatric Reviews* 2006;2:225-231.
- 12964 *Filler G, Bokenkamp A, Hofmann W, Le Bricon T, Martinez-Bru C, Grubb A. Cystatin C as a marker of GFR--history, indications, and future research. *Clin Biochem*. 2005;38:1-8.
- 5342 *Filler G, Foster J, Acker A, Lepage N, Akbari A, Ehrich JH. The Cockcroft-Gault formula should not be used in children. *Kidney Int* 2005;67:2321-4.
- 5343 *Filler G, Lepage N. Should the Schwartz formula for estimation of GFR be replaced by cystatin C formula? *Pediatr Nephrol* 2003;18:981-5.

- 1168 *Filler G, Priem F, Lepage N, Sinha P, Vollmer I, Clark H, et al. Beta-trace protein, cystatin C, beta(2)-microglobulin, and creatinine compared for detecting impaired glomerular filtration rates in children. *Clin Chem* 2002;48:729-36.
- 1169 *Filler G, Priem F, Vollmer I, Gellermann J, Jung K. Diagnostic sensitivity of serum cystatin for impaired glomerular filtration rate. *Pediatr Nephrol* 1999;13:501-5.
- 51 *Filler G, Sharma AP. How to monitor renal function in pediatric solid organ transplant recipients. *Pediatr Transplant* 2008;12:393-401.
- 1172 *Finney H, Newman DJ, Price CP. Adult reference ranges for serum cystatin C, creatinine and predicted creatinine clearance. *Ann Clin Biochem* 2000;37 (Pt 1):49-59.
- 1899 *Fliser D, Ritz E. Serum cystatin C concentration as a marker of renal dysfunction in the elderly. *Am J Kidney Dis* 2001;37:79-83.
- 5376 *Flodin M, Hansson LO, Larsson A. Evaluation of Dade Behring N Latex Cystatin C reagent on Abbott ci8200. *Ups J Med Sci* 2006;111:209-13.
- 5377 *Flodin M, Jonsson AS, Hansson LO, Danielsson LA, Larsson A. Evaluation of Gentian cystatin C reagent on Abbott Ci8200 and calculation of glomerular filtration rate expressed in mL/min/1.73 m(2) from the cystatin C values in mg/L. *Scand J Clin Lab Invest* 2007;67:560-7.
- 5382 *Fommei E, Bruselli L, Ripoli A, Gimelli A, Ghione S, Giorgetti A, et al. Single-shot cardiorenal scintigraphy with 99mTc-tetrofosmin: a dynamic characterization at rest and during adenosine infusion. *J Nucl Med* 2009;50:1288-95.
- 5388 *Fontserè N, Bonal J, Navarro M, Riba J, Fraile M, Torres F, et al. A comparison of prediction equations for estimating glomerular filtration rate in adult patients with chronic kidney disease stages 4-5. Effect of nutritional status and age. *Nephron Clin Pract* 2006;104:c160-8.
- 5390 *Fontserè N, Esteve V, Saurina A, Pou M, Barba N, Deulofeu R, et al. The search for a new marker of renal function in older patients with chronic kidney disease stages 3-4: usefulness of cystatin C-based equations. *Nephron Clin Pract* 2009;112:c164-70.
- 5391 *Fontserè N, Salinas I, Bonal J, Bayes B, Riba J, Torres F, et al. Are prediction equations for glomerular filtration rate useful for the long-term monitoring of type 2 diabetic patients? *Nephrol Dial Transplant* 2006;21:2152-8.
- 5392 *Ford L, Berg J. Delay in separating blood samples affects creatinine measurement using the Roche kinetic Jaffe method. *Ann Clin Biochem* 2008;45:83-7.
- 5398 *Fotopoulos AD, Katopodis K, Balafa O, Katsaraki A, Kalaitzidis R, Siamopoulos KC. Individual renal function in polycystic kidney disease: a follow-up study. *Clin Nucl Med* 2001;26:518-24.
- 13679 *Francoz C, Prie D, Abdelrazek W, Moreau R, Mandot A, Belghiti J, et al. Inaccuracies of creatinine and creatinine-based equations in candidates for liver transplantation with low creatinine: impact on the model for end-stage liver disease score. *Liver Transpl.* 2010;16:1169-77.
- 5440 *Froissart M, Rossert J, Jacquot C, Paillard M, Houillier P. Predictive performance of the modification of diet in renal disease and Cockcroft-Gault equations for estimating renal function. *J Am Soc Nephrol* 2005;16:763-73.
- 5452 *Furth SL, Cole SR, Moxey-Mims M, Kaskel F, Mak R, Schwartz G, et al. Design and methods of the Chronic Kidney Disease in Children (CKiD) prospective cohort study. *Clin J Am Soc Nephrol* 2006;1:1006-15.
- 5455 *Gabutti L, Ferrari N, Mombelli G, Marone C. Does cystatin C improve the precision of Cockcroft and Gault's creatinine clearance estimation? *J Nephrol* 2004;17:673-8.
- 13502 *Gansevoort RT, De Jong PE. Challenges for the present CKD classification system. *Current Opinion in Nephrology and Hypertension* 2010;19:308-314.

- 14039 *Gao A, Cachat F, Faouzi M, Mosig D, Girardin E, Chehade H. Assessment of glomerular filtration rate in children: From the new revised Schwartz formula to a new generalized formula. *Pediatric Nephrology* 2011;26:1577.
- 14040 *Garcovich M, Tsochatzis E, Georgiadis D, Germani G, Fede G, Davenport A, et al. Reliability of serum creatinine in assessing renal function is diminished in patients with cirrhosis as compared with patients with organic renal disease. *J Hepatol* 2011;54:S68-S69.
- 14041 *Garcovich M, Tsochatzis E, Georgiadis D, Germani G, Fede G, Burroughs AK, et al. Reliability of serum creatinine in assessing renal function is reduced in patients with cirrhosis as compared with patients with organic renal disease. *Gastroenterology* 2011;140:S954.
- 14042 *Garcovich M, Tsochatzis E, Manousou P, Germani G, Fede G, Fatourou E, et al. Reliability of serum creatinine in assessing renal function is reduced in patients with cirrhosis as compared with patients with organic renal disease. *Digestive and Liver Disease* 2011;43:S101.
- 5484 *Garlipp CR, Bottini PV, Queiroz RR, Afaz SH. Estimating glomerular filtration rate by measurement of cystatin C serum concentrations: a comparison with chrome-EDTA clearance. *Clin Chim Acta* 2008;393:125-7.
- 5508 *Gera M, Slezak JM, Rule AD, Larson TS, Stegall MD, Cosio FG. Assessment of changes in kidney allograft function using creatinine-based estimates of glomerular filtration rate. *Am J Transplant* 2007;7:880-7.
- 5511 *Gerhardt T, Poge U, Stoffel-Wagner B, Ahrendt M, Wolff M, Spengler U, et al. Estimation of glomerular filtration rates after orthotopic liver transplantation: Evaluation of cystatin C-based equations. *Liver Transpl* 2006;12:1667-72.
- 5512 *Gerhardt T, Terjung B, Knipper P, Palmedo H, Woitas RP, Kalff J, et al. Renal impairment after liver transplantation - a pilot trial of calcineurin inhibitor-free vs. calcineurin inhibitor sparing immunosuppression in patients with mildly impaired renal function after liver transplantation. *Eur J Med Res* 2009;14:210-5.
- 14043 *Gibson P, Shammas A, Cada M, Licht C, Gupta A. Do nuclear medicine gfr studies contribute to therapeutic decision making in pediatric oncology patients receiving chemotherapy? *Pediatric Blood and Cancer* 2011;56:937-938.
- 1198 *Giles PD, Fitzmaurice DA. Formula estimation of glomerular filtration rate: have we gone wrong? *BMJ* 2007;334:1198-200.
- 13503 *Glasscock RJ. Estimated glomerular filtration rate: Time for a performance review? *Kidney international* 2009;75:1001-1003.
- 5553 *Godfrey T, Cuadrado MJ, Fofi C, Abbs I, Khamashta MA, Nunan T, et al. Chromium-51 ethylenediamine tetraacetic acid glomerular filtration rate: a better predictor than glomerular filtration rate calculated by the Cockcroft-Gault formula for renal involvement in systemic lupus erythematosus patients. *Rheumatology (Oxford)* 2001;40:324-8.
- 12992 *Gourishankar S, Courtney M, Jhangri GS, Cembrowski G, Pannu N. Serum cystatin C performs similarly to traditional markers of kidney function in the evaluation of donor kidney function prior to and following unilateral nephrectomy. *Nephrol Dial Transplant*. 2008;23:3004-9.
- 5597 *Grassi G, Abdelkawy H, Barsotti M, Paleologo G, Tregnaghi C, Rizzo G, et al. Living kidney transplantation: evaluation of renal function and morphology of potential donors. *Transplant Proc* 2009;41:1121-4.
- 1928 *Graves JW. Diagnosis and management of chronic kidney disease. *Mayo Clin Proc* 2008;83:1064-9.
- 5607 *Gretz N, Schock D, Sadick M, Pill J. Bias and precision of estimated glomerular filtration rate in children. *Pediatr Nephrol* 2007;22:167-9.
- 5611 *Gronroos MH, Jahnukainen T, Irjala K, Harkonen R, Hurme S, Mottonen M, et al. Comparison of glomerular function tests in children with cancer. *Pediatr Nephrol* 2008;23:797-803.

- 14044 *Grubb A. Cystatin C- and creatinine-based GFR-prediction equations for children and adults. *Clinical Biochemistry* 2011;44:501-502.
- 14045 *Grubb A. Cystatin C for estimation of renal function in CKD. *Clinical Chemistry and Laboratory Medicine* 2011;49:S28.
- 1215 *Grubb A, Bjork J, Lindstrom V, Sterner G, Bondesson P, Nyman U. A cystatin C-based formula without anthropometric variables estimates glomerular filtration rate better than creatinine clearance using the Cockcroft-Gault formula. *Scand J Clin Lab Invest* 2005;65:153-62.
- 13741 *Grubb A, Blirup-Jensen S, Lindstrom V, Schmidt C, Althaus H, Zegers I. First certified reference material for cystatin C in human serum ERM-DA471/IFCC. *Clin Chem Lab Med* 2010;48:1619-21.
- 12995 *Grubb A, Nordin G. Notable steps in obtaining improved estimates for glomerular filtration rate. *Clinical Chemistry* 2006;52:169-170.
- 5617 *Grubb A, Nyman U, Bjork J, Lindstrom V, Rippe B, Sterner G, et al. Simple cystatin C-based prediction equations for glomerular filtration rate compared with the modification of diet in renal disease prediction equation for adults and the Schwartz and the Counahan-Barratt prediction equations for children. *Clin Chem* 2005;51:1420-31.
- 5618 *Grubb A, Simonsen O, Sturfelt G, Truedsson L, Thysell H. Serum concentration of cystatin C, factor D and beta 2-microglobulin as a measure of glomerular filtration rate. *Acta Med Scand* 1985;218:499-503.
- 12996 *Gruev T, Chakalarovski K, Stojceva-Taneva O, Grueva A, Trencova K. Effects of glucocorticoid immunosuppression on serum cystatin C levels. *Journal of Medical Biochemistry* 2009;28:191-196.
- 5671 *Hallan S, Asberg A, Lindberg M, Johnsen H. Validation of the Modification of Diet in Renal Disease formula for estimating GFR with special emphasis on calibration of the serum creatinine assay. *Am J Kidney Dis* 2004;44:84-93.
- 1222 *Hallan S, Astor B, Lydersen S. Estimating glomerular filtration rate in the general population: the second Health Survey of Nord-Trondelag (HUNT II). *Nephrol Dial Transplant* 2006;21:1525-33.
- 5693 *Hansson LO, Wadstrom J, Lipcsey M, Biglarnia A, Larsson A. [Current Swedish renal function tests must be better. Uncertain diagnosis with routine formulas--great risk of wrong drug dosages]. *Lakartidningen* 2008;105:731-4.
- 1946 *Harambat J, Ranchin B, Dubourg L, Liutkus A, Hadj-Haissa A, Rivet C, et al. Renal function in pediatric liver transplantation: a long-term follow-up study. *Transplantation* 2008;86:1028-34.
- 1230 *Hari P, Bagga A, Mahajan P, Lakshmy R. Effect of malnutrition on serum creatinine and cystatin C levels. *Pediatr Nephrol* 2007;22:1757-61.
- 5697 *Harmoinen A, Lehtimaki T, Korpela M, Turjanmaa V, Saha H. Diagnostic accuracies of plasma creatinine, cystatin C, and glomerular filtration rate calculated by the Cockcroft-Gault and Levey (MDRD) formulas. *Clin Chem* 2003;49:1223-5.
- 5698 *Harmoinen AP, Kouri TT, Wirta OR, Lehtimaki TJ, Rantalaiho V, Turjanmaa VM, et al. Evaluation of plasma cystatin C as a marker for glomerular filtration rate in patients with type 2 diabetes. *Clin Nephrol* 1999;52:363-70.
- 5709 *Harzallah K, Sellam A, Mhiri A, Slim I, Halouas M, Mezzigue C, et al. Creatinine clearance estimation after kidney transplantation: an analysis of three methods. *Transplant Proc* 2007;39:2571-3.
- 1950 *Hayashi T, Nitta K, Hatano M, Nakauchi M, Nihei H. The serum cystatin C concentration measured by particle-enhanced immunonephelometry is well correlated with inulin clearance in patients with various types of glomerulonephritis. *Nephron* 1999;82:90-2.
- 5729 *Hazar V, Gungor O, Guven AG, Aydin F, Akbas H, Gungor F, et al. Renal function after hematopoietic stem cell transplantation in children. *Pediatr Blood Cancer* 2009;53:197-202.

- 5731 *Heaf JG, Iversen J. Uses and limitations of renal scintigraphy in renal transplantation monitoring. *Eur J Nucl Med* 2000;27:871-9.
- 13854 *Hebert PL, Nori US, Bhatt UY, Hebert LA. A modest proposal for improving the accuracy of creatinine-based GFR-estimating equations. *Nephrol Dial Transplant* 2011;26:2426-8.
- 13010 *Heilman RL, Mazur MJ. Cystatin C as a more sensitive indicator of diminished glomerular filtration rate. *Liver Transplantation* 2005;11:264-266.
- 1236 *Helin I, Axenram M, Grubb A. Serum cystatin C as a determinant of glomerular filtration rate in children. *Clin Nephrol* 1998;49:221-5.
- 13504 *Helou R. Should We Continue to Use the Cockcroft-Gault Formula? *Nephron - Clinical Practice* 2010;116:c172-c186.
- 5776 *Herget-Rosenthal S, Bokenkamp A, Hofmann W. How to estimate GFR-serum creatinine, serum cystatin C or equations? *Clin Biochem* 2007;40:153-61.
- 5780 *Herlitz H, Harris K, Risler T, Boner G, Bernheim J, Chanard J, et al. The effects of an ACE inhibitor and a calcium antagonist on the progression of renal disease: the Nephros Study. *Nephrol Dial Transplant* 2001;16:2158-65.
- 5792 *Herrera-Gutierrez ME, Seller-Perez G, Banderas-Bravo E, Munoz-Bono J, Lebron-Gallardo M, Fernandez-Ortega JF. Replacement of 24-h creatinine clearance by 2-h creatinine clearance in intensive care unit patients: a single-center study. *Intensive Care Med* 2007;33:1900-6.
- 5795 *Herts BR, Schneider E, Obuchowski N, Poggio E, Jain A, Baker ME. Probability of reduced renal function after contrast-enhanced CT: a model based on serum creatinine level, patient age, and estimated glomerular filtration rate. *AJR Am J Roentgenol* 2009;193:494-500.
- 5830 *Hjorth L, Wiebe T, Karpman D. Correct evaluation of renal glomerular filtration rate requires clearance assays. *Pediatr Nephrol* 2002;17:847-51.
- 13680 *Hjorth L, Wiebe T, Karpman D. Hyperfiltration evaluated by glomerular filtration rate at diagnosis in children with cancer. *Pediatr Blood Cancer*. 2011.
- 1246 *Hoek FJ, Kemperman FA, Krediet RT. A comparison between cystatin C, plasma creatinine and the Cockcroft and Gault formula for the estimation of glomerular filtration rate. *Nephrol Dial Transplant* 2003;18:2024-31.
- 5842 *Hoek FJ, Korevaar JC, Dekker FW, Boeschoten EW, Krediet RT. Estimation of residual glomerular filtration rate in dialysis patients from the plasma cystatin C level. *Nephrol Dial Transplant* 2007;22:1633-8.
- 5848 *Hogg RJ, Furth S, Lemley KV, Portman R, Schwartz GJ, Coresh J, et al. National Kidney Foundation's Kidney Disease Outcomes Quality Initiative clinical practice guidelines for chronic kidney disease in children and adolescents: evaluation, classification, and stratification. *Pediatrics* 2003;111:1416-21.
- 5850 *Hojs R, Bevc S, Antolinc B, Gorenjak M, Puklavec L. Serum cystatin C as an endogenous marker of renal function in the elderly. *Int J Clin Pharmacol Res* 2004;24:49-54.
- 5851 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Serum cystatin C as an endogenous marker of renal function in patients with mild to moderate impairment of kidney function. *Nephrol Dial Transplant* 2006;21:1855-62.
- 5852 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Serum cystatin C-based equation compared to serum creatinine-based equations for estimation of glomerular filtration rate in patients with chronic kidney disease. *Clin Nephrol* 2008;70:10-7.
- 5853 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Serum cystatin C as an endogenous marker of renal function in patients with chronic kidney disease. *Ren Fail* 2008;30:181-6.

13505 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Serum cystatin C-based formulas for prediction of glomerular filtration rate in patients with chronic kidney disease. *Nephron - Clinical Practice* 2010;114:c118-c126.

13875 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Kidney function estimating equations in patients with chronic kidney disease. *Int J Clin Pract* 2011;65:458-64.

5861 *Holmquist P, Torffvit O, Sjoblad S. Metabolic status in diabetes mellitus affects markers for glomerular filtration rate. *Pediatr Nephrol* 2003;18:536-40.

13685 *Horio M, Imai E, Yasuda Y, Watanabe T, Matsuo S. Modification of the CKD epidemiology collaboration (CKD-EPI) equation for Japanese: accuracy and use for population estimates. *Am J Kidney Dis* 2010;56:32-8.

13855 *Horio M, Imai E, Yasuda Y, Watanabe T, Matsuo S. Performance of serum cystatin C versus serum creatinine as a marker of glomerular filtration rate as measured by inulin renal clearance. *Clin Exp Nephrol* 2011.

5879 *Hornum M, Burton CM, Iversen M, Hovind P, Hilsted L, Feldt-Rasmussen B. Decline in 51Cr-labelled EDTA measured glomerular filtration rate following lung transplantation. *Nephrol Dial Transplant* 2007;22:3616-22.

13506 *Hottelart C, Achard JM, Fournier A. Creatininemia during fibrate therapy: A reply to Lipscombe et al. [3]. *Clinical nephrology* 2001;56:85-86.

13782 *Hsu CY, Bansal N. Measured GFR as "gold standard"--all that glitters is not gold? *Clin J Am Soc Nephrol* 2011;6:1813-4.

13856 *Hsu CY, Probert K, Xie D, Hamm L, He J, Miller E, et al. Measured GFR Does Not Outperform Estimated GFR in Predicting CKD-related Complications. *J Am Soc Nephrol* 2011;22:1931-1937.

13857 *Huang SH, Macnab JJ, Sontrop JM, Filler G, Gallo K, Lindsay RM, et al. Performance of the creatinine-based and the cystatin C-based glomerular filtration rate (GFR) estimating equations in a heterogenous sample of patients referred for nuclear GFR testing. *Transl Res* 2011;157:357-67.

13577 *Huang SH, Sharma AP, Yasin A, Lindsay RM, Clark WF, Filler G. Hyperfiltration Affects Accuracy of Creatinine eGFR Measurement. *Clin J Am Soc Nephrol*. 2011;6:274-280.

5938 *Ibrahim H, Mondress M, Tello A, Fan Y, Koopmeiners J, Thomas W. An alternative formula to the Cockcroft-Gault and the modification of diet in renal diseases formulas in predicting GFR in individuals with type 1 diabetes. *J Am Soc Nephrol* 2005;16:1051-60.

5939 *Ibrahim HN, Rogers T, Tello A, Matas A. The performance of three serum creatinine-based formulas in estimating GFR in former kidney donors. *Am J Transplant* 2006;6:1479-85.

61 *Ibrahim S, Rashid L, Darai M. Modification of Diet in Renal Disease equation underestimates glomerular filtration rate in Egyptian kidney donors. *Exp Clin Transplant* 2008;6:144-8.

13837 *Iliadis F, Didangelos T, Ntemka A, Makedou A, Moralidis E, Gotzamani-Psarakou A, et al. Glomerular filtration rate estimation in patients with type 2 diabetes: creatinine- or cystatin C-based equations? *Diabetologia* 2011.

2002 *Imai E, Horio M, Nitta K, Yamagata K, Iseki K, Hara S, et al. Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. *Clin Exp Nephrol* 2007;11:41-50.

1266 *Imai E, Horio M, Nitta K, Yamagata K, Iseki K, Tsukamoto Y, et al. Modification of the Modification of Diet in Renal Disease (MDRD) Study equation for Japan. *Am J Kidney Dis* 2007;50:927-37.

13858 *Inker LA, Eckfeldt J, Levey AS, Leiendecker-Foster C, Rynders G, Manzi J, et al. Expressing the CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration) Cystatin C Equations for Estimating GFR With Standardized Serum Cystatin C Values. *Am J Kidney Dis* 2011;58:682-4.

5965 *International Federation of Clinical C, Laboratory M, Working Group on Standardization of Glomerular Filtration Rate A, Panteghini M, Myers GL, Miller WG, et al. The importance of metrological traceability on the validity of creatinine measurement as an index of renal function. *Clin Chem Lab Med* 2006;44:1287-92.

5979 *Issa N, Meyer KH, Arrigain S, Choure G, Fatica RA, Nurko S, et al. Evaluation of creatinine-based estimates of glomerular filtration rate in a large cohort of living kidney donors. *Transplantation* 2008;86:223-30.

5983 *Itoh K. Comparison of methods for determination of glomerular filtration rate: Tc-99m-DTPA renography, predicted creatinine clearance method and plasma sample method. *Ann Nucl Med* 2003;17:561-5.

5984 *Itoh K, Tsushima S, Tsukamoto E, Tamaki N. Reappraisal of single-sample and gamma camera methods for determination of the glomerular filtration rate with 99mTc-DTPA. *Ann Nucl Med* 2000;14:143-50.

13681 *Ix JH, Wassel CL, Stevens LA, Beck GJ, Froissart M, Navis G, et al. Equations to estimate creatinine excretion rate: the CKD epidemiology collaboration. *Clin J Am Soc Nephrol*. 2011;6:184-91.

6005 *Jacobi D, Lavigne C, Halimi JM, Fierrard H, Andres C, Couet C, et al. Variability in creatinine excretion in adult diabetic, overweight men and women: consequences on creatinine-based classification of renal disease. *Diabetes Res Clin Pract* 2008;80:102-7.

6008 *Jacobsen P, Tarnow L, Carstensen B, Hovind P, Poirier O, Parving HH. Genetic variation in the Renin-Angiotensin system and progression of diabetic nephropathy. *J Am Soc Nephrol* 2003;14:2843-50.

13652 *Janssen A, Dumoulin A, De Waele JJ, Vanholder R, Waterloos MA, Hoste EA. Assessment of increased glomerular filtration rate in ICU patients. *SOURCE Intensive Care Medicine* (2010) 36 SUPPL. 2 (S402). Date of Publication: September 2010 CONFERENCE NAME 23rd Annual Congress of the European Society of Intensive Care Medicine, ESICM CONFERENCE LOCATION Barcelona, Spain CONFERENCE DATE 2010-10-09 to 2010-10-13.

14046 *Johnson S, Rishi R, Andone A, Khawandi W, Al-Said J, Gletsu-Miller N, et al. Determinants and functional significance of renal parenchymal volume in adults. *Clinical Journal of the American Society of Nephrology* 2011;6:70-76.

6057 *Johnson TM, 2nd, Sands JM, Ouslander JG. A prospective evaluation of the glomerular filtration rate in older adults with frequent nighttime urination. *J Urol* 2002;167:146-50.

13682 *Jones GR. Use of the CKD-EPI equation for estimation of GFR in an Australian cohort. *Pathology*. 2010;42:487-8.

6062 *Jones GR, Imam SK. Validation of the revised MDRD formula and the original Cockcroft and Gault formula for estimation of the glomerular filtration rate using Australian data. *Pathology* 2009;41:379-82.

13838 *Jones MA, Golightly LK, Stolpman NM. Use of recalibrated serum creatinine concentrations for adjustment of drug dosages: determination of values compatible with conventional dosing recommendations. *Ann Pharmacother* 2011;45:748-56.

6065 *Jonsson AS, Flodin M, Hansson LO, Larsson A. Estimated glomerular filtration rate (eGFR_{CystC}) from serum cystatin C shows strong agreement with iohexol clearance in patients with low GFR. *Scand J Clin Lab Invest* 2007;67:801-9.

6067 *Jonsson O, Herlitz H, Lindholm E, Lindstedt G, Pazooki D, Tornqvist H, et al. Prophylaxis against bone loss in Kock reservoir patients with reduced glomerular filtration rate. *Scand J Urol Nephrol* 2005;39:200-5.

3507 *Kalaitzakis E, Bjornsson E. Renal function and cognitive impairment in patients with liver cirrhosis. *Scand J Gastroenterol* 2007;42:1238-44.

1294 *Kallner A, Ayling PA, Khatami Z. Does eGFR improve the diagnostic capability of S-Creatinine concentration results? A retrospective population based study. *Int J Med Sci* 2008;5:9-17.

6103 *Kallner A, Khatami Z. How does the MDRD Study equation compare with serum creatinine in routine healthcare? Anatomy of MDRD-eGFR. *Scand J Clin Lab Invest Suppl* 2008;241:39-45.

- 2015 *Kamar N, Van TT, Ribes D, Modesto A, Cointault O, Lavayssiere L, et al. Renal function and histology in kidney transplant patients receiving tacrolimus and sirolimus or mycophenolate mofetil. *Exp Clin Transplant* 2006;4:532-6.
- 6109 *Kamata K, Tochikubo O. Estimation of 24-h urinary sodium excretion using lean body mass and overnight urine collected by a pipe-sampling method. *J Hypertens* 2002;20:2191-7.
- 609 *Kang YS, Han KH, Han SY, Kim HK, Cha DR. Characteristics of population with normal serum creatinine impaired renal function and: the validation of a MDRD formula in a healthy general population. *Clin Nephrol* 2005;63:258-66.
- 6120 *Kapoor A, Kumar S, Gulati S, Gambhir S, Sethi RS, Sinha N. The role of theophylline in contrast-induced nephropathy: a case-control study. *Nephrol Dial Transplant* 2002;17:1936-41.
- 6124 *Karanikas G, Schutz M, Szabo M, Becherer A, Wiesner K, Dudczak R, et al. Isotopic renal function studies in severe hypothyroidism and after thyroid hormone replacement therapy. *Am J Nephrol* 2004;24:41-5.
- 6125 *Karawajczyk M, Ramklint M, Larsson A. Reduced cystatin C-estimated GFR and increased creatinine-estimated GFR in comparison with iohexol-estimated GFR in a hyperthyroid patient: A case report. *J Med Case Reports* 2008;2:66.
- 6132 *Karstila K, Harmoinen AP, Lehtimaki TJ, Korpela MM, Mustonen JT, Saha HH. Measurement of the kidney function in patients with rheumatoid arthritis: plasma cystatin C versus ⁵¹Cr-EDTA clearance. *Nephron Clin Pract* 2008;108:c284-90.
- 6156 *Kemperman FA, Krediet RT, Arisz L. Formula-derived prediction of the glomerular filtration rate from plasma creatinine concentration. *Nephron* 2002;91:547-58.
- 6157 *Kemperman FA, Silberbusch J, Slaats EH, Prins AM, Krediet RT, Arisz L. Follow-up of GFR estimated from plasma creatinine after cimetidine administration in patients with diabetes mellitus type 2. *Clin Nephrol* 2000;54:255-60.
- 6160 *Kemperman FA, Surachno J, Krediet RT, Arisz L. Cimetidine improves prediction of the glomerular filtration rate by the Cockcroft-Gault formula in renal transplant recipients. *Transplantation* 2002;73:770-4.
- 1306 *Kielstein JT, Boger RH, Bode-Boger SM, Frolich JC, Haller H, Ritz E, et al. Marked increase of asymmetric dimethylarginine in patients with incipient primary chronic renal disease. *J Am Soc Nephrol* 2002;13:170-6.
- 6181 *Kielstein JT, Martens-Lobenhoffer J, Vollmer S, Bode-Boger SM. L-Arginine, ADMA, SDMA, creatinine, MDRD formula: detour to renal function testing. *J Nephrol* 2008;21:959-61.
- 2026 *Kielstein JT, Salpeter SR, Bode-Boeger SM, Cooke JP, Fliser D. Symmetric dimethylarginine (SDMA) as endogenous marker of renal function--a meta-analysis. *Nephrol Dial Transplant* 2006;21:2446-51.
- 6185 *Kilpatrick ES, Keevil BG, Addison GM. Does adjustment of GFR to extracellular fluid volume improve the clinical utility of cystatin C? *Arch Dis Child* 2000;82:499-502.
- 6206 *Kingdon EJ, Knight CJ, Dustan K, Irwin AG, Thomas M, Powis SH, et al. Calculated glomerular filtration rate is a useful screening tool to identify scleroderma patients with renal impairment. *Rheumatology (Oxford)* 2003;42:26-33.
- 63 *Kleinert J, Lorenz M, Hauser AC, Becherer A, Staudenherz A, Fodinger M, et al. Measurement of renal function in patients with Fabry disease. *Acta Paediatr Suppl* 2005;94:19-23; discussion 9-10.
- 6229 *Kletzel M, Pirich L, Haut P, Cohn RA. Comparison of Tc-99 measurement of glomerular filtration rate vs. calculated creatinine clearance to assess renal function pretransplant in pediatric patients undergoing hematopoietic stem cell transplantation. *Pediatr Transplant* 2005;9:584-8.
- 2035 *Koch M, Beien A, Fusha Ller A, Zitta S, Haastert B, Trapp R. Impact of age, body mass index, insulin resistance and proteinuria on the kidney function in obese patients with Type 2 diabetes and renal insufficiency. *Clin Nephrol* 2008;69:10-7.

- 13653 *Koetje PM, Spaan JJ, Kooman JP, Spaanderman ME, Peeters LL. Pregnancy Reduces the Accuracy of the Estimated Glomerular Filtration Rate Based on Cockcroft-Gault and MDRD Formulas. *Reprod Sci*. 2011;18:456-462.
- 6247 *Komenda P, Beaulieu M, Seccombe D, Levin A. Regional implementation of creatinine measurement standardization. *J Am Soc Nephrol* 2008;19:164-9.
- 6256 *Kopple JD, Greene T, Chumlea WC, Hollinger D, Maroni BJ, Merrill D, et al. Relationship between nutritional status and the glomerular filtration rate: results from the MDRD study. *Kidney Int* 2000;57:1688-703.
- 628 *Kottgen A, Selvin E, Stevens LA, Levey AS, Van Lente F, Coresh J. Serum cystatin C in the United States: the Third National Health and Nutrition Examination Survey (NHANES III). *Am J Kidney Dis* 2008;51:385-94.
- 6275 *Kramer H, Palmas W, Kestenbaum B, Cushman M, Allison M, Astor B, et al. Chronic kidney disease prevalence estimates among racial/ethnic groups: the Multi-Ethnic Study of Atherosclerosis. *Clin J Am Soc Nephrol* 2008;3:1391-7.
- 6285 *Krieser D, Rosenberg AR, Kainer G, Naidoo D. The relationship between serum creatinine, serum cystatin C and glomerular filtration rate in pediatric renal transplant recipients: a pilot study. *Pediatr Transplant* 2002;6:392-5.
- 6290 *Kronenberg F, Kuen E, Ritz E, Konig P, Kraatz G, Lhotta K, et al. Apolipoprotein A-IV serum concentrations are elevated in patients with mild and moderate renal failure. *J Am Soc Nephrol* 2002;13:461-9.
- 2041 *Kuan Y, Hossain M, Surman J, El Nahas AM, Haylor J. GFR prediction using the MDRD and Cockcroft and Gault equations in patients with end-stage renal disease. *Nephrol Dial Transplant* 2005;20:2394-401.
- 13557 *Kukla A, El-Shahawi Y, Leister E, Kasiske B, Mauer M, Matas A, et al. GFR-estimating models in kidney transplant recipients on a steroid-free regimen. In: *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*; 2010. p 1653-61.
- 6314 *Kuriyan R, Kurpad AV. Prediction of total body muscle mass from simple anthropometric measurements in young Indian males. *Indian J Med Res* 2004;119:121-8.
- 6315 *Kuriyan R, Thomas T, Kurpad AV. Total body muscle mass estimation from bioelectrical impedance analysis & simple anthropometric measurements in Indian men. *Indian J Med Res* 2008;127:441-6.
- 6321 *Kuypers DR, Neumayer HH, Fritsche L, Budde K, Rodicio JL, Vanrenterghem Y, et al. Calcium channel blockade and preservation of renal graft function in cyclosporine-treated recipients: a prospective randomized placebo-controlled 2-year study. *Transplantation* 2004;78:1204-11.
- 13507 *Kwong YT, Stevens LA, Selvin E, Zhang YL, Greene T, Van Lente F, et al. Imprecision of urinary iothalamate clearance as a gold-standard measure of GFR decreases the diagnostic accuracy of kidney function estimating equations. In: *American journal of kidney diseases : the official journal of the National Kidney Foundation*; 2010. p 39-49.
- 1330 *Kyhse-Andersen J, Schmidt C, Nordin G, Andersson B, Nilsson-Ehle P, Lindstrom V, et al. Serum cystatin C, determined by a rapid, automated particle-enhanced turbidimetric method, is a better marker than serum creatinine for glomerular filtration rate. *Clin Chem* 1994;40:1921-6.
- 6345 *Lamb EJ, Tomson CR, Roderick PJ, Clinical Sciences Reviews Committee of the Association for Clinical B. Estimating kidney function in adults using formulae. *Ann Clin Biochem* 2005;42:321-45.
- 6347 *Lamb EJ, Webb MC, Simpson DE, Coakley AJ, Newman DJ, O'Riordan SE. Estimation of glomerular filtration rate in older patients with chronic renal insufficiency: is the modification of diet in renal disease formula an improvement? *J Am Geriatr Soc* 2003;51:1012-7.
- 6348 *Lamb EJ, Wood J, Stowe HJ, O'Riordan SE, Webb MC, Dalton RN. Susceptibility of glomerular filtration rate estimations to variations in creatinine methodology: a study in older patients. *Ann Clin Biochem* 2005;42:11-8.

- 13508 *Lane BR, Demirjian S, Weight CJ, Larson BT, Poggio ED, Campbell SC. Performance of the Chronic Kidney Disease-Epidemiology Study Equations for Estimating Glomerular Filtration Rate Before and After Nephrectomy. *Journal of Urology* 2010;183:896-902.
- 6362 *Lankisch P, Wessalowski R, Maisonneuve P, Haghgu M, Hermsen D, Kramm CM. Serum cystatin C is a suitable marker for routine monitoring of renal function in pediatric cancer patients, especially of very young age. *Pediatr Blood Cancer* 2006;46:767-72.
- 6363 *Laroche ML, Charmes JP, Marcheix A, Bouthier F, Merle L. Estimation of glomerular filtration rate in the elderly: Cockcroft-Gault formula versus modification of diet in renal disease formula. *Pharmacotherapy* 2006;26:1041-6.
- 6365 *Larsson A. Cystatin C: an emerging glomerular filtration rate marker. *Scand J Clin Lab Invest* 2005;65:89-91.
- 5 *Larsson A, Flodin M, Hansson LO, Carlsson L. Patient selection has a strong impact on cystatin C and Modification of Diet in Renal Disease (MDRD) estimated glomerular filtration rate. *Clin Biochem* 2008;41:1355-61.
- 6368 *Larsson A, Malm J, Grubb A, Hansson LO. Calculation of glomerular filtration rate expressed in mL/min from plasma cystatin C values in mg/L. *Scand J Clin Lab Invest* 2004;64:25-30.
- 14047 *Laskin BL, Goebel J, Bissler JJ, Khoury JC, Davies SM, Jodele S. Cystatin C is a more sensitive estimate of kidney function than creatinine in pediatric autologous bone marrow transplantation. *Biology of Blood and Marrow Transplantation* 2011;17:S261.
- 6388 *Le Bricon T, Leblanc I, Benlakehal M, Gay-Bellile C, Erlich D, Boudaoud S. Evaluation of renal function in intensive care: plasma cystatin C vs. creatinine and derived glomerular filtration rate estimates. *Clin Chem Lab Med* 2005;43:953-7.
- 6389 *Le Bricon T, Thervet E, Benlakehal M, Bousquet B, Legendre C, Erlich D. Changes in plasma cystatin C after renal transplantation and acute rejection in adults. *Clin Chem* 1999;45:2243-9.
- 6391 *le Riche M, Zemlin AE, Erasmus RT, Davids MR. An audit of 24-hour creatinine clearance measurements at Tygerberg Hospital and comparison with prediction equations. *S Afr Med J* 2007;97:968-70.
- 6403 *Lee BM, Yoon SN, Oh CK, Kim JH, Kim SJ, Kim H, et al. Fractional creatinine clearance of the donated kidney using Cockcroft-Gault formula as a predictor of graft function after living donor transplantation. *Transplant Proc* 2006;38:1974-6.
- 13654 *Lee CS, Cha RH, Lim YH, Kim H, Song KH, Gu N, et al. Ethnic coefficients for glomerular filtration rate estimation by the Modification of Diet in Renal Disease study equations in the Korean population. *J Korean Med Sci*. 2010;25:1616-25.
- 6404 *Lee D, Levin A, Roger SD, McMahon LP. Longitudinal analysis of performance of estimated glomerular filtration rate as renal function declines in chronic kidney disease. *Nephrol Dial Transplant* 2009;24:109-16.
- 6407 *Lee EK, Cho YM, Kim JT, Koo BK, Cho HY, Ku YH, et al. Derivation of a new equation for estimating creatinine clearance by using fat-free mass and serum creatinine concentration in Korean patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract* 2009;83:44-9.
- 13509 *Lee JC, Kang IM, Chou CY, Tseng YH, Huang CC, Shih CM, et al. Difference between estimated glomerulofiltration rate by modification of diet in renal diseases and Cockcroft-Gault formula in general population. *Journal of Internal Medicine of Taiwan* 2009;20:148-154.
- 1346 *Leger F, Bouissou F, Coulais Y, Tafani M, Chatelut E. Estimation of glomerular filtration rate in children. *Pediatr Nephrol* 2002;17:903-7.
- 271 *Lemann J, Bidani AK, Bain RP, Lewis EJ, Rohde RD. Use of the serum creatinine to estimate glomerular filtration rate in health and early diabetic nephropathy. Collaborative Study Group of Angiotensin Converting Enzyme Inhibition in Diabetic Nephropathy. *Am J Kidney Dis* 1990;16:236-43.

- 13510 *Lemley KV. Pediatric nephrology: Estimating GFR in children: Schwartz redux. *Nature Reviews Nephrology* 2009;5:310-311.
- 6439 *Leung YY, Lo KM, Tam LS, Szeto CC, Li EK, Kun EW. Estimation of glomerular filtration rate in patients with systemic lupus erythematosus. *Lupus* 2006;15:276-81.
- 7 *Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med* 1999;130:461-70.
- 13511 *Levey AS, Coresh J, Balk E, Kausz A, Levin A, Steffes MW, et al. Clinical guidelines National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification [corrected] [published erratum appears in ANN INTERN MED 2003 Oct 7;139(7):605]. *Annals of Internal Medicine* 2003;139:137-47.
- 13512 *Levey AS, Coresh J, Greene T, Stevens LA, Zhang Y, Hendriksen S. Chronic Kidney Disease Epidemiology Collaboration. Using standardized serum creatinine values in the Modification of Diet in Renal Disease study equation for estimating glomerular filtration rate (*Annals of Internal Medicine* (2006) 145, (247-254)). *Annals of Internal Medicine* 2008;149:519.
- 6444 *Levey AS, Coresh J, Greene T, Stevens LA, Zhang YL, Hendriksen S, et al. Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate. *Ann Intern Med* 2006;145:247-54.
- 13513 *Levey AS, Stevens LA. Estimating GFR Using the CKD Epidemiology Collaboration (CKD-EPI) Creatinine Equation: More Accurate GFR Estimates, Lower CKD Prevalence Estimates, and Better Risk Predictions. *American Journal of Kidney Diseases* 2010;55:622-627.
- 6446 *Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF, 3rd, Feldman HI, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;150:604-12.
- 6450 *Lewis J, Agodoa L, Cheek D, Greene T, Middleton J, O'Connor D, et al. Comparison of cross-sectional renal function measurements in African Americans with hypertensive nephrosclerosis and of primary formulas to estimate glomerular filtration rate. *Am J Kidney Dis* 2001;38:744-53.
- 3601 *Li H, Zhang X, Xu G, Wang X, Zhang C. Determination of reference intervals for creatinine and evaluation of creatinine-based estimating equation for Chinese patients with chronic kidney disease. *Clin Chim Acta* 2009;403:87-91.
- 13394 *Li HX, Xu GB, Wang XJ, Zhang XC, Yang JM. Diagnostic accuracy of various glomerular filtration rates estimating equations in patients with chronic kidney disease and diabetes. *Chinese Medical Journal* 2010;123:745-751.
- 6463 *Li L, Greene T. Varying coefficients model with measurement error. *Biometrics* 2008;64:519-26.
- 66 *Lim WH, Lim EM, McDonald S. Lean body mass-adjusted Cockcroft and Gault formula improves the estimation of glomerular filtration rate in subjects with normal-range serum creatinine. *Nephrology (Carlton)* 2006;11:250-6.
- 6500 *Lindegaard MW, Aass N, Bue ES, Theodorsen L, Fossa SD. Glomerular filtration rate, 131I-hippuran clearance and estimated creatinine clearance in cancer patients. *Br J Cancer* 1991;64:401-5.
- 6503 *Ling Q, Xu X, Li J, Wu J, Chen J, Xie H, et al. A new serum cystatin C-based equation for assessing glomerular filtration rate in liver transplantation. *Clin Chem Lab Med* 2008;46:405-10.
- 6504 *Ling Q, Xu X, Li JJ, Chen J, Shen JW, Zheng SS. Alternative definition of acute kidney injury following liver transplantation: based on serum creatinine and cystatin C levels. *Transplant Proc* 2007;39:3257-60.
- 6511 *Linzbach S, Samigullin A, Yilmaz S, Tsioga M, Zeiher AM, Spyridopoulos I. Role of N-terminal pro-brain natriuretic peptide and cystatin C to estimate renal function in patients with and without heart failure. *Am J Cardiol* 2009;103:1128-33.

- 13657 *Lipsey M, Furebring M, Rubertsson S, Larsson A. Significant differences when using creatinine, modification of diet in renal disease, or cystatin C for estimating glomerular filtration rate in ICU patients. *Ups J Med Sci.* 2010.
- 6514 *Lippi G, Banfi G, Luca Salvagno G, Montagnana M, Franchini M, Cesare Guidi G. Comparison of creatinine-based estimations of glomerular filtration rate in endurance athletes at rest. *Clin Chem Lab Med* 2008;46:235-9.
- 6516 *Lippi G, De Franceschi L, Salvagno GL, Montagnana M, Guidi GC. Estimation of glomerular filtration rate by the modification of diet in renal disease (MDRD) equation in patients with sickle cell disease. *Clin Chem Lab Med* 2008;46:1200-1.
- 13658 *Lippi G, Franchini M, Targher G. Detection of chronic kidney disease in hospitalized patients: Is one estimating glomerular filtration rate equation better than another? *European Journal of Internal Medicine* 2011;22:119-120.
- 6518 *Lippi G, Targher G, Salvagno GL, Montagnana M, Guidi GC. Detecting of chronic kidney disease in older people by the MDRD and MCQ formulas. *Age Ageing* 2008;37:722.
- 13659 *Liu X, Lv L, Wang C, Shi CG, Cheng CL, Tang H, et al. Comparison of prediction equations to estimate glomerular filtration rate in Chinese patients with chronic kidney disease. *Intern Med J.* 2010.
- 13660 *Liu Y, Xu GB. Trueness investigation of routine creatinine assays on nine homogeneous systems in Beijing demonstrates an encouraging outcome that meets clinical requirements. *Chin Med J (Engl).* 2010;123:2364-9.
- 6548 *Long SE, Thompson WL, Sonnemaker RE, Pounds BK, Burdine JA. Tc-99m glucoheptonate estimation of glomerular filtration rate. Correlation with endogenous creatinine clearance. *Clin Nucl Med* 1984;9:271-4.
- 6556 *Louvar DW, Rogers TB, Bailey RF, Matas AJ, Ibrahim HN. Cystatin C is not superior to creatinine-based models in estimating glomerular filtration rate in former kidney donors. *Transplantation* 2007;84:1112-7.
- 14048 *Luangchosiri C, Sobhonslidsuk A, Klyprayong P, Meemaew P, Sangcakul A, Chittamma A, et al. Estimating GFR with creatinine and cystatin C-based equations in cirrhotic patients. *Hepatology International* 2011;5:357.
- 2083 *Luke DR, Halstenson CE, Opsahl JA, Matzke GR. Validity of creatinine clearance estimates in the assessment of renal function. *Clin Pharmacol Ther* 1990;48:503-8.
- 6584 *Ma YC, Zuo L, Chen JH, Luo Q, Yu XQ, Li Y, et al. Modified glomerular filtration rate estimating equation for Chinese patients with chronic kidney disease. *J Am Soc Nephrol* 2006;17:2937-44.
- 1369 *Ma YC, Zuo L, Chen JH, Luo Q, Yu XQ, Li Y, et al. Improved GFR estimation by combined creatinine and cystatin C measurements. *Kidney Int* 2007;72:1535-42.
- 6586 *Ma YC, Zuo L, Wang M, Zhou YH, Zhang CL, Xu GB, et al. [The application of glomerular filtration rate estimation equations in different stages of chronic kidney disease]. *Zhonghua Nei Ke Za Zhi* 2005;44:285-9.
- 13839 *Maahs DM, Jalal D, McFann K, Rewers M, Snell-Bergeon JK. Systematic shifts in cystatin C between 2006 and 2010. *Clin J Am Soc Nephrol* 2011;6:1952-5.
- 6592 *MacAulay J, Thompson K, Kiberd BA, Barnes DC, Peltekian KM. Serum creatinine in patients with advanced liver disease is of limited value for identification of moderate renal dysfunction: are the equations for estimating renal function better? *Can J Gastroenterol* 2006;20:521-6.
- 6593 *MacDiarmid SA, McIntyre WJ, Anthony A, Bailey RR, Turner JG, Arnold EP. Monitoring of renal function in patients with spinal cord injury. *BJU Int* 2000;85:1014-8.
- 6594 *Macdonald J, Marcora S, Jibani M, Roberts G, Kumwenda M, Glover R, et al. GFR estimation using cystatin C is not independent of body composition. *Am J Kidney Dis* 2006;48:712-9.

- 13395 *Macdonald JH, Marcora SM, Jibani M, Roberts G, Kumwenda MJ, Glover R, et al. Bioelectrical impedance can be used to predict muscle mass and hence improve estimation of glomerular filtration rate in non-diabetic patients with chronic kidney disease. *Nephrology Dialysis Transplantation* 2006;21:3481-3487.
- 13396 *Macdonald JH, Marcora SM, Kumwenda MJ, Jibani M, Roberts G, Glover R, et al. The relationship between estimated glomerular filtration rate, demographic and anthropometric variables is mediated by muscle mass in non-diabetic patients with chronic kidney disease. *Nephrology Dialysis Transplantation* 2006;21:3488-3494.
- 6598 *Macisaac RJ, Tsalamandris C, Thomas MC, Premaratne E, Panagiotopoulos S, Smith TJ, et al. Estimating glomerular filtration rate in diabetes: a comparison of cystatin-C- and creatinine-based methods. *Diabetologia* 2006;49:1686-9.
- 1371 *Macisaac RJ, Tsalamandris C, Thomas MC, Premaratne E, Panagiotopoulos S, Smith TJ, et al. The accuracy of cystatin C and commonly used creatinine-based methods for detecting moderate and mild chronic kidney disease in diabetes. *Diabet Med* 2007;24:443-8.
- 6600 *Mackelaite L, Alsauskas ZC, Ranganna K. Renal failure in patients with cirrhosis. *Med Clin North Am* 2009;93:855-69, viii.
- 13859 *Macunluoglu B, Gokce I, Atakan A, Demirci M, Ari E, Topuzoglu A, et al. A comparison of different methods for the determination of glomerular filtration rate in elderly patients with chronic renal failure. *Int Urol Nephrol* 2011;43:257-63.
- 13661 *Madder RD, Hickman L, Crimmins GM, Puri M, Marinescu V, Safian RD. Estimated glomerular filtration rates are not useful for serial assessment of renal function in patients with atherosclerotic renal artery stenosis. *Journal of the American College of Cardiology* 2010;56:B93.
- 1373 *Mafham MM, Niculescu-Duvaz I, Barron J, Emberson JR, Dockrell ME, Landray MJ, et al. A practical method of measuring glomerular filtration rate by iohexol clearance using dried capillary blood spots. *Nephron Clin Pract* 2007;106:c104-12.
- 1374 *Mahajan S, Mukhiya GK, Singh R, Tiwari SC, Kalra V, Bhowmik DM, et al. Assessing glomerular filtration rate in healthy Indian adults: a comparison of various prediction equations. *J Nephrol* 2005;18:257-61.
- 6615 *Mahajan S, Mukhiya GK, Singh R, Tiwari SC, Kalra V, Guleria S, et al. Assessing suitability for renal donation: can equations predicting glomerular filtration rate substitute for a reference method in the Indian population? *Nephron Clin Pract* 2005;101:c128-33.
- 2087 *Maillard N, Mariat C, Bonneau C, Mehdi M, Thibaudin L, Laporte S, et al. Cystatin C-based equations in renal transplantation: moving toward a better glomerular filtration rate prediction? *Transplantation* 2008;85:1855-8.
- 13397 *Maillard N, Mehdi M, Thibaudin L, Berthoux F, Alamartine E, Mariat C. Creatinine-based GFR predicting equations in renal transplantation: Reassessing the tubular secretion effect. *Nephrology Dialysis Transplantation* 2010;25:3076-3082.
- 14049 *Maillard N, Moranne O, Thibaudin L, Laurent B, Alamartine E, Mariat C. One-year GFR as a surrogate for long term allograft survival: Estimated versus measured GFR. *American Journal of Transplantation* 2011;11:400.
- 279 *Mak RH, Al Dahhan J, Azzopardi D, Bosque M, Chantler C, Haycock GB. Measurement of glomerular filtration rate in children after renal transplantation. *Kidney Int* 1983;23:410-3.
- 6632 *Malmrose LC, Gray SL, Pieper CF, Blazer DG, Rowe JW, Seeman TE, et al. Measured versus estimated creatinine clearance in a high-functioning elderly sample: MacArthur Foundation Study of Successful Aging. *J Am Geriatr Soc* 1993;41:715-21.
- 3636 *Marco MP, Muray S, Valdivielso JM, Fernandez E. Occult chronic kidney disease: discordance among different methods used to estimate glomerular filtration rate in a healthy population. *Clin Nephrol* 2009;71:475-81.

2095 *Mariat C, Alamartine E, Barthelemy JC, De Filippis JP, Thibaudin D, Berthoux P, et al. Assessing renal graft function in clinical trials: can tests predicting glomerular filtration rate substitute for a reference method? *Kidney Int* 2004;65:289-97.

6666 *Marouf R, Mojiminiyi O, Abdella N, Kortom M, Al Wazzan H. Comparison of renal function markers in Kuwaiti patients with sickle cell disease. *J Clin Pathol* 2006;59:345-51.

6671 *Marshall MR, Song Q, Ma TM, MacDonell SG, Kasabov NK. Evolving connectionist system versus algebraic formulas for prediction of renal function from serum creatinine. *Kidney Int* 2005;67:1944-54.

13860 *Martin JH, Fay MF, Udy A, Roberts J, Kirkpatrick C, Ungerer J, et al. Pitfalls of using estimations of glomerular filtration rate in an intensive care population. *Intern Med J* 2011;41:537-43.

6677 *Martin L, Chatelut E, Boneu A, Rostaing L, Roussilhes C, Caselles O, et al. Improvement of the Cockcroft-Gault equation for predicting glomerular filtration in cancer patients. *Bull Cancer* 1998;85:631-6.

14050 *Martinez-Martinez MU, Mandeville PB, Llamazares-Azuara L, Abud-Mendoza C. Assessment of renal function in systemic lupus erythematosus (SLE) patients. *Lupus* 2011;20:394.

2099 *Martini S, Prevot A, Mosig D, Werner D, van Melle G, Guignard JP. Glomerular filtration rate: measure creatinine and height rather than cystatin C! *Acta Paediatr* 2003;92:1052-7.

6683 *Marx GM, Blake GM, Galani E, Steer CB, Harper SE, Adamson KL, et al. Evaluation of the Cockcroft-Gault, Jelliffe and Wright formulae in estimating renal function in elderly cancer patients. *Ann Oncol* 2004;15:291-5.

13735 *Mathisen UD, Melsom T, Ingebretsen OC, Jenssen T, Njolstad I, Solbu MD, et al. Estimated GFR associates with cardiovascular risk factors independently of measured GFR. *J Am Soc Nephrol* 2011;22:927-37.

2105 *Matsuo S, Imai E, Horio M, Yasuda Y, Tomita K, Nitta K, et al. Revised equations for estimated GFR from serum creatinine in Japan. *Am J Kidney Dis* 2009;53:982-92.

13398 *Matsuo S, Yasuda Y, Imai E, Horio M. Current status of estimated glomerular filtration rate (eGFR) equations for Asians and an approach to create a common eGFR equation. *Nephrology* 2010;15:45-48.

6700 *Mattman A, Eintracht S, Mock T, Schick G, Seccombe DW, Hurley RM, et al. Estimating pediatric glomerular filtration rates in the era of chronic kidney disease staging. *J Am Soc Nephrol* 2006;17:487-96.

6701 *Matz R. Prediction equation for glomerular filtration rate. *Ann Intern Med* 1999;131:629; author reply 630.

6706 *Mauss S, Berger F, Kuschak D, Henke J, Hegener P, Wolf E, et al. Cystatin C as a marker of renal function is affected by HIV replication leading to an underestimation of kidney function in HIV patients. *Antivir Ther* 2008;13:1091-5.

6710 *Mazur MJ, Rea DJ, Griffin MD, Larson TS, Prieto M, Gloor JM, et al. Decline in native renal function early after bladder-drained pancreas transplantation alone. *Transplantation* 2004;77:844-9.

2110 *McDougal WS, Koch MO. Accurate determination of renal function in patients with intestinal urinary diversions. *J Urol* 1986;135:1175-8.

138 *McElduff A, Shuter B, Cooper R, Davies L, Fulcher G, Hoschl R, et al. Measuring renal function in patients with diabetes mellitus. *J Diabetes Complications* 1997;11:225-9.

6722 *McKillop DJ, Cairns B, Duly E, Van Drimmelen M, Ryan M. The effect of serum creatinine method choice on estimated glomerular filtration rate determined by the abbreviated MDRD formula. *Ann Clin Biochem* 2006;43:220-2.

6723 *McLigeyo SO. Calculation of creatinine clearance from plasma creatinine. *East Afr Med J* 1993;70:3-5.

3655 *McNamara NV, Chen R, Janu MR, Bwititi P, Car G, Seibel M. Early renal failure detection by cystatin C in Type 2 diabetes mellitus: varying patterns of renal analyte expression. *Pathology* 2009;41:269-75.

- 6745 *Mention K, Lahoche-Manucci A, Bonneville M, Pruvot FR, Declerck N, Foulard M, et al. Renal function outcome in pediatric liver transplant recipients. *Pediatr Transplant* 2005;9:201-7.
- 13662 *Michels WM, Grootendorst DC, Verduijn M, Elliott EG, Dekker FW, Krediet RT. Performance of the Cockcroft-Gault, MDRD, and new CKD-EPI formulas in relation to GFR, age, and body size. *Clin J Am Soc Nephrol*. 2010;5:1003-9.
- 13134 *Miller WG. Estimating equations for glomerular filtration rate in children: Laboratory considerations. *Clinical Chemistry* 2009;55:402-403.
- 13399 *Mingat N, Allard J, Villard E, Guillotreau J, Paparel P, Roche JB, et al. Which is the most accurate biomarker to assess renal functions in neurogenic patients? *European Urology, Supplements* 2010;9:180-181.
- 6806 *Mohler JL, Ellison MF, Flanigan RC. Creatinine clearance prediction in spinal cord injury patients: comparison of 6 prediction equations. *J Urol* 1988;139:706-9.
- 1402 *Mojiminiyi OA, Abdella N. Evaluation of cystatin C and beta-2 microglobulin as markers of renal function in patients with type 2 diabetes mellitus. *J Diabetes Complications* 2003;17:160-8.
- 6821 *Montgomery MJ, Beringer PM, Louie SG, Gill MA. Estimation of creatinine clearance in patients with metastatic ovarian cancer. *Ther Drug Monit* 2000;22:695-700.
- 6823 *Montini G, Cosmo L, Amici G, Mussap M, Zacchello G. Plasma cystatin C values and inulin clearances in premature neonates. *Pediatr Nephrol* 2001;16:463-5.
- 6836 *Morgan C, Senthilselvan A, Bamforth F, Hoskinson M, Gowrishankar M. Correlation between cystatin C- and renal scan-determined glomerular filtration rate in children with spina bifida. *Pediatr Nephrol* 2008;23:329-32.
- 1407 *Morris MC, Allanby CW, Toseland P, Haycock GB, Chantler C. Evaluation of a height/plasma creatinine formula in the measurement of glomerular filtration rate. *Arch Dis Child* 1982;57:611-5.
- 2131 *Mpio I, Laville M, Hadj-Aissa A, Fauvel JP. Predicted creatinine clearance to evaluate glomerular filtration rate in black Caribbean subjects. *Nephrol Dial Transplant* 2003;18:1307-10.
- 13840 *Murata K, Baumann NA, Saenger AK, Larson TS, Rule AD, Lieske JC. Relative performance of the MDRD and CKD-EPI equations for estimating glomerular filtration rate among patients with varied clinical presentations. *Clin J Am Soc Nephrol* 2011;6:1963-72.
- 6890 *Murthy K, Stevens LA, Stark PC, Levey AS. Variation in the serum creatinine assay calibration: a practical application to glomerular filtration rate estimation. *Kidney Int* 2005;68:1884-7.
- 144 *Mussap M, Dalla Vestra M, Fioretto P, Saller A, Varagnolo M, Nosadini R, et al. Cystatin C is a more sensitive marker than creatinine for the estimation of GFR in type 2 diabetic patients. *Kidney Int* 2002;61:1453-61.
- 1415 *Mussap M, Ruzzante N, Varagnolo M, Plebani M. Quantitative automated particle-enhanced immunonephelometric assay for the routine measurement of human cystatin C. *Clin Chem Lab Med* 1998;36:859-65.
- 6900 *Myara I, Lahiani F, Cosson C, Duboust A, Moatti N. Estimated creatinine clearance by the formula of Gault and Cockcroft in renal transplantation. *Nephron* 1989;51:426-7.
- 6908 *Nagao S, Fujiwara K, Imafuku N, Kagawa R, Kozuka Y, Oda T, et al. Difference of carboplatin clearance estimated by the Cockcroft-Gault, Jelliffe, Modified-Jelliffe, Wright or Chatelut formula. *Gynecol Oncol* 2005;99:327-33.
- 1422 *Nakai K, Kikuchi M, Fujimoto K, Kaneko Y, Omori S, Suwabe A. Serum levels of cystatin C in patients with malignancy. *Clin Exp Nephrol* 2008;12:132-9.
- 6926 *Nankivell BJ, Chapman JR, Allen RD. Predicting glomerular filtration rate after simultaneous pancreas and kidney transplantation. *Clin Transplant* 1995;9:129-34.

6927 *Nankivell BJ, Gruenewald SM, Allen RD, Chapman JR. Predicting glomerular filtration rate after kidney transplantation. *Transplantation* 1995;59:1683-9.

14051 *Nehus EJ, Ross J, Laskin BL, Goebel J. Cystatin C and serum creatinine similarly estimate graft function in pediatric kidney transplant recipients. *American Journal of Transplantation* 2011;11:494-495.

6954 *Newman DJ, Thakkar H, Edwards RG, Wilkie M, White T, Grubb AO, et al. Serum cystatin C measured by automated immunoassay: a more sensitive marker of changes in GFR than serum creatinine. *Kidney Int* 1995;47:312-8.

14052 *Ng DKS, Schwartz GJ, Jacobson LP, Palella FJ, Margolick JB, Warady BA, et al. Universal GFR determination based on two time points during plasma iothexol disappearance. *Kidney Int* 2011;80:423-430.

6956 *Nguyen HT, Shannon AG, Coates PA, Owens DR. Estimation of glomerular filtration rate in type II (non-insulin dependent) diabetes mellitus patients. *IMA J Math Appl Med Biol* 1997;14:151-60.

13400 *Nguyen MT, Maynard SE, Kimmel PL. Misapplications of commonly used kidney equations: Renal physiology in practice. *Clinical Journal of the American Society of Nephrology* 2009;4:528-534.

6965 *Nicoll SR, Sainsbury R, Bailey RR, King A, Frampton C, Elliot JR, et al. Assessment of creatinine clearance in healthy subjects over 65 years of age. *Nephron* 1991;59:621-5.

6969 *Nielsen S, Rehling M, Schmitz A, Mogensen CE. Validity of rapid estimation of glomerular filtration rate in type 2 diabetic patients with normal renal function. *Nephrol Dial Transplant* 1999;14:615-9.

14053 *Nigam R, Mathur M. A comparison of estimated creatinine clearance and measured glomerular filtration rate (TC99MDTPA clearance) in Indians. *Clinical Chemistry and Laboratory Medicine* 2011;49:S566.

6974 *Nilsson-Ehle P, Grubb A. New markers for the determination of GFR: iothexol clearance and cystatin C serum concentration. *Kidney Int Suppl* 1994;47:S17-9.

6982 *Nix DE, Erstad BL, Nakazato PZ, Barletta JF, Matthias KR, Krueger TS. Estimation of creatinine clearance in end-stage liver disease. *Ann Pharmacother* 2006;40:900-8.

6984 *Nobrega AM, Gomes CP, Lemos CC, Bregman R. Is it possible to use modification of diet in renal disease (MDRD) equation in a Brazilian population? *J Nephrol* 2006;19:196-9.

12853 *Nyman U, Bjork J, Lindstrom V, Grubb A. The Lund-Malmo creatinine-based glomerular filtration rate prediction equation for adults also performs well in children. *Scand J Clin Lab Invest* 2008;1-9.

7010 *Nyman U, Bjork J, Sterner G, Back SE, Carlson J, Lindstrom V, et al. Standardization of p-creatinine assays and use of lean body mass allow improved prediction of calculated glomerular filtration rate in adults: a new equation. *Scand J Clin Lab Invest* 2006;66:451-68.

7012 *Nyman U, Grubb A, Sterner G, Bjork J. Different equations to combine creatinine and cystatin C to predict GFR. Arithmetic mean of existing equations performs as well as complex combinations. *Scand J Clin Lab Invest* 2009;69:619-27.

13558 *Nyman U, Grubb A, Sterner G, Bjork J. The CKD-EPI and MDRD equations to estimate GFR. Validation in the Swedish Lund-Malmo Study cohort. *Scand J Clin Lab Invest* 2011;71:129-38.

7015 *Ocheke IE, Agaba EI. Discrepancy between predicted and measured GFR in hospitalized Nigerian children. *J Trop Pediatr* 2006;52:335-40.

7017 *Odoze C, Morange S, Portugal H, Berland Y, Dussol B. Cystatin C is not more sensitive than creatinine for detecting early renal impairment in patients with diabetes. *Am J Kidney Dis* 2001;38:310-6.

13401 *Olsson CL, Rippe B, Bendz H. A comparison of plasma cystatin C and plasma creatinine for the screening of renal function in lithium-treated patients. *Clinical nephrology* 2010;74:132-140.

1450 *O'Riordan SE, Webb MC, Stowe HJ, Simpson DE, Kandarpa M, Coakley AJ, et al. Cystatin C improves the detection of mild renal dysfunction in older patients. *Ann Clin Biochem* 2003;40:648-55.

2163 *Orlando R, Floreani M, Padrini R, Palatini P. Evaluation of measured and calculated creatinine clearances as glomerular filtration markers in different stages of liver cirrhosis. *Clin Nephrol* 1999;51:341-7.

13402 *Orskov B, Borresen ML, Feldt-Rasmussen B, Ostergaard O, Laursen I, Strandgaard S. Estimating glomerular filtration rate using the new CKD-EPI equation and other equations in patients with autosomal dominant polycystic kidney disease. *American journal of nephrology* 2010;31:53-57.

13403 *Owiredu WKBA, Ephraim RKD, Amidu N, Eghan Jr BA, Quaye L. Predictive performance of renal function equations among Ghanaians presenting with chronic kidney disease. *Journal of Medical Sciences* 2008;8:491-497.

7103 *Ozmen S, Kaplan MA, Kaya H, Akin D, Danis R, Kizilkan B, et al. Role of lean body mass for estimation of glomerular filtration rate in patients with chronic kidney disease with various body mass indices. *Scand J Urol Nephrol* 2009;43:171-6.

13404 *Pai MP. Estimating the Glomerular Filtration Rate in Obese Adult Patients for Drug Dosing. *Advances in Chronic Kidney Disease* 2010;17:e53-e62.

7109 *Pai MP, Norenberg JP, Anderson T, Goade DW, Rodvold KA, Telepak RA, et al. Influence of morbid obesity on the single-dose pharmacokinetics of daptomycin. *Antimicrob Agents Chemother* 2007;51:2741-7.

7115 *Paleologo G, Abdelkawy H, Barsotti M, Basha A, Bernabini G, Bianchi A, et al. Kidney dimensions at sonography are correlated with glomerular filtration rate in renal transplant recipients and in kidney donors. *Transplant Proc* 2007;39:1779-81.

14054 *Pande SA, Malik UM, Mumtaz K, Jain M, Dissanayake I, Bloom EJ, et al. Estimated GFR as a measure of dialytic clearance in patients on peritoneal dialysis. *Blood Purification* 2011;31:223.

14055 *Park EJ, Pai MP, Zhang J, Dong T, Ko CW, Crentsil V, et al. Estimation of creatinine clearance (ECLCR) across all weight strata. *Clinical Pharmacology and Therapeutics* 2011;89:S32-S33.

7159 *Pedone C, Semeraro R, Chiurco D, D'Andria F, Gigante M, Coppola A, et al. Reliability of equations to estimate glomerular filtration rate in the very old. *Aging Clin Exp Res* 2008;20:496-502.

7165 *Pequignot R, Belmin J, Chauvelier S, Gaubert JY, Konrat C, Duron E, et al. Renal Function in Older Hospital Patients Is More Accurately Estimated Using the Cockcroft-Gault Formula Than the Modification Diet in Renal Disease Formula. *J Am Geriatr Soc* 2009.

7177 *Perlemoine C, Beauvieux MC, Rigalleau V, Baillet L, Barthes N, Derache P, et al. Interest of cystatin C in screening diabetic patients for early impairment of renal function. *Metabolism* 2003;52:1258-64.

1472 *Peters AM, Bird NJ, Halsall I, Peters C, Michell AR. Evaluation of the Modification of Diet in Renal Disease equation (eGFR) against simultaneous, dual-marker multi-sample measurements of glomerular filtration rate. *Ann Clin Biochem* 2009;46:58-64.

7198 *Pham-Huy A, Leonard M, Lepage N, Halton J, Filler G. Measuring glomerular filtration rate with cystatin C and beta-trace protein in children with spina bifida. *J Urol* 2003;169:2312-5.

13841 *Pieroni L, Delanaye P, Boutten A, Bargnoux AS, Rozet E, Delatour V, et al. A multicentric evaluation of IDMS-traceable creatinine enzymatic assays. *Clin Chim Acta* 2011;412:2070-2075.

2181 *Pierrat A, Gravier E, Saunders C, Caira MV, Ait-Djafer Z, Legras B, et al. Predicting GFR in children and adults: a comparison of the Cockcroft-Gault, Schwartz, and modification of diet in renal disease formulas. *Kidney Int* 2003;64:1425-36.

7221 *Planet G. Comparison of creatinine clearance estimates with routine measured clearance. *Clin Biochem* 2007;40:124-7.

- 2186 *Plebani M, Dall'Amico R, Mussap M, Montini G, Ruzzante N, Marsilio R, et al. Is serum cystatin C a sensitive marker of glomerular filtration rate (GFR)? A preliminary study on renal transplant patients. *Ren Fail* 1998;20:303-9.
- 7226 *Podracka L, Feber J, Lepage N, Filler G. Intra-individual variation of cystatin C and creatinine in pediatric solid organ transplant recipients. *Pediatr Transplant* 2005;9:28-32.
- 2190 *Poge U, Gerhardt T, Stoffel-Wagner B, Klehr HU, Sauerbruch T, Woitas RP. Calculation of glomerular filtration rate based on cystatin C in cirrhotic patients. *Nephrol Dial Transplant* 2006;21:660-4.
- 7228 *Poge U, Gerhardt T, Stoffel-Wagner B, Palmedo H, Klehr HU, Sauerbruch T, et al. Cystatin C-based calculation of glomerular filtration rate in kidney transplant recipients. *Kidney Int* 2006;70:204-10.
- 7229 *Poge U, Gerhardt T, Stoffel-Wagner B, Palmedo H, Klehr HU, Sauerbruch T, et al. Prediction of glomerular filtration rate in renal transplant recipients: cystatin C or modification of diet in renal disease equation? *Clin Transplant* 2006;20:200-5.
- 7230 *Poge U, Gerhardt T, Stoffel-Wagner B, Palmedo H, Klehr HU, Sauerbruch T, et al. Can modifications of the MDRD formula improve the estimation of glomerular filtration rate in renal allograft recipients? *Nephrol Dial Transplant* 2007;22:3610-5.
- 7232 *Poge U, Gerhardt T, Woitas RP. Equations to estimate GFR using serum cystatin C in kidney transplant recipients. *Am J Kidney Dis* 2008;52:383-4.
- 7234 *Poge U, Stoschus B, Stoffel-Wagner B, Gerhardt T, Klehr HU, Sauerbruch T, et al. Cystatin C as an endogenous marker of glomerular filtration rate in renal transplant patients. *Kidney Blood Press Res* 2003;26:55-60.
- 7236 *Poggio ED, Nef PC, Wang X, Greene T, Van Lente F, Dennis VW, et al. Performance of the Cockcroft-Gault and modification of diet in renal disease equations in estimating GFR in ill hospitalized patients. *Am J Kidney Dis* 2005;46:242-52.
- 1485 *Poggio ED, Rule AD. Can we do better than a single estimated GFR threshold when screening for chronic kidney disease? *Kidney Int* 2007;72:534-6.
- 1486 *Poggio ED, Wang X, Greene T, Van Lente F, Hall PM. Performance of the modification of diet in renal disease and Cockcroft-Gault equations in the estimation of GFR in health and in chronic kidney disease. *J Am Soc Nephrol* 2005;16:459-66.
- 7238 *Poggio ED, Wang X, Weinstein DM, Issa N, Dennis VW, Braun WE, et al. Assessing glomerular filtration rate by estimation equations in kidney transplant recipients. *Am J Transplant* 2006;6:100-8.
- 7255 *Poole SG, Dooley MJ, Rischin D. A comparison of bedside renal function estimates and measured glomerular filtration rate (Tc99mDTPA clearance) in cancer patients. *Ann Oncol* 2002;13:949-55.
- 13405 *Pottel H, Hoste L, Martens F. New insights in glomerular filtration rate formulas and chronic kidney disease classification. *Clinica Chimica Acta* 2010;411:1341-1347.
- 13406 *Pottel H, Mottaghy FM, Zaman Z, Martens F. On the relationship between glomerular filtration rate and serum creatinine in children. *Pediatric Nephrology* 2010;25:927-934.
- 13861 *Praditpornsilpa K, Townamchai N, Chaiwatanarat T, Tiranathanagul K, Katawatin P, Susantitaphong P, et al. The need for robust validation for MDRD-based glomerular filtration rate estimation in various CKD populations. *Nephrol Dial Transplant* 2011;26:2780-5.
- 7285 *Premaratne E, Maclsaac RJ, Finch S, Panagiotopoulos S, Ekinci E, Jerums G. Serial measurements of cystatin C are more accurate than creatinine-based methods in detecting declining renal function in type 1 diabetes. *Diabetes Care* 2008;31:971-3.
- 14056 *Prestidge C, Chilvers MA, Davidson AGF, Cho E, McMahon V, White CT. Renal function in pediatric cystic fibrosis patients in the first decade of life. *Pediatric Nephrology* 2011;26:605-612.

2206 *Price M. Comparison of creatinine clearance to inulin clearance in the determination of glomerular filtration rate. *J Urol* 1972;107:339-40.

2207 *Price M, Van Pilsum JF, Markland C, Boen JR. Comparison of creatinine clearance with inulin clearance in paraplegic patients. *Proc Annu Clin Spinal Cord Inj Conf* 1967;16:176-80.

7295 *Prigent A. Monitoring renal function and limitations of renal function tests. *Semin Nucl Med* 2008;38:32-46.

13407 *Probst LA, Darko W, Smith A, Cwikla GM. Pitfalls of the application of the modification of diet in renal disease equation to drug-dosing practices: A tertiary care teaching hospital experience. *Hospital Pharmacy* 2008;43:564-570.

2209 *Proulx NL, Akbari A, Garg AX, Rostom A, Jaffey J, Clark HD. Measured creatinine clearance from timed urine collections substantially overestimates glomerular filtration rate in patients with liver cirrhosis: a systematic review and individual patient meta-analysis. *Nephrol Dial Transplant* 2005;20:1617-22.

7301 *Pucci L, Triscornia S, Lucchesi D, Fotino C, Pellegrini G, Pardini E, et al. Cystatin C and estimates of renal function: searching for a better measure of kidney function in diabetic patients. *Clin Chem* 2007;53:480-8.

13862 *Quinn MP, Cardwell CR, Kee F, Maxwell AP, Savage G, McCarron P, et al. The finding of reduced estimated glomerular filtration rate is associated with increased mortality in a large UK population. *Nephrol Dial Transplant* 2011;26:875-80.

7317 *Quinn MP, Rainey A, Cairns KJ, Marshall AH, Savage G, Kee F, et al. The practical implications of using standardized estimation equations in calculating the prevalence of chronic kidney disease. *Nephrol Dial Transplant* 2008;23:542-8.

7324 *Rafey MA, Lipkowitz MS. How accurate are formulas that calculate creatinine clearance? *Nat Clin Pract Urol* 2007;4:188-9.

7332 *Raj GV, Iasonos A, Herr H, Donat SM. Formulas calculating creatinine clearance are inadequate for determining eligibility for Cisplatin-based chemotherapy in bladder cancer. *J Clin Oncol* 2006;24:3095-100.

7336 *Raju DL, Grover VK, Shoker A. Limitations of glomerular filtration rate equations in the renal transplant patient. *Clin Transplant* 2005;19:259-68.

14057 *Ramanathan K, Padmanaban G. Serum cystatin C a new marker of glomerular filtration rate. *Biosciences Biotechnology Research Asia* 2011;8:345-347.

7345 *Randers E, Kristensen JH, Erlandsen EJ, Danielsen H. Serum cystatin C as a marker of the renal function. *Scand J Clin Lab Invest* 1998;58:585-92.

7350 *Rashid ST, Salman M, Agarwal S, Hamilton G. Occult renal impairment is common in patients with peripheral vascular disease and normal serum creatinine. *Eur J Vasc Endovasc Surg* 2006;32:294-9.

7360 *Ravasi G, Lauriola M, Tinelli C, Brandolini M, Uglietti A, Maserati R. Comparison of glomerular filtration rate estimates vs. 24-h creatinine clearance in HIV-positive patients. *HIV Med* 2009;10:219-28.

14058 *Riad S, Kukla AK, Leister EC, Matas AJ, Ibrahim HN. Utility of the CKD-EPI equation in estimating GFR in former kidney donors. *American Journal of Transplantation* 2011;11:310.

7414 *Rigalleau V, Beauvieux MC, Le Moigne F, Lasseur C, Chauveau P, Raffaitin C, et al. Cystatin C improves the diagnosis and stratification of chronic kidney disease, and the estimation of glomerular filtration rate in diabetes. *Diabetes Metab* 2008;34:482-9.

13408 *Rigalleau V, Chauveau P, Lasseur C, Raffaitin C, Perlemoine C, Barthe N, et al. Glomerular filtration rate prediction using lean mass is unsuccessful in diabetic subjects [3]. *Nephrology Dialysis Transplantation* 2006;21:1443-1444.

- 7416 *Rigalleau V, Lasseur C, Perlemoine C, Barthe N, Raffaitin C, de La Faille R, et al. A simplified Cockcroft-Gault formula to improve the prediction of the glomerular filtration rate in diabetic patients. *Diabetes Metab* 2006;32:56-62.
- 7417 *Rigalleau V, Lasseur C, Perlemoine C, Barthe N, Raffaitin C, Liu C, et al. Estimation of glomerular filtration rate in diabetic subjects: Cockcroft formula or modification of Diet in Renal Disease study equation? *Diabetes Care* 2005;28:838-43.
- 1510 *Rigalleau V, Lasseur C, Raffaitin C, Beauvieux MC, Barthe N, Chauveau P, et al. Normoalbuminuric renal-insufficient diabetic patients: a lower-risk group. *Diabetes Care* 2007;30:2034-9.
- 13409 *Rigalleau V, Lasseur C, Raffaitin C, Perlemoine C, Barthe N, Chauveau P, et al. The Mayo Clinic quadratic equation improves the prediction of glomerular filtration rate in diabetic subjects. *Nephrology Dialysis Transplantation* 2007;22:813-818.
- 7420 *Rimon E, Kagansky N, Cojocaru L, Gindin J, Schattner A, Levy S. Can creatinine clearance be accurately predicted by formulae in octogenarian in-patients? *QJM* 2004;97:281-7.
- 7425 *Risch L, Blumberg A, Huber A. Rapid and accurate assessment of glomerular filtration rate in patients with renal transplants using serum cystatin C. *Nephrol Dial Transplant* 1999;14:1991-6.
- 7427 *Risch L, Drexel H, Huber AR. Differences in glomerular filtration rate estimates by 2 cystatin C-based equations. *Clin Chem* 2005;51:2211-2.
- 7428 *Risch L, Huber AR. Assessing glomerular filtration rate in renal transplant recipients by estimates derived from serum measurements of creatinine and cystatin C. *Clin Chim Acta* 2005;356:204-11.
- 2226 *Robert S, Zarowitz BJ. Is there a reliable index of glomerular filtration rate in critically ill patients? *DICP* 1991;25:169-78.
- 2227 *Robert S, Zarowitz BJ, Peterson EL, Dumler F. Predictability of creatinine clearance estimates in critically ill patients. *Crit Care Med* 1993;21:1487-95.
- 7441 *Robertshaw M, Lai KN, Swaminathan R. Prediction of creatinine clearance from plasma creatinine: comparison of five formulae. *Br J Clin Pharmacol* 1989;28:275-80.
- 7448 *Rodby RA, Ali A, Rohde RD, Lewis EJ. Renal scanning 99mTc diethylene-triamine pentaacetic acid glomerular filtration rate (GFR) determination compared with iothalamate clearance GFR in diabetics. The Collaborative Study Group for The study of Angiotensin-Converting Enzyme Inhibition in Diabetic Nephropathy. *Am J Kidney Dis* 1992;20:569-73.
- 7454 *Rodrigo E, Fernandez-Fresnedo G, Ruiz JC, Pinera C, Heras M, de Francisco AL, et al. Assessment of glomerular filtration rate in transplant recipients with severe renal insufficiency by Nankivell, Modification of Diet in Renal Disease (MDRD), and Cockcroft-Gault equations. *Transplant Proc* 2003;35:1671-2.
- 7470 *Rolin HA, 3rd, Hall PM, Wei R. Inaccuracy of estimated creatinine clearance for prediction of iothalamate glomerular filtration rate. *Am J Kidney Dis* 1984;4:48-54.
- 2244 *Roos JF, Doust J, Tett SE, Kirkpatrick CM. Diagnostic accuracy of cystatin C compared to serum creatinine for the estimation of renal dysfunction in adults and children--a meta-analysis. *Clin Biochem* 2007;40:383-91.
- 7501 *Rossing P, Astrup AS, Smidt UM, Parving HH. Monitoring kidney function in diabetic nephropathy. *Diabetologia* 1994;37:708-12.
- 7502 *Rossing P, Rossing K, Gaede P, Pedersen O, Parving HH. Monitoring kidney function in type 2 diabetic patients with incipient and overt diabetic nephropathy. *Diabetes Care* 2006;29:1024-30.
- 14059 *Ruggenenti P, Remuzzi G. Chronic kidney disease: Staging CKD: Can we rely on estimated GFRs? *Nature Reviews Nephrology* 2011;7:307-308.

13410 *Rule AD. The CKD-EPI equation for estimating GFR from serum creatinine: Real improvement or more of the same? *Clinical Journal of the American Society of Nephrology* 2010;5:951-953.

7538 *Rule AD, Bailey KR, Schwartz GL, Khosla S, Lieske JC, Melton LJ, 3rd. For estimating creatinine clearance measuring muscle mass gives better results than those based on demographics. *Kidney Int* 2009;75:1071-8.

7539 *Rule AD, Bergstralh EJ, Slezak JM, Bergert J, Larson TS. Glomerular filtration rate estimated by cystatin C among different clinical presentations. *Kidney Int* 2006;69:399-405.

1523 *Rule AD, Gussak HM, Pond GR, Bergstralh EJ, Stegall MD, Cosio FG, et al. Measured and estimated GFR in healthy potential kidney donors. *Am J Kidney Dis* 2004;43:112-9.

13411 *Rule AD, Gussak HM, Pond GR, Bergstralh EJ, Stegall MD, Cosio FG, et al. Transplantation Measured and Estimated GFR in Healthy Potential Kidney Donors. *American Journal of Kidney Diseases* 2004;43:112-119.

13412 *Rule AD, Larson TS, Bergstralh EJ, Sequist TD. Popular prediction equation underestimates glomerular filtration rate. *Journal of Clinical Outcomes Management* 2005;12:68-74.

1524 *Rule AD, Larson TS, Bergstralh EJ, Slezak JM, Jacobsen SJ, Cosio FG. Using serum creatinine to estimate glomerular filtration rate: accuracy in good health and in chronic kidney disease. *Ann Intern Med* 2004;141:929-37.

7546 *Rule AD, Torres VE, Chapman AB, Grantham JJ, Guay-Woodford LM, Bae KT, et al. Comparison of methods for determining renal function decline in early autosomal dominant polycystic kidney disease: the consortium of radiologic imaging studies of polycystic kidney disease cohort. *J Am Soc Nephrol* 2006;17:854-62.

7550 *Rumpfeld M, Johnson DW. A population-specific formula predicting creatinine excretion in continuous peritoneal dialysis. *Perit Dial Int* 2003;23:406-8; author reply 408.

7569 *Saleem M, Florkowski CM, George PM. Comparison of the Mayo Clinic Quadratic Equation with the Modification of Diet in Renal Disease equation and radionuclide glomerular filtration rate in a clinical setting. *Nephrology (Carlton)* 2008;13:684-8.

7570 *Saleem M, Florkowski CM, George PM, Woltersdorf WW. Comparison of two prediction equations with radionuclide glomerular filtration rate: validation in routine use. *Ann Clin Biochem* 2006;43:309-13.

13413 *Salgado JV, Neves FA, Bastos MG, Franca AK, Brito DJ, Santos EM, et al. Monitoring renal function: Measured and estimated glomerular filtration rates - A review. *Brazilian Journal of Medical and Biological Research* 2010;43:528-536.

7573 *Sampson MJ, Drury PL. Accurate estimation of glomerular filtration rate in diabetic nephropathy from age, body weight, and serum creatinine. *Diabetes Care* 1992;15:609-12.

7576 *Samyn M, Cheeseman P, Bevis L, Taylor R, Samaroo B, Buxton-Thomas M, et al. Cystatin C, an easy and reliable marker for assessment of renal dysfunction in children with liver disease and after liver transplantation. *Liver Transpl* 2005;11:344-9.

2258 *Sansoe G, Ferrari A, Castellana CN, Bonardi L, Villa E, Manenti F. Cimetidine administration and tubular creatinine secretion in patients with compensated cirrhosis. *Clin Sci (Lond)* 2002;102:91-8.

7588 *Sanusi AA, Akinsola A, Ajayi AA. Creatinine clearance estimation from serum creatinine values: evaluation and comparison of five prediction formulae in Nigerian patients. *Afr J Med Med Sci* 2000;29:7-11.

7591 *Saracino A, Morrone LF, Suriano V, Niccoli-Asabella A, Ramunni A, Fanelli M, et al. A simple method for correcting overestimated glomerular filtration rate in obese subjects evaluated by the Cockcroft and Gault formula: a comparison with ⁵¹Cr EDTA clearance. *Clin Nephrol* 2004;62:97-103.

13414 *Schaeffner ES, Van Der Giet M, Gaedeke J, Tolle M, Ebert N, Kuhlmann MK, et al. The Berlin initiative study: The methodology of exploring kidney function in the elderly by combining a longitudinal and cross-sectional approach. *European Journal of Epidemiology* 2010;25:203-210.

- 7621 *Schiff J, Paraskevas S, Keith D, Tchervenkov J, Metrakos P, Cantarovich M. Prediction of the glomerular filtration rate using equations in kidney-pancreas transplant patients receiving cimetidine. *Transplantation* 2006;81:469-72.
- 7646 *Schuck O, Horackova M. Relationship between an increase in serum creatinine or a doubling in serum creatinine and a decrease in glomerular filtration rate estimated using prediction equations. *Int J Clin Pharmacol Ther* 2008;46:40-1.
- 2278 *Schuck O, Teplan V, Mareckova O, Skibova J, Stolova M. MDRD formulas for GFR estimation. Is there any difference among them in prediction of renal inulin clearance? *Clin Nephrol* 2005;64:326-7.
- 2279 *Schuck O, Teplan V, Mareckova O, Skibova J, Stolova M. Estimation of glomerular filtration rate based on the modification of diet in renal disease equation in patients with chronic renal failure. *Kidney Blood Press Res* 2005;28:63-7.
- 2283 *Schuck O, Teplan V, Sibova J, Stolova M. Predicting the glomerular filtration rate from serum creatinine, serum cystatin C and the Cockcroft and Gault formula with regard to drug dosage adjustment. *Int J Clin Pharmacol Ther* 2004;42:93-7.
- 13415 *Schut A, Anguelov A, Aussel C, Mameri S, Royand F, Vincent JP. Commonly used estimates of creatinine clearance are inadequate for the very elderly. *Journal of Nutrition, Health and Aging* 2009;13:145-148.
- 7664 *Schwartz GJ. Does kL/PCr estimate GFR, or does GFR determine k? *Pediatr Nephrol* 1992;6:512-5.
- 1553 *Schwartz GJ, Brion LP, Spitzer A. The use of plasma creatinine concentration for estimating glomerular filtration rate in infants, children, and adolescents. *Pediatr Clin North Am* 1987;34:571-90.
- 7666 *Schwartz GJ, Feld LG, Langford DJ. A simple estimate of glomerular filtration rate in full-term infants during the first year of life. *J Pediatr* 1984;104:849-54.
- 7667 *Schwartz GJ, Furth S, Cole SR, Warady B, Munoz A. Glomerular filtration rate via plasma iothexol disappearance: pilot study for chronic kidney disease in children. *Kidney Int* 2006;69:2070-7.
- 155 *Schwartz GJ, Furth SL. Glomerular filtration rate measurement and estimation in chronic kidney disease. *Pediatr Nephrol* 2007;22:1839-48.
- 7669 *Schwartz GJ, Gauthier B. A simple estimate of glomerular filtration rate in adolescent boys. *J Pediatr* 1985;106:522-6.
- 16 *Schwartz GJ, Haycock GB, Edelmann CM, Jr., Spitzer A. A simple estimate of glomerular filtration rate in children derived from body length and plasma creatinine. *Pediatrics* 1976;58:259-63.
- 4239 *Schwartz GJ, Munoz A, Schneider MF, Mak RH, Kaskel F, Warady BA, et al. New equations to estimate GFR in children with CKD. *J Am Soc Nephrol* 2009;20:629-37.
- 13245 *Schwartz GJ, Work DF. Measurement and estimation of GFR in children and adolescents. *Clin J Am Soc Nephrol*. 2009;4:1832-43.
- 7682 *Sebasky M, Kukla A, Leister E, Guo H, Akkina SK, El-Shahawy Y, et al. Appraisal of GFR-estimating equations following kidney donation. *Am J Kidney Dis* 2009;53:1050-8.
- 13795 *Segarra A, de la Torre J, Ramos N, Quiroz A, Garjau M, Torres I, et al. Assessing Glomerular Filtration Rate in Hospitalized Patients: A Comparison Between CKD-EPI and Four Cystatin C-Based Equations. *Clin J Am Soc Nephrol* 2011;6:2411-20.
- 7694 *Seikaly MG, Browne R, Bajaj G, Arant BS, Jr. Limitations to body length/serum creatinine ratio as an estimate of glomerular filtration in children. *Pediatr Nephrol* 1996;10:709-11.
- 7695 *Seikaly MG, Browne R, Simonds N, Atkins C, Alexander SR. Glomerular filtration rate in children following renal transplantation. *Pediatr Transplant* 1998;2:231-5.

- 7696 *Sekar KC, Leonard JC, Puffinburger WR, Burow DM, Duke JC. Rapid determination of glomerular filtration rate in infants using Tc-99m DTPA. *Clin Nucl Med* 1992;17:550-2.
- 14060 *Senulkowska R, Roszkowska-Blaim M. A modification of Schwartz formula for enzymatic creatinine assay. *Pediatric Nephrology* 2011;26:1610.
- 7707 *Serdar MA, Kurt I, Ozcelik F, Urhan M, Ilgan S, Yenicesu M, et al. A practical approach to glomerular filtration rate measurements: creatinine clearance estimation using cimetidine. *Ann Clin Lab Sci* 2001;31:265-73.
- 7710 *Seronie-Vivien S, Bouissou F, Dattez S, Coulais Y, Hanser AM, Hym B, et al. Modeling the variability of creatinine measurements improves estimates of the glomerular filtration rate. *Clin Chem Lab Med* 2008;46:215-8.
- 7711 *Seronie-Vivien S, Delanaye P, Pieroni L, Mariat C, Froissart M, Cristol JP, et al. Cystatin C: current position and future prospects. *Clin Chem Lab Med* 2008;46:1664-86.
- 7715 *Seronie-Vivien S, Toullec S, Malard L, Thomas F, Durrand V, Chatelut E. Contribution of the MDRD equation and of cystatin C for renal function estimates in cancer patients. *Med Oncol* 2006;23:63-73.
- 13252 *Sharma AP, Filler G. Monitoring kidney function and renal disease in children following transplant. *Pediatric Health* 2009;3:155-163.
- 7729 *Sharma AP, Kathiravelu A, Nadarajah R, Yasin A, Filler G. Body mass does not have a clinically relevant effect on cystatin C eGFR in children. *Nephrol Dial Transplant* 2009;24:470-4.
- 13863 *Sharma AP, Yasin A, Garg AX, Filler G. Diagnostic accuracy of cystatin C-based eGFR equations at different GFR levels in children. *Clin J Am Soc Nephrol* 2011;6:1599-608.
- 13664 *Shikanov S, Clark MA, Raman JD, Smith B, Kaag M, Russo P, et al. Chronic kidney disease epidemiology collaboration versus modification of diet in renal disease equations for renal function evaluation in patients undergoing partial nephrectomy. *Journal of Urology* 2010;184:1867-1871.
- 7764 *Shoker A, Hossain MA, Koru-Sengul T, Raju DL, Cockcroft D. Performance of creatinine clearance equations on the original Cockcroft-Gault population. *Clin Nephrol* 2006;66:89-97.
- 7767 *Shull BC, Haughey D, Koup JR, Baliah T, Li PK. A useful method for predicting creatinine clearance in children. *Clin Chem* 1978;24:1167-9.
- 13864 *Silveiro SP, Araujo GN, Ferreira MN, Souza FD, Yamaguchi HM, Camargo EG. Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) Equation Pronouncedly Underestimates Glomerular Filtration Rate in Type 2 Diabetes. *Diabetes Care* 2011.
- 13865 *Simon J, Amde M, Poggio ED. Interpreting the estimated glomerular filtration rate in primary care: benefits and pitfalls. *Cleve Clin J Med* 2011;78:189-95.
- 7783 *Simonsen O, Grubb A, Thysell H. The blood serum concentration of cystatin C (gamma-trace) as a measure of the glomerular filtration rate. *Scand J Clin Lab Invest* 1985;45:97-101.
- 7788 *Sinton TJ, De Leacy EA, Cowley DM. Comparison of 51Cr EDTA clearance with formulae in the measurement of glomerular filtration rate. *Pathology* 1986;18:445-7.
- 13262 *Sirwal IA, Banday KA, Reshi AR, Bhat MA, Wani MM. Estimation of glomerular filtration rate (GFR). *JK Science* 2004;6:121-123.
- 7793 *Sjostrom PA, Jones IL, Tidman MA. Cystatin C as a filtration marker--haemodialysis patients expose its strengths and limitations. *Scand J Clin Lab Invest* 2009;69:65-72.
- 7792 *Sjöström P, Tidman M, Jones I. Determination of the production rate and non-renal clearance of cystatin C and estimation of the glomerular filtration rate from the serum concentration of cystatin C in humans. *Scand J Clin Lab Invest* 2005;65:111-24.

- 7799 *Skinner R, Cole M, Pearson AD, Keir MJ, Price L, Wyllie RA, et al. Inaccuracy of glomerular filtration rate estimation from height/plasma creatinine ratio. *Arch Dis Child* 1994;70:387-90.
- 7801 *Skruzacek PA, Szewc RG, Nolan CR, 3rd, Riley DJ, Lee S, Pergola PE. Prediction of GFR in liver transplant candidates. *Am J Kidney Dis* 2003;42:1169-76.
- 7806 *Sladen RN. Accurate estimation of glomerular filtration in the intensive care unit: another Holy Grail? *Crit Care Med* 1993;21:1424-7.
- 14061 *Sladen RN. Chronic kidney disease: The silent enemy? *Anesth Analg* 2011;112:1277-1279.
- 7812 *Smilde TD, van Veldhuisen DJ, Navis G, Voors AA, Hillege HL. Drawbacks and prognostic value of formulas estimating renal function in patients with chronic heart failure and systolic dysfunction. *Circulation* 2006;114:1572-80.
- 7819 *Smith JS, Roizen MF, Cahalan MK, Benefiel DJ, Beaupre PN, Sohn YJ, et al. Does anesthetic technique make a difference? Augmentation of systolic blood pressure during carotid endarterectomy: effects of phenylephrine versus light anesthesia and of isoflurane versus halothane on the incidence of myocardial ischemia. *Anesthesiology* 1988;69:846-53.
- 2307 *Smith MC, Moran P, Ward MK, Davison JM. Assessment of glomerular filtration rate during pregnancy using the MDRD formula. *BJOG* 2008;115:109-12.
- 7821 *Smith SA. Estimation of glomerular filtration rate from the serum creatinine concentration. *Postgrad Med J* 1988;64:204-8.
- 13416 *Soares A, Eyff T, Campani R, Ritter L, Weinert L, Camargo J, et al. Performance of the CKD Epidemiology Collaboration (CKD-EPI) and the Modification of Diet in Renal Disease (MDRD) Study Equations in Healthy South Brazilians. *American Journal of Kidney Diseases* 2010;55:1162-1163.
- 7828 *Sobh M, Neamatallah A, Sheashaa H, Akl A, Osman Y, Gad H, et al. Sobh formula: a new formula for estimation of creatinine clearance in healthy subjects and patients with chronic renal disease. *Int Urol Nephrol* 2005;37:403-8.
- 7843 *Sorof JM, Goldstein SL, Brewer ED, Steiger HM, Portman RJ. Serial estimation of glomerular filtration rate in children after renal transplant. *Pediatr Nephrol* 1999;13:737-41.
- 7854 *Springate JE, Christensen SL, Feld LG. Serum creatinine level and renal function in children. *Am J Dis Child* 1992;146:1232-5.
- 7857 *Srinivas S, Annigeri RA, Mani MK, Rao BS, Kowdle PC, Seshadri R. Estimation of glomerular filtration rate in South Asian healthy adult kidney donors. *Nephrology (Carlton)* 2008;13:440-6.
- 7859 *Srivastava T, Alon US, Althahabi R, Garg U. Impact of standardization of creatinine methodology on the assessment of glomerular filtration rate in children. *Pediatr Res* 2009;65:113-6.
- 13417 *Staples A, Leblond R, Watkins S, Wong C, Brandt J. Validation of the revised Schwartz estimating equation in a predominantly non-CKD population. *Pediatric Nephrology* 2010;25:2321-2326.
- 13275 *Steffes MW. Estimating GFR From Cystatin C in Serum: Seeking to Enhance Its Clinical Application. *American Journal of Kidney Diseases* 2006;48:842-843.
- 7884 *Sterner G, Bjork J, Carlson J, Grubb A, Nyman U. Validation of a new plasma cystatin C-based formula and the Modification of Diet in Renal Disease creatinine-based formula for determination of glomerular filtration rate. *Scand J Urol Nephrol* 2009;43:242-9.
- 206 *Sterner G, Frennby B, Mansson S, Nyman U, Van Westen D, Almen T. Determining 'true' glomerular filtration rate in healthy adults using infusion of inulin and comparing it with values obtained using other clearance techniques or prediction equations. *Scand J Urol Nephrol* 2008;42:278-85.

- 13665 *Stevens LA, Claybon MA, Schmid CH, Chen J, Horio M, Imai E, et al. Evaluation of the Chronic Kidney Disease Epidemiology Collaboration equation for estimating the glomerular filtration rate in multiple ethnicities. *Kidney Int.* 2011;79:555-62.
- 7890 *Stevens LA, Coresh J, Feldman HI, Greene T, Lash JP, Nelson RG, et al. Evaluation of the modification of diet in renal disease study equation in a large diverse population. *J Am Soc Nephrol* 2007;18:2749-57.
- 13276 *Stevens LA, Coresh J, Greene T, Levey AS. Assessing kidney function--measured and estimated glomerular filtration rate. *N Engl J Med.* 2006;354:2473-83.
- 7891 *Stevens LA, Coresh J, Schmid CH, Feldman HI, Froissart M, Kusek J, et al. Estimating GFR using serum cystatin C alone and in combination with serum creatinine: a pooled analysis of 3,418 individuals with CKD. *Am J Kidney Dis* 2008;51:395-406.
- 13385 *Stevens LA, Levey AS. Measured GFR as a confirmatory test for estimated GFR. *J Am Soc Nephrol* 2009;20:2305-13.
- 13418 *Stevens LA, Levey AS. Current Status and Future Perspectives for CKD Testing. *American Journal of Kidney Diseases* 2009;53:S17-S26.
- 13842 *Stevens LA, Li S, Kurella Tamura M, Chen SC, Vassalotti JA, Norris KC, et al. Comparison of the CKD Epidemiology Collaboration (CKD-EPI) and Modification of Diet in Renal Disease (MDRD) study equations: risk factors for and complications of CKD and mortality in the Kidney Early Evaluation Program (KEEP). *Am J Kidney Dis* 2011;57:S9-16.
- 7894 *Stevens LA, Manzi J, Levey AS, Chen J, Deysher AE, Greene T, et al. Impact of creatinine calibration on performance of GFR estimating equations in a pooled individual patient database. *Am J Kidney Dis* 2007;50:21-35.
- 13419 *Stevens LA, Nolin T, Levey AS. In Reply to 'Estimated GFR for Drug Dosing: A Bedside Formula', 'Drug Dose Adjustments in Patients With Renal Impairment', 'Use of the MDRD Study Equation for Drug Dosing', and 'Estimated GFR vs Creatinine Clearance for Drug Dosing'. *American Journal of Kidney Diseases* 2009;54:985-986.
- 7895 *Stevens LA, Nolin TD, Richardson MM, Feldman HI, Lewis JB, Rodby R, et al. Comparison of drug dosing recommendations based on measured GFR and kidney function estimating equations. *Am J Kidney Dis* 2009;54:33-42.
- 13420 *Stevens LA, Padala S, Levey AS. Advances in glomerular filtration rate-estimating equations. *Current Opinion in Nephrology and Hypertension* 2010;19:298-307.
- 13421 *Stevens LA, Schmid CH, Greene T, Zhang Y, Beck GJ, Froissart M, et al. Comparative performance of the CKD Epidemiology Collaboration (CKD-EPI) and the Modification of Diet in Renal Disease (MDRD) Study equations for estimating GFR levels above 60 mL/min/1.73 m². *American Journal of Kidney Diseases* 2010;56:486-495.
- 13351 *Stevens LA, Schmid CH, Zhang YL, Coresh J, Manzi J, Landis R, et al. Development and validation of GFR-estimating equations using diabetes, transplant and weight. *Nephrol Dial Transplant* 2010;25:449-57.
- 1587 *Stickle D, Cole B, Hock K, Hruska KA, Scott MG. Correlation of plasma concentrations of cystatin C and creatinine to inulin clearance in a pediatric population. *Clin Chem* 1998;44:1334-8.
- 14062 *Stojanoski S, Gjorceva DP, Gruev T, Ristevska-Miceva S, Ristevska N. Impact of thyroid dysfunction on serum Cystatin C, serum Creatinine and Glomerular Filtration Rate. *Macedonian Journal of Medical Sciences* 2011;4:25-30.
- 13422 *Stoves J, Lindley EJ, Barnfield MC, Burniston MT, Newstead CG. MDRD equation estimates of glomerular filtration rate in potential living kidney donors and renal transplant recipients with impaired graft function [5]. *Nephrology Dialysis Transplantation* 2002;17:2036-2037.

7941 *Sun X, Chen Y, Chen X, Wang J, Xi C, Lin S, et al. Change of glomerular filtration rate in healthy adults with aging. *Nephrology (Carlton)* 2009;14:506-13.

14063 *Taal MW. Predicting renal risk in the general population: Do we have the right formula? *Clinical Journal of the American Society of Nephrology* 2011;6:1523-1525.

13866 *Takata Y, Ansai T, Yoshihara A, Miyazaki H. Glomerular filtration rate and 10-year mortality in a 70-year-old community-dwelling Japanese population. *Aging Clin Exp Res* 2011;23:223-30.

1604 *Tan GD, Lewis AV, James TJ, Altmann P, Taylor RP, Levy JC. Clinical usefulness of cystatin C for the estimation of glomerular filtration rate in type 1 diabetes: reproducibility and accuracy compared with standard measures and iohexol clearance. *Diabetes Care* 2002;25:2004-9.

13423 *Tan JC, Ho B, Busque S, Blouch K, Derby G, Efron B, et al. Imprecision of creatinine-based GFR estimates in uninephric kidney donors. *Clinical Journal of the American Society of Nephrology* 2010;5:497-502.

14064 *Taner T, Iqbal CW, Cosio FG, Stegall MD, Prieto M, Ishitani MB. Impact of donor age, obesity and hypertension on remaining kidney volume. *American Journal of Transplantation* 2011;11:308.

7997 *Tangri N, Alam A, Giannetti N, Deedwardes MB, Cantarovich M. Predicting glomerular filtration rate in heart transplant recipients using serum creatinine-based equations with cimetidine. *J Heart Lung Transplant* 2008;27:905-9.

8011 *Taylor GO, Bamgboye EA, Oyediran AB, Longe O. Serum creatinine and prediction formulae for creatinine clearance. *Afr J Med Med Sci* 1982;11:175-81.

8012 *Taylor TP, Wang W, Shrayyef MZ, Cheek D, Hutchison FN, Gadegbeku CA. Glomerular filtration rate can be accurately predicted using lean mass measured by dual-energy X-ray absorptiometry. *Nephrol Dial Transplant* 2006;21:84-7.

13424 *Tello A, Mondress M, Fan Y, Thomas W, Steffes M, Ibrahim H. The utility of serum creatinine based formulas in predicting the glomerular filtration rate in Type 1 diabetics with normal serum creatinine. [abstract no: SU-PO169]. In: *Journal of the American Society of Nephrology*; 2004. p 569a.

8021 *Tenstad O, Roald AB, Grubb A, Aukland K. Renal handling of radiolabelled human cystatin C in the rat. *Scand J Clin Lab Invest* 1996;56:409-14.

13666 *Tent H, Rook M, Stevens LA, Van Son WJ, Van Pelt LJ, Hofker HS, et al. Renal function equations before and after living kidney donation: A within-individual comparison of performance at different levels of renal function. *Clinical Journal of the American Society of Nephrology* 2010;5:1960-1968.

13425 *Teo BW, Xu H, Shuter B, Wang D, Li J, Sinha AK, et al. CKD-EPI equation is more accurate than mdrdequation in a multi-ethnic Asian population. *American Journal of Kidney Diseases* 2010;55:A106.

13426 *Teo BW, Xu H, Shuter B, Wang D, Li J, Sinha AK, et al. Ethnic coefficients are not required for the MDRD-equation in a multi-ethnic Asian population. *American Journal of Kidney Diseases* 2010;55:A106.

14065 *Teo BW, Xu H, Wang D, Li J, Sinha AK, Shuter B, et al. GFR estimating equations in a multiethnic asian population. *American Journal of Kidney Diseases* 2011;58:56-63.

312 *Tessitore N, Lo Schiavo C, Corgnati A, Previato G, Valvo E, Lupo A, et al. 125I-iothalamate and creatinine clearances in patients with chronic renal diseases. *Nephron* 1979;24:41-5.

14066 *Textor SC. Atherosclerotic renal artery stenosis: Flaws in estimated glomerular filtration rate and the problem of progressive kidney injury. *Circulation: Cardiovascular Interventions* 2011;4:213-215.

13291 *Thomas C, Thomas L. Renal failure--measuring the glomerular filtration rate. *Dtsch Arztebl Int.* 2009;106:849-54.

1613 *Thomas L, Huber AR. Renal function--estimation of glomerular filtration rate. *Clin Chem Lab Med* 2006;44:1295-302.

- 8059 *Tidman M, Sjostrom P, Jones I. A Comparison of GFR estimating formulae based upon s-cystatin C and s-creatinine and a combination of the two. *Nephrol Dial Transplant* 2008;23:154-60.
- 13292 *Tidman M, Sjostrom P, Jones I. Plasma cystatin C for estimating residual GFR (rGFR) in dialysis patients. *Nephrol Dial Transplant*. 2008;23:1072-3; author reply 1073.
- 13427 *Tinti F, Umbro I, Mecule A, Rossi M, Merli M, Nofroni I, et al. RIFLE Criteria and Hepatic Function in the Assessment of Acute Renal Failure in Liver Transplantation. *Transplantation proceedings* 2010;42:1233-1236.
- 13428 *Toffaletti JG. Improving the clinical value of estimating glomerular filtration rate by reporting all values: A contrarian viewpoint. *Nephron - Clinical Practice* 2010;115:c177-c181.
- 13429 *Tomer A, Odar-Cederlof I, Kallner A, Akner G. Renal function in community-dwelling frail elderly. Comparison between measured and predicted glomerular filtration rate in the elderly and proposal for a new cystatin C-based prediction equation. *Aging - Clinical and Experimental Research* 2008;20:216-225.
- 13430 *Tondel C, Ramaswami U, Aakre KM, Wijburg F, Bouwman M, Svarstad E. Monitoring renal function in children with Fabry disease: Comparisons of measured and creatinine-based estimated glomerular filtration rate. *Nephrology Dialysis Transplantation* 2010;25:1507-1513.
- 4197 *Torner A, Odar-Cederlof I, Kallner A, Akner G. Renal function in community-dwelling frail elderly. Comparison between measured and predicted glomerular filtration rate in the elderly and proposal for a new cystatin C-based prediction equation. *Aging Clin Exp Res* 2008;20:216-25.
- 2355 *Toto RD. Conventional measurement of renal function utilizing serum creatinine, creatinine clearance, inulin and para-aminohippuric acid clearance. *Curr Opin Nephrol Hypertens* 1995;4:505-9; discussion 503-4.
- 8090 *Toto RD, Kirk KA, Coresh J, Jones C, Appel L, Wright J, et al. Evaluation of serum creatinine for estimating glomerular filtration rate in African Americans with hypertensive nephrosclerosis: results from the African-American Study of Kidney Disease and Hypertension (AASK) Pilot Study. *J Am Soc Nephrol* 1997;8:279-87.
- 14067 *Tragulpiankit P, Montakantikul P, Chulavatnatol S, Chansirikarnjana S, Thawinwang A. Measured versus estimated creatinine clearance in elderly patients. *Journal of Applied Therapeutic Research* 2011;8:43-51.
- 8104 *Traynor JP, McManus SK, Mactier RA, Scottish Renal R. Derived equations are not precise enough to predict the adequacy of creatinine clearance in peritoneal dialysis patients. *Am J Kidney Dis* 2002;40:1036-44.
- 8110 *Trollfors B, Alestig K, Jagenburg R. Prediction of glomerular filtration rate from serum creatinine, age, sex and body weight. *Acta Med Scand* 1987;221:495-8.
- 8111 *Trollfors B, Alestig K, Jagenburg R. Prediction of glomerular filtration rate from changes in serum creatinine. *Scand J Clin Lab Invest* 1988;48:85-9.
- 4251 *Tsinalis D, Thiel GT. An easy to calculate equation to estimate GFR based on inulin clearance. *Nephrol Dial Transplant* 2009;24:3055-61.
- 13431 *Tsinalis D, Thiel GT. Reply. *Nephrology Dialysis Transplantation* 2010;25:1346-1347.
- 13432 *Tsochatzis E, Manousou P, Thomas M, Jones K, Fatourou E, Fede G, et al. Systematic error in MELD calculation by using serum creatinine measurement in cirrhosis. *SOURCE Hepatology* (2009) 50 (597A) CONFERENCE NAME 60th Annual Meeting of the American Association for the Study of Liver Diseases: The Liver Meeting 2009 CONFERENCE LOCATION Boston, MA CONFERENCE DATE 2009-10-30 to 2009-11-03.
- 8137 *Twomey PJ, Reynolds TM. The MDRD formula and validation. *QJM* 2006;99:804-5.
- 8144 *Tzamaloukas AH, Murata GH. Computational formulas for clearance indices in continuous ambulatory peritoneal dialysis. *Perit Dial Int* 1996;16:13-4.
- 8145 *Tzamaloukas AH, Murata GH. Creatinine excretion in continuous peritoneal dialysis: a systematic error of the Cockcroft-Gault formula. *Am J Kidney Dis* 2001;38:862-6.

- 8146 *Tzamaloukas AH, Murata GH. A population-specific formula predicting creatinine excretion in continuous peritoneal dialysis. *Perit Dial Int* 2002;22:67-72.
- 14068 *Ueda K, Miguel M, Wilson J, Bohannon L, Katznelson S, Lu L, et al. Comparison of estimated glomerular filtration rate (eGFR) calculations to radiolabeled measured GFR in kidney transplant recipients. *American Journal of Transplantation* 2011;11:216.
- 13843 *Uemura O, Honda M, Matsuyama T, Ishikura K, Hataya H, Yata N, et al. Age, gender, and body length effects on reference serum creatinine levels determined by an enzymatic method in Japanese children: a multicenter study. *Clin Exp Nephrol* 2011;15:694-9.
- 13814 *Uemura O, Ushijima K, Nagai T, Yamada T, Yamakawa S, Hibi Y, et al. Measurements of serum cystatin C concentrations underestimate renal dysfunction in pediatric patients with chronic kidney disease. *Clin Exp Nephrol* 2011;15:535-8.
- 8158 *Urbaniak J, Weyde W, Smolska D, Zagocka E, Klak R, Kusztal M, et al. S-cystatin C formulae or combination of s-cystatin C and s-creatinine formulae do not improve prediction of GFR. *Nephrol Dial Transplant* 2008;23:2425-6; author reply 2426-7.
- 8167 *Vachvanichsanong P, Saeteu P, Geater A. Simple estimation of the glomerular filtration rate in sick Thai children. *Nephrology (Carlton)* 2003;8:251-5.
- 8188 *Waller DG, Fleming JS, Ramsey B, Gray J. The accuracy of creatinine clearance with and without urine collection as a measure of glomerular filtration rate. *Postgrad Med J* 1991;67:42-6.
- 8189 *Waller DG, Juer L, Fleming JS. Measurement of glomerular function during cisplatin therapy. *Br J Clin Pharmacol* 1991;31:271-4.
- 8192 *Walser M. Assessment of renal function and progression of disease. *Curr Opin Nephrol Hypertens* 1994;3:564-7.
- 8193 *Walser M. Assessing renal function from creatinine measurements in adults with chronic renal failure. *Am J Kidney Dis* 1998;32:23-31.
- 917 *Walser M, Drew HH, Guldán JL. Prediction of glomerular filtration rate from serum creatinine concentration in advanced chronic renal failure. *Kidney Int* 1993;44:1145-8.
- 8209 *Van Biesen W, Vanholder R, Veys N, Verbeke F, Delanghe J, De Bacquer D, et al. The importance of standardization of creatinine in the implementation of guidelines and recommendations for CKD: implications for CKD management programmes. *Nephrol Dial Transplant* 2006;21:77-83.
- 8211 *van de Ree MA, Christiaan G, Huisman MV, van der Vijver JC, Meinders AE. Monitoring renal function in obese patients with type 2 diabetes mellitus in daily practice. *Diabetes Nutr Metab* 2001;14:66-70.
- 2378 *van den Anker JN, de Groot R, Broerse HM, Sauer PJ, van der Heijden BJ, Hop WC, et al. Assessment of glomerular filtration rate in preterm infants by serum creatinine: comparison with inulin clearance. *Pediatrics* 1995;96:1156-8.
- 13867 *van den Brand JA, van Boekel GA, Willems HL, Kiemeneij LA, den Heijer M, Wetzels JF. Introduction of the CKD-EPI equation to estimate glomerular filtration rate in a Caucasian population. *Nephrol Dial Transplant* 2011;26:3176-81.
- 8217 *Van Den Noortgate N, Velghe A, Petrovic M, Vandewiele C, Lameire N, Voet D, et al. The role of ultrasonography in the assessment of renal function in the elderly. *J Nephrol* 2003;16:658-62.
- 8219 *Van Den Noortgate NJ, Janssens WH, Delanghe JR, Afschrift MB, Lameire NH. Serum cystatin C concentration compared with other markers of glomerular filtration rate in the old old. *J Am Geriatr Soc* 2002;50:1278-82.

8223 *van Deventer HE, George JA, Paiker JE, Becker PJ, Katz IJ. Estimating glomerular filtration rate in black South Africans by use of the modification of diet in renal disease and Cockcroft-Gault equations. *Clin Chem* 2008;54:1197-202.

13610 *van Deventer HE, Paiker JE, Katz IJ, George JA. A comparison of cystatin C- and creatinine-based prediction equations for the estimation of glomerular filtration rate in black South Africans. *Nephrol Dial Transplant*. 2011;1553-1558.

8231 *Van Lente F, Suit P. Assessment of renal function by serum creatinine and creatinine clearance: glomerular filtration rate estimated by four procedures. *Clin Chem* 1989;35:2326-30.

13868 *Van Pottelbergh G, Vaes B, Morelle J, Jadoul M, Wallemacq P, Degryse J. Estimating GFR in the oldest old: does it matter what equation we use? *Age Ageing* 2011;40:401-5.

13433 *van Pottelbergh G, van Heden L, Mathei C, Degryse J. Methods to evaluate renal function in elderly patients: A systematic literature review. *Age and Ageing* 2010;39:542-548.

2385 *van Rossum LK, Mathot RA, Cransberg K, Zietse R, Vulto AG. Estimation of the glomerular filtration rate in children: which algorithm should be used? *Pediatr Nephrol* 2005;20:1769-75.

8250 *Wang JY, Lu YS, Wang SJ, Cheng CH, Shu KH, Lian JD. Comparison and correlation of measurements of glomerular filtration rates by Tc-99m DTPA and 24-hour creatinine clearance. *Zhonghua Yi Xue Za Zhi (Taipei)* 1995;55:432-7.

8252 *Wang X, Lewis J, Appel L, Cheek D, Contreras G, Faulkner M, et al. Validation of creatinine-based estimates of GFR when evaluating risk factors in longitudinal studies of kidney disease. *J Am Soc Nephrol* 2006;17:2900-9.

13869 *Wang X, Xu G, Li H, Liu Y, Wang F. Reference intervals for serum creatinine with enzymatic assay and evaluation of four equations to estimate glomerular filtration rate in a healthy Chinese adult population. *Clin Chim Acta* 2011;412:1793-7.

13434 *Ware RE, Rees RC, Sarnaik SA, Iyer RV, Alvarez OA, Casella JF, et al. Renal function in infants with sickle cell anemia: baseline data from the BABY HUG trial. In: *The Journal of pediatrics*; 2010. p 66-70.e1.

8279 *Waz WR, Quattrin T, Feld LG. Serum creatinine, height, and weight do not predict glomerular filtration rate in children with IDDM. *Diabetes Care* 1993;16:1067-70.

13825 *Weinert LS, Camargo EG, Soares AA, Silveiro SP. Review: Glomerular filtration rate estimation: performance of serum cystatin C-based prediction equations. *Clin Chem Lab Med* 2011.

14069 *Weinert LS, Camargo EG, Soares AA, Silveiro SP. Glomerular filtration rate estimation: Performance of serum cystatin C-based prediction equations. *Clinical Chemistry and Laboratory Medicine* 2011;49:1761-1771.

1654 *Verhave JC, Balje-Volkers CP, Hillege HL, de Zeeuw D, de Jong PE. The reliability of different formulae to predict creatinine clearance. *J Intern Med* 2003;253:563-73.

8314 *Verhave JC, Fesler P, Ribstein J, du Cailar G, Mimran A. Estimation of renal function in subjects with normal serum creatinine levels: influence of age and body mass index. *Am J Kidney Dis* 2005;46:233-41.

8318 *Vervoort G, Klein Gunnewiek JM, Willems HL, Wetzels JF. Effect of creatinine assay standardization on the performance of Cockcroft-Gault and MDRD formula in predicting GFR. *Nephrol Dial Transplant* 2006;21:2998-9.

1655 *Vervoort G, Willems HL, Wetzels JF. Assessment of glomerular filtration rate in healthy subjects and normoalbuminuric diabetic patients: validity of a new (MDRD) prediction equation. *Nephrol Dial Transplant* 2002;17:1909-13.

86 *Wharton WW, 3rd, Sondeen JL, McBiles M, Gradwohl SE, Wade CE, Ciceri DP, et al. Measurement of glomerular filtration rate in ICU patients using 99mTc-DTPA and inulin. *Kidney Int* 1992;42:174-8.

- 8333 *White C, Akbari A, Hussain N, Dinh L, Filler G, Lepage N, et al. Estimating glomerular filtration rate in kidney transplantation: a comparison between serum creatinine and cystatin C-based methods. *J Am Soc Nephrol* 2005;16:3763-70.
- 8334 *White C, Akbari A, Hussain N, Dinh L, Filler G, Lepage N, et al. Chronic kidney disease stage in renal transplantation classification using cystatin C and creatinine-based equations. *Nephrol Dial Transplant* 2007;22:3013-20.
- 8335 *White CA, Akbari A, Doucette S, Fergusson D, Hussain N, Dinh L, et al. A novel equation to estimate glomerular filtration rate using beta-trace protein. *Clin Chem* 2007;53:1965-8.
- 8336 *White CA, Akbari A, Doucette S, Fergusson D, Hussain N, Dinh L, et al. Estimating GFR using serum beta trace protein: accuracy and validation in kidney transplant and pediatric populations. *Kidney Int* 2009.
- 13318 *White CA, Akbari A, Doucette S, Fergusson D, Knoll GA. Estimating Glomerular Filtration Rate in Kidney Transplantation: Is the New Chronic Kidney Disease Epidemiology Collaboration Equation Any Better? *Clin Chem*. 2009.
- 13694 *White CA, Akbari A, Doucette S, Fergusson D, Knoll GA. Estimating glomerular filtration rate in kidney transplantation: is the new chronic kidney disease epidemiology collaboration equation any better? *Clin Chem* 2010;56:474-7.
- 2401 *White CA, Huang D, Akbari A, Garland J, Knoll GA. Performance of creatinine-based estimates of GFR in kidney transplant recipients: a systematic review. *Am J Kidney Dis* 2008;51:1005-15.
- 8351 *Vickery S, Stevens PE, Dalton RN, van Lente F, Lamb EJ. Does the ID-MS traceable MDRD equation work and is it suitable for use with compensated Jaffe and enzymatic creatinine assays? *Nephrol Dial Transplant* 2006;21:2439-45.
- 8358 *Wieczorowska-Tobis K, Mossakowska M, Niemir Z, Breborowicz A, Oreopoulos DG. Discrepancies in creatinine clearance in centenarians when calculated by two different mathematical formulas. *Nephrol Dial Transplant* 2002;17:2274-5.
- 8360 *Wieczorowska-Tobis K, Niemir ZI, Guzik P, Mossakowska M. Kidney function estimated with different formulas in centenarians. *Rocz Akad Med Bialymst* 2004;49:219-21.
- 8385 *Willems D, Wolff F, Mekhali F, Gillet C. Cystatin C for early detection of renal impairment in diabetes. *Clin Biochem* 2009;42:108-10.
- 2409 *Willems HL, Hilbrands LB, van de Calseyde JF, Monnens LA, Swinkels DW. Is serum cystatin C the marker of choice to predict glomerular filtration rate in paediatric patients? *Ann Clin Biochem* 2003;40:60-4.
- 14070 *Winearls CG, Lamb EJ. Chronic kidney disease: The CKD-EPI equation to estimate GFR - Better alchemy? *Nature Reviews Nephrology* 2011;7:127-128.
- 8399 *Vinge E, Lindergard B, Nilsson-Ehle P, Grubb A. Relationships among serum cystatin C, serum creatinine, lean tissue mass and glomerular filtration rate in healthy adults. *Scand J Clin Lab Invest* 1999;59:587-92.
- 8402 *Winston JA. Assessing kidney function in HIV infection. *AIDS Read* 2007;17:257-61, 264.
- 8404 *Virga G, Gaspari F, Thomaseth K, Cara M, Mastrosimone S, Rossi V. A new equation for estimating renal function using age, body weight and serum creatinine. *Nephron Clin Pract* 2007;105:c43-53.
- 8405 *Virga G, La Milia V, Russo R, Bonfante L, Cara M, Nordio M. Comparison between creatinine-based equations for estimating total creatinine clearance in peritoneal dialysis: a multicentre study. *Nephrol Dial Transplant* 2009.
- 8406 *Visvardis G, Griveas I, Zilidou R, Papadopoulou D, Mitsopoulos E, Kyriklidou P, et al. Glomerular filtration rate estimation in renal transplant patients based on serum cystatin-C levels: comparison with other markers of glomerular filtration rate. *Transplant Proc* 2004;36:1757-9.

- 209 *Work DF, Schwartz GJ. Estimating and measuring glomerular filtration rate in children. *Curr Opin Nephrol Hypertens* 2008;17:320-5.
- 8428 *Wright JG, Boddy AV, Highley M, Fenwick J, McGill A, Calvert AH. Estimation of glomerular filtration rate in cancer patients. *Br J Cancer* 2001;84:452-9.
- 13870 *Wuerzner G, Bochud M, Giusti V, Burnier M. Measurement of glomerular filtration rate in obese patients: pitfalls and potential consequences on drug therapy. *Obes Facts* 2011;4:238-43.
- 321 *Wulkan R, den Hollander J, Berghout A. Cystatin C: unsuited to use as a marker of kidney function in the intensive care unit. *Crit Care* 2005;9:531-2; author reply 531-2.
- 8438 *Wuyts B, Bernard D, Van den Noortgate N, Van de Walle J, Van Vlem B, De Smet R, et al. Reevaluation of formulas for predicting creatinine clearance in adults and children, using compensated creatinine methods. *Clin Chem* 2003;49:1011-4.
- 8440 *Xia LH, Bing XG, An XT. Serum cystatin C assay for the detection of early renal impairment in diabetic patients. *J Clin Lab Anal* 2004;18:31-5.
- 13612 *Xirouchakis E, Marelli L, Cholongitas E, Manousou P, Calvaruso V, Pleguezuelo M, et al. Comparison of cystatin C and creatinine-based glomerular filtration rate formulas with ⁵¹Cr-EDTA clearance in patients with cirrhosis. *Clin J Am Soc Nephrol*. 2011;6:84-92.
- 13435 *Xu H, Teo BW, Shuter B, Wang D, Li J, Sinha AK, et al. An ethnic Chinese coefficient for the re-expressed MDRD equation using standardized creatinine. *American Journal of Kidney Diseases* 2010;55:A112.
- 13436 *Xu H, Teo BW, Wang D, Li J, Sethi S, Lee E. Standardized serum creatinine with cystatin C improves accuracy of GFR estimates in Asians. *American Journal of Kidney Diseases* 2010;55:A112.
- 8447 *Xun L, Cheng W, Hua T, Chenggang S, Zhujiang C, Zengchun Y, et al. Assessing glomerular filtration rate (GFR) in elderly Chinese patients with chronic kidney disease (CKD): A comparison of various predictive equations. *Arch Gerontol Geriatr* 2009.
- 8459 *Yap HK, Sundram FX, Yip WC, Tay JS, Ang ES, Toh HJ, et al. Estimation of glomerular filtration rate in children using 99m-technetium DTPA. A comparison with 51-chromium EDTA clearance. *Acta Paediatr Scand* 1985;74:579-83.
- 13331 *Yeo Y, Han DJ, Moon DH, Park JS, Yang WS, Chang JW, et al. Suitability of the IDMS-Traceable MDRD Equation Method to Estimate GFR in Early Postoperative Renal Transplant Recipients. *Nephron Clin Pract*. 2010;114:108-117.
- 8467 *Ylinen EA, Ala-Houhala M, Harmoinen AP, Knip M. Cystatin C as a marker for glomerular filtration rate in pediatric patients. *Pediatr Nephrol* 1999;13:506-9.
- 14071 *Yu B, Zhou C, Bandinelli S. Combining multiple continuous tests for the diagnosis of kidney impairment in the absence of a gold standard. *Statistics in Medicine* 2011;30:1712-1721.
- 8495 *Zahran A, El-Husseini A, Shoker A. Can cystatin C replace creatinine to estimate glomerular filtration rate? A literature review. *Am J Nephrol* 2007;27:197-205.
- 2424 *Zahran A, Hossain MA, Kora MA, Galal AZ, Shoker A. Validation of the Virga GFR equation in a renal transplant population. *Nephron Clin Pract* 2008;109:c140-7.
- 2426 *Zahran A, Qureshi M, Shoker A. Comparison between creatinine and cystatin C-based GFR equations in renal transplantation. *Nephrol Dial Transplant* 2007;22:2659-68.
- 8505 *Zapletal C, Lorenz MW, Woeste G, Wullstein C, Golling M, Bechstein WO. Predicting creatinine clearance by a simple formula following live-donor kidney transplantation. *Transpl Int* 2004;17:490-4.
- 8506 *Zappitelli M, Joseph L, Gupta IR, Bell L, Paradis G. Validation of child serum creatinine-based prediction equations for glomerular filtration rate. *Pediatr Nephrol* 2007;22:272-81.

- 1691 *Zappitelli M, Parvex P, Joseph L, Paradis G, Grey V, Lau S, et al. Derivation and validation of cystatin C-based prediction equations for GFR in children. *Am J Kidney Dis* 2006;48:221-30.
- 13667 *Zappitelli M, Zhang X, Foster BJ. Estimating glomerular filtration rate in children at serial follow-up when height is unknown. *Clin J Am Soc Nephrol*. 2010;5:1763-9.
- 2428 *Zarowitz BJ, Robert S, Peterson EL. Prediction of glomerular filtration rate using aminoglycoside clearance in critically ill medical patients. *Ann Pharmacother* 1992;26:1205-10.
- 8530 *Zhao WY, Zeng L, Zhu YH, Wang LM, Zhou MS, Han S, et al. A comparison of prediction equations for estimating glomerular filtration rate in Chinese potential living kidney donors. *Clin Transplant* 2009;23:469-75.
- 8534 *Zietse R, Blankestijn PJ, Pos B, Balk AH, Derkx FH, Weimar W, et al. Optimising glomerular filtration rate and effective renal plasma flow measurements using a simple pharmacokinetic model. *Clin Nephrol* 1995;43:29-34.
- 2432 *Zitta S, Holzer H, Reibnegger G, Estelberger W. Comment on 'Comparison between creatinine and cystatin C-based GFR equations in renal transplantation'. *Nephrol Dial Transplant* 2008;23:1764; author reply 1765.
- 8543 *Zubairi AM, Hussain A. The glomerular filtration rate: comparison of various predictive equations based on serum creatinine with conventional creatinine clearance test in Pakistani population. *J Pak Med Assoc* 2008;58:182-5.
- 8545 *Zuo L, Ma YC, Zhou YH, Wang M, Xu GB, Wang HY. Application of GFR-estimating equations in Chinese patients with chronic kidney disease. *Am J Kidney Dis* 2005;45:463-72.
- 8546 *Zuo L, Ying C, Wang M, Zhang CL, Wang RF, Wang HY. Prediction of two-sample (99m)Tc-diethylene triamine pentaacetic acid plasma clearance from single-sample method. *Ann Nucl Med* 2005;19:399-405.

Kapitel 3.3

Record nr, författare, artikel mm.

- 12857 Now for the renal side of the story. Prevention and early detection of kidney disease could save your life. *Harv Health Lett.* 2005;30:4-5.
- 12859 New kidney function test a better predictor of risk. *WHO Drug Information* 2005;19:115.
- 12860 Use of cystatin C measurement in evaluating kidney function. *Nephrology* 2005;10:S157-S167.
- 13532 Direct measurement of glomerular filtration rate. *Nephrology* 2005;10:S168-S171.
- 12862 Improved GFR estimates with a new model that incorporates both cystatin C and creatinine. *Nature Clinical Practice Nephrology* 2008;4:120-121.
- 12863 Plasma cystatin C correlates with GFR in patients with anorexia nervosa. *Nature Clinical Practice Nephrology* 2009;5:118.
- 326 *Abdella N, Mojiminiyi OA, Akanji AO. Homocysteine and endogenous markers of renal function in type 2 diabetic patients without coronary heart disease. *Diabetes Res Clin Pract* 2000;50:177-85.
- 12864 *Abrahamsson K, Jodal U, Sixt R, Olsson I, Sillen U. Estimation of renal function in children and adolescents with spinal dysraphism. *J Urol.* 2008;179:2407-9.
- 12866 *Abu-Omar Y, Mussa S, Naik MJ, MacCarthy N, Standing S, Taggart DP. Evaluation of Cystatin C as a marker of renal injury following on-pump and off-pump coronary surgery. *Eur J Cardiothorac Surg.* 2005;27:893-8.
- 12867 *Adolph E, Holdt-Lehmann B, Chatterjee T, Paschka S, Prott A, Schneider H, et al. Renal Insufficiency Following Radiocontrast Exposure Trial (REINFORCE): a randomized comparison of sodium bicarbonate versus sodium chloride hydration for the prevention of contrast-induced nephropathy. *Coron Artery Dis.* 2008;19:413-9.
- 212 *Agarwal R. Ambulatory GFR measurement with cold iohalamate in adults with chronic kidney disease. *Am J Kidney Dis* 2003;41:752-9.
- 12868 *Ahlstrom A, Tallgren M, Peltonen S, Pettila V. Evolution and predictive power of serum cystatin C in acute renal failure. *Clin Nephrol.* 2004;62:344-50.
- 12869 *Aho PS, Niemi T, Lindgren L, Lepantalo M. Endovascular vs open AAA repair: similar effects on renal proximal tubular function. *Scand J Surg.* 2004;93:52-6.
- 12870 *Akbari A, Lepage N, Keely E, Clark HD, Jaffey J, MacKinnon M, et al. Cystatin-C and beta trace protein as markers of renal function in pregnancy. *BJOG.* 2005;112:575-8.
- 4313 *Akbas SH, Yavuz A, Tuncer M, Ruhi C, Gurkan A, Cetinkaya R, et al. Serum cystatin C as an index of renal function in kidney transplant patients. *Transplant Proc* 2004;36:99-101.
- 4317 *Aksun SA, Ozmen D, Ozmen B, Parildar Z, Mutaf I, Turgan N, et al. Beta2-microglobulin and cystatin C in type 2 diabetes: assessment of diabetic nephropathy. *Exp Clin Endocrinol Diabetes* 2004;112:195-200.
- 12871 *Al Wakeel JS, Hammad D, Al Suwaida A, Tarif N, Chaudhary A, Isnani A, et al. Validation of predictive equations for glomerular filtration rate in the Saudi population. *Saudi J Kidney Dis Transpl.* 2009;20:1030-7.
- 969 *Al Wakeel JS, Memon NA, Chaudhary A, Mitwalli AH, Tarif N, Isnani A, et al. Normal reference levels of serum cystatin C in Saudi adults. *Saudi J Kidney Dis Transpl* 2008;19:361-70.
- 12872 *Albayrak S, Ordu S, Ozhan H, Yazici M, Aydin M, Alemdar R, et al. Effect of olmesartan medoxomil on cystatin C level, left ventricular hypertrophy and diastolic function. *Blood Press.* 2009:1-5.

- 12873 *Alnaeb ME, Youssef F, Mikhailidis DP, Hamilton G. Short-term lipid-lowering treatment with atorvastatin improves renal function but not renal blood flow indices in patients with peripheral arterial disease. *Angiology*. 2006;57:65-71.
- 12874 *Al-Tonbary YA, Hammad AM, Zaghloul HM, El-Sayed HE, Abu-Hashem E. Pretreatment cystatin C in children with malignancy: can it predict chemotherapy-induced glomerular filtration rate reduction during the induction phase? *J Pediatr Hematol Oncol*. 2004;26:336-41.
- 12876 *Alvarez O, Zilleruelo G, Wright D, Montane B, Lopez-Mitnik G. Serum cystatin C levels in children with sickle cell disease. *Pediatr Nephrol*. 2006;21:533-7.
- 12877 *Ambrosch A, Dierkes J, Lobmann R, Kuhne W, Konig W, Luley C, et al. Relation between homocysteinaemia and diabetic neuropathy in patients with Type 2 diabetes mellitus. *Diabet Med*. 2001;18:185-92.
- 12878 *Andersen TB, Erlandsen EJ, Frokiaer J, Eskild-Jensen A, Brochner-Mortensen J. Comparison of within- and between-subject variation of serum cystatin C and serum creatinine in children aged 2-13 years. *Scand J Clin Lab Invest*. 2010.
- 4366 *Andersen TB, Eskild-Jensen A, Frokiaer J, Brochner-Mortensen J. Measuring glomerular filtration rate in children; can cystatin C replace established methods? A review. *Pediatr Nephrol* 2009;24:929-41.
- 14023 *Andersen TB, Jodal L, Boegsted M, Erlandsen EJ, Morsing A, Frokiaer J, et al. GFR Prediction From Cystatin C and Creatinine in Children: Effect of Including Body Cell Mass. *American Journal of Kidney Diseases* 2011.
- 1716 *Ansquer JC, Dalton RN, Causse E, Crimet D, Le Malicot K, Foucher C. Effect of fenofibrate on kidney function: a 6-week randomized crossover trial in healthy people. *Am J Kidney Dis* 2008;51:904-13.
- 12879 *Anwar W, Gueant JL, Abdelmoutaleb I, Adjalla C, Gerard P, Lemoel G, et al. Hyperhomocysteinemia is related to residual glomerular filtration and folate, but not to methylenetetrahydrofolate-reductase and methionine synthase polymorphisms, in supplemented end-stage renal disease patients undergoing hemodialysis. *Clin Chem Lab Med*. 2001;39:747-52.
- 12880 *Arbeiter K. Neonate or preterm infants with acute renal failure - Generally undertreated? *Wiener Klinische Wochenschrift* 2008;120:125-127.
- 12881 *Arimoto T, Takeishi Y, Niizeki T, Takabatake N, Okuyama H, Fukui A, et al. Cystatin C, a novel measure of renal function, is an independent predictor of cardiac events in patients with heart failure. *J Card Fail*. 2005;11:595-601.
- 3183 *Armanjil D, Yurdakok M, Canpolat FE, Korkmaz A, Yigit S, Tekinalp G. Determination of reference values for plasma cystatin C and comparison with creatinine in premature infants. *Pediatr Nephrol* 2008;23:2081-3.
- 4198 *Aronson JK. Drugs and Renal Insufficiency. *Medicine* 2003;31:103-109.
- 4412 *Arpegard J, Ostergren J, de Faire U, Hansson LO, Svensson P. Cystatin C--a marker of peripheral atherosclerotic disease? *Atherosclerosis* 2008;199:397-401.
- 170 *Artunc FH, Fischer IU, Risler T, Erley CM. Improved estimation of GFR by serum cystatin C in patients undergoing cardiac catheterization. *Int J Cardiol* 2005;102:173-8.
- 12882 *Asic-Buturovic B, Cavaljuga S. Cystatin C as a marker for detection of early renal failure in diabetes type 2 patients. *Bosn J Basic Med Sci*. 2005;5:68-71.
- 12883 *Asselbergs FW, Mozaffarian D, Katz R, Kestenbaum B, Fried LF, Gottdiener JS, et al. Association of renal function with cardiac calcifications in older adults: the cardiovascular health study. *Nephrol Dial Transplant*. 2009;24:834-40.
- 12884 *Astor BC, Levey AS, Stevens LA, Van Lente F, Selvin E, Coresh J. Method of glomerular filtration rate estimation affects prediction of mortality risk. *J Am Soc Nephrol*. 2009;20:2214-22.

- 12885 *Aydn F, Tezcan G, Gungor O, Cengiz AK, Hazar V, Akman S, et al. Can serum cystatin C reflect the glomerular filtration rate accurately in pediatric patients under chemotherapeutic treatment? A comparative study with Tc-99m DTPA two-plasma sample method. *Nucl Med Commun*. 2010.
- 14072 *Aygün B, Mortier NA, Smeltzer MP, Hankins JS, Ware RE. Glomerular hyperfiltration and albuminuria in children with sickle cell anemia. *Pediatric Nephrology* 2011;26:1285-1290.
- 997 *Babay Z, Al-Wakeel J, Addar M, Mittwalli A, Tarif N, Hammad D, et al. Serum cystatin C in pregnant women: reference values, reliable and superior diagnostic accuracy. *Clin Exp Obstet Gynecol* 2005;32:175-9.
- 13583 *Bacchetta J, Cochat P, Rognant N, Ranchin B, Hadj-Aissa A, Dubourg L. Which Creatinine and Cystatin C Equations Can Be Reliably Used in Children? *Clin J Am Soc Nephrol*. 2011;6:552-560.
- 12886 *Bachorzewska-Gajewska H, Malyszko J, Sitniewska E, Malyszko JS, Dobrzycki S. Neutrophil gelatinase-associated lipocalin (NGAL) correlations with cystatin C, serum creatinine and eGFR in patients with normal serum creatinine undergoing coronary angiography. *Nephrol Dial Transplant*. 2007;22:295-6.
- 12887 *Bachorzewska-Gajewska H, Malyszko J, Sitniewska E, Malyszko JS, Pawlak K, Mysliwiec M, et al. Could neutrophil-gelatinase-associated lipocalin and cystatin C predict the development of contrast-induced nephropathy after percutaneous coronary interventions in patients with stable angina and normal serum creatinine values? *Kidney Blood Press Res*. 2007;30:408-15.
- 12888 *Bachorzewska-Gajewska H, Malyszko J, Sitniewska E, Malyszko JS, Poniatowski B, Pawlak K, et al. NGAL (neutrophil gelatinase-associated lipocalin) and cystatin C: are they good predictors of contrast nephropathy after percutaneous coronary interventions in patients with stable angina and normal serum creatinine? *Int J Cardiol*. 2008;127:290-1.
- 4451 *Back SE, Lundahl T, Isaksson A. [Regarding kidney function assessment (GFR) with formulas]. *Lakartidningen* 2008;105:1780-1.
- 12889 *Bagshaw SM. Novel biomarkers for early diagnosis of acute kidney injury. *Expert Opinion on Medical Diagnostics* 2008;2:1041-1054.
- 12890 *Bagshaw SM, Bellomo R. Early diagnosis of acute kidney injury. *Current Opinion in Critical Care* 2007;13:638-644.
- 12891 *Bagshaw SM, Gibney RT. Conventional markers of kidney function. *Crit Care Med*. 2008;36:S152-8.
- 1001 *Bahar A, Yilmaz Y, Unver S, Gocmen I, Karademir F. Reference values of umbilical cord and third-day cystatin C levels for determining glomerular filtration rates in newborns. *J Int Med Res* 2003;31:231-5.
- 4464 *Bakoush O, Grubb A, Rippe B. Inaccuracy of GFR predictions by plasma cystatin C in patients without kidney dysfunction and in advanced kidney disease. *Clin Nephrol* 2008;69:331-8.
- 12892 *Balik M, Jabor A, Waldauf P, Kolar M, Pavlisova M, Brest'an D, et al. Cystatin C as a marker of residual renal function during continuous hemodiafiltration. *Kidney Blood Press Res*. 2005;28:14-9.
- 1002 *Bandaranayake N, Ankrah-Tetteh T, Wijeratne S, Swaminathan R. Intra-individual variation in creatinine and cystatin C. *Clin Chem Lab Med* 2007;45:1237-9.
- 12893 *Banfi G, Del Fabbro M, d'Eril GM, Melegati G. Reliability of cystatin C in estimating renal function in rugby players. *Ann Clin Biochem*. 2009;46:428.
- 4485 *Bardi E, Bobok I, Olah AV, Olah E, Kappelmayer J, Kiss C. Cystatin C is a suitable marker of glomerular function in children with cancer. *Pediatr Nephrol* 2004;19:1145-7.
- 13584 *Bargnoux AS, Cavalier E, Cristol JP, Simon N, Dupuy AM, Garrigue V, et al. Cystatin C is a reliable marker for estimation of glomerular filtration rate in renal transplantation: validation of a new turbidimetric assay using monospecific sheep antibodies. *Clin Chem Lab Med*. 2011;49:265-270.

- 13832 *Bargnoux AS, Perrin M, Garrigue V, Badiou S, Dupuy AM, Mourad G, et al. Analytical performances of cystatin C turbidimetric assay: which impact on accuracy of glomerular filtration rate estimation in renal transplantation? *Clin Chem Lab Med* 2011.
- 4490 *Barraclough K, Er L, Ng F, Harris M, Montaner J, Levin A. A comparison of the predictive performance of different methods of kidney function estimation in a well-characterized HIV-infected population. *Nephron Clin Pract* 2009;111:c39-48.
- 4492 *Barraclough LH, Field C, Wieringa G, Swindell R, Livsey JE, Davidson SE. Estimation of renal function -- what is appropriate in cancer patients? *Clin Oncol (R Coll Radiol)* 2008;20:721-6.
- 12896 *Baxmann AC, Ahmed MS, Marques NC, Menon VB, Pereira AB, Kirsztajn GM, et al. Influence of muscle mass and physical activity on serum and urinary creatinine and serum cystatin C. *Clin J Am Soc Nephrol*. 2008;3:348-54.
- 4529 *Beauvieux MC, Le Moigne F, Lasseur C, Raffaitin C, Perlemoine C, Barthe N, et al. New predictive equations improve monitoring of kidney function in patients with diabetes. *Diabetes Care* 2007;30:1988-94.
- 4531 *Beck M, Graf C, Ellenrieder B, Bokenkamp A, Huber A, Hecher K, et al. Long-term outcome of kidney function after twin-twin transfusion syndrome treated by intrauterine laser coagulation. *Pediatr Nephrol* 2005;20:1657-9.
- 13603 *Becker BN, Vassalotti JA. A Software Upgrade: CKD Testing in 2010. *American Journal of Kidney Diseases* 2010;55:8-10.
- 12897 *Bell M, Granath F, Martensson J, Lofberg E, Ekblom A, Martling CR. Cystatin C is correlated with mortality in patients with and without acute kidney injury. *Nephrol Dial Transplant*. 2009;24:3096-102.
- 12899 *Ben-Dov IZ. Why choose serum cystatin C levels over serum creatinine levels as a serologic marker of kidney function? *Ann Intern Med*. 2008;149:284; author reply 284.
- 1750 *Benohr P, Grenz A, Hartmann JT, Muller GA, Blaschke S. Cystatin C--a marker for assessment of the glomerular filtration rate in patients with cisplatin chemotherapy. *Kidney Blood Press Res* 2006;29:32-5.
- 13585 *Berding G, Geisler S, Melter M, Marquardt P, Luhr A, Scheller F, et al. Estimation of glomerular filtration rate in liver-transplanted children: comparison of simplified procedures using ⁵¹Cr-EDTA and endogenous markers with Sapirstein's method as a reference standard. *Pediatr Transplant*. 2010;14:786-95.
- 12900 *Berenguer-Pina JJ, Noguera J, Canizares F, Hernandez-Bernal C, Velasco-Conesa E, Lozano-Postigo C, et al. Cystatin C: a marker of glomerular filtration rate in liver transplantation. *Transplant Proc*. 2002;34:268-9.
- 1754 *Berg UB, Ericzon BG, Nemeth A. Renal function before and long after liver transplantation in children. *Transplantation* 2001;72:631-7.
- 12901 *Berghout A, Wulkan RW, Den Hollander JG, Risch L, Drexel H, Huber AR, et al. Cystatin C and the risk of death [1] (multiple letters). *New England Journal of Medicine* 2005;353:842-844.
- 4578 *Beringer PM, Hidayat L, Heed A, Zheng L, Owens H, Benitez D, et al. GFR estimates using cystatin C are superior to serum creatinine in adult patients with cystic fibrosis. *J Cyst Fibros* 2009;8:19-25.
- 13586 *Beringer PM, Owens H, Nguyen A, Mordwinkin N, Louie S, Mak M, et al. Estimation of glomerular filtration rate by using serum cystatin C and serum creatinine concentrations in patients with human immunodeficiency virus. *Pharmacotherapy*. 2010;30:1004-10.
- 13546 *Berman N, Hostetter TH. Comparing the Cockcroft-Gault and MDRD equations for calculation of GFR and drug doses in the elderly: Commentary. *Nature Clinical Practice Nephrology* 2007;3:644-645.
- 14027 *Bevc S, Hojs R, Ekart R. Creatinine-based formula (MDRD) versus cystatin c-based formula (simple cystatin C formula) for estimation of GFR in patients with chronic kidney disease. *European Journal of Internal Medicine* 2011;22:S12.

- 13844 *Bevc S, Hojs R, Ekart R, Gorenjak M, Puklavec L. Simple cystatin C formula compared to sophisticated CKD-EPI formulas for estimation of glomerular filtration rate in the elderly. *Ther Apher Dial* 2011;15:261-8.
- 13845 *Bhavsar NA, Appel LJ, Kusek JW, Contreras G, Bakris G, Coresh J, et al. Comparison of Measured GFR, Serum Creatinine, Cystatin C, and Beta-Trace Protein to Predict ESRD in African Americans With Hypertensive CKD. *Am J Kidney Dis* 2011.
- 4601 *Biancofiore G, Pucci L, Cerutti E, Penno G, Pardini E, Esposito M, et al. Cystatin C as a marker of renal function immediately after liver transplantation. *Liver Transpl* 2006;12:285-91.
- 12902 *Bibbins-Domingo K, Chertow GM, Fried LF, Odden MC, Newman AB, Kritchevsky SB, et al. Renal function and heart failure risk in older black and white individuals: the Health, Aging, and Body Composition Study. *Arch Intern Med*. 2006;166:1396-402.
- 12903 *Bicik Z, Bahcebasi T, Kulaksizoglu S, Yavuz O. The efficacy of cystatin C assay in the prediction of glomerular filtration rate. Is it a more reliable marker for renal failure? *Clin Chem Lab Med*. 2005;43:855-61.
- 4 *Biglarnia AR, Wadstrom J, Larsson A. Decentralized glomerular filtration rate (GFR) estimates in healthy kidney donors show poor correlation and demonstrate the need for improvement in quality and standardization of GFR measurements in Sweden. *Scand J Clin Lab Invest* 2007;67:227-35.
- 12904 *Bittl JA. Damage control for renal artery stenting. *Circulation* 2008;117:2724-2726.
- 13847 *Blufpand HN, Tromp J, Abbink FC, Stoffel-Wagner B, Bouman AA, Schouten-van Meeteren AY, et al. Cystatin C more accurately detects mildly impaired renal function than creatinine in children receiving treatment for malignancy. *Pediatr Blood Cancer* 2011;57:262-7.
- 12905 *Bokenkamp A, De Jong PE, Verhave JC, Curhan GC. Kidney function itself, and not cystatin C, is correlated with height and weight [3] (multiple letters). *Kidney International* 2005;67:777-779.
- 3230 *Bokenkamp A, Dieterich C, Dressler F, Muhlhaus K, Gembruch U, Bald R, et al. Fetal serum concentrations of cystatin C and beta2-microglobulin as predictors of postnatal kidney function. *Am J Obstet Gynecol* 2001;185:468-75.
- 3231 *Bokenkamp A, Domanetzki M, Zinck R, Schumann G, Brodehl J. Reference values for cystatin C serum concentrations in children. *Pediatr Nephrol* 1998;12:125-9.
- 1768 *Bokenkamp A, Domanetzki M, Zinck R, Schumann G, Byrd D, Brodehl J. Cystatin C--a new marker of glomerular filtration rate in children independent of age and height. *Pediatrics* 1998;101:875-81.
- 12908 *Bokenkamp A, Ozden N, Dieterich C, Schumann G, Ehrich JH, Brodehl J. Cystatin C and creatinine after successful kidney transplantation in children. *Clin Nephrol*. 1999;52:371-6.
- 12911 *Boldt J, Wolf M. Identification of renal injury in cardiac surgery: the role of kidney-specific proteins. *J Cardiothorac Vasc Anesth*. 2008;22:122-32.
- 13547 *Bolke E, Gripp S, Matuschek C, Hermsen D, Peiper M, Budach W. Use of cystatin C as a marker to detect renal insufficiency in head and neck cancer patients with cisplatin-based chemotherapy. *Journal of Clinical Oncology* 2009;27:e17048.
- 13872 *Bolke E, Schieren G, Gripp S, Steinbach G, Peiper M, Orth K, et al. Cystatin C - a fast and reliable biomarker for glomerular filtration rate in head and neck cancer patients. *Strahlenther Onkol* 2011;187:191-201.
- 12913 *Borges RL, Hirota AH, Quinto BM, Ribeiro AB, Zanella MT, Batista MC. Is Cystatin C a Useful Marker in the Detection of Diabetic Kidney Disease? *Nephron Clin Pract*. 2009;114:c127-c134.
- 12914 *Borrows R, Cockwell P. Measuring renal function in solid organ transplant recipients. *Transplantation* 2007;83:529-531.
- 1040 *Bosio M, Bianchi ML, Lloyd SE, Thakker RV. A familial syndrome due to Arg648Stop mutation in the X-linked renal chloride channel gene. *Pediatr Nephrol* 1999;13:278-83.

- 3240 *Bostom AG, Bausserman L, Jacques PF, Liaugaudas G, Selhub J, Rosenberg IH. Cystatin C as a determinant of fasting plasma total homocysteine levels in coronary artery disease patients with normal serum creatinine. *Arterioscler Thromb Vasc Biol* 1999;19:2241-4.
- 12915 *Bostom AG, Dworkin LD. Cystatin C measurement: improved detection of mild decrements in glomerular filtration rate versus creatinine-based estimates? *Am J Kidney Dis*. 2000;36:205-7.
- 12916 *Bostom AG, Gohh RY, Bausserman L, Hakas D, Jacques PF, Selhub J, et al. Serum cystatin C as a determinant of fasting total homocysteine levels in renal transplant recipients with a normal serum creatinine. *J Am Soc Nephrol*. 1999;10:164-6.
- 4674 *Boudville N, Salama M, Jeffrey GP, Ferrari P. The inaccuracy of cystatin C and creatinine-based equations in predicting GFR in orthotopic liver transplant recipients. *Nephrol Dial Transplant* 2009;24:2926-30.
- 4680 *Bouvet Y, Bouissou F, Coulais Y, Seronie-Vivien S, Tafani M, Decramer S, et al. GFR is better estimated by considering both serum cystatin C and creatinine levels. *Pediatr Nephrol* 2006;21:1299-306.
- 1042 *Brandis M, Krause PH, Hunermund G, Bahlmann J. [Rapidly progressive glomerulonephritis in a child. Clinical and histological observation over 3 1/2 years (author's transl)]. *Monatsschr Kinderheilkd* 1976;124:85-9.
- 13873 *Brinkert F, Kemper MJ, Briem-Richter A, van Husen M, Treszl A, Ganschow R. High prevalence of renal dysfunction in children after liver transplantation: non-invasive diagnosis using a cystatin C-based equation. *Nephrol Dial Transplant* 2011;26:1407-12.
- 13587 *Bronden B, Eyjolfsson A, Blomquist S, Dardashti A, Ederoth P, Bjursten H. Evaluation of cystatin C with iohexol clearance in cardiac surgery. *Acta Anaesthesiol Scand*. 2011;55:196-202.
- 13588 *Brown S. Re: estimation of glomerular filtration rate with creatinine-based versus cystatin C-based equations in kidney transplant recipients. *Iran J Kidney Dis*. 2010;4:169-70.
- 12919 *Bulla M. Symposium on problems in renal insufficiency and replacement therapy in children: Munster, October 4th - 5th, 2000. *Nieren- und Hochdruckkrankheiten* 2000;29:459-468.
- 1791 *Burkhardt H, Bojarsky G, Gladisch R. Diagnostic efficiency of cystatin C and serum creatinine as markers of reduced glomerular filtration rate in the elderly. *Clin Chem Lab Med* 2002;40:1135-8.
- 1793 *Burkhardt H, Bojarsky G, Gretz N, Gladisch R. Creatinine clearance, Cockcroft-Gault formula and cystatin C: estimators of true glomerular filtration rate in the elderly? *Gerontology* 2002;48:140-6.
- 401 *Buysschaert M, Joudi I, Wallemacq P, Hermans MP. Comparative performance of serum cystatin-c versus serum creatinine in diabetic subjects. *Diabetes Metab* 2003;29:377-83.
- 12920 *Buysschaert M, Joudi I, Wallemacq P, Hermans MP. Performance of serum cystatin-C versus serum creatinine in subjects with type 1 diabetes. *Diabetes Care*. 2003;26:1320.
- 12907 *Bökenkamp A, Laarman CARC, Braam KI, Van Wijk JAE, Kors WA, Kool M, et al. Effect of corticosteroid therapy on low-molecular-weight protein markers of kidney function [5]. *Clinical Chemistry* 2007;53:2219-2221.
- 12909 *Bökenkamp A, Van Wijk JAE, Lentze MJ, Stoffel-Wagner B. Effect of corticosteroid therapy on serum cystatin C and (beta)(2)-microglobulin concentrations. *Clinical Chemistry* 2002;48:1123-1126.
- 12921 *Cabarkapa V, Stosic Z, Deric M, Vucurevic-Ristic L, Zeravica R, Ilincic B. Serum cystatin C in estimating glomerular filtration rate. *Journal of Medical Biochemistry* 2008;27:46-51.
- 12923 *Campo A, Lanfranco G, Gramaglia L, Goia F, Cottino R, Giusto V. Could plasma cystatin C be useful as a marker of hemodialysis low molecular weight proteins removal? *Nephron Clin Pract*. 2004;98:c79-82.
- 13548 *Cantarovich M, Yoshida EM, Peltekian KM, Marotta PJ, Greig PD, Kneteman NM, et al. Poor prediction of the glomerular filtration rate using current formulas in de novo liver transplant patients. *Transplantation* 2006;82:433-436.

13848 *Carter JL, Lane CE, Fan SL, Lamb EJ. Estimation of residual glomerular filtration rate in peritoneal dialysis patients using cystatin C: comparison with ⁵¹Cr-EDTA clearance. *Nephrol Dial Transplant* 2011;26:3729-3732.

4828 *Carter JL, O'Riordan SE, Eaglestone GL, Delaney MP, Lamb EJ. Chronic kidney disease prevalence in a UK residential care home population. *Nephrol Dial Transplant* 2008;23:1257-64.

3265 *Cataldi L, Mussap M, Bertelli L, Ruzzante N, Fanos V, Plebani M. Cystatin C in healthy women at term pregnancy and in their infant newborns: relationship between maternal and neonatal serum levels and reference values. *Am J Perinatol* 1999;16:287-95.

4841 *Cepeda FJ, Fernandez E, Pobes A, Banos LM. [Utility of cystatin-C in hospitalized patients. Comparing with different methods of assessing renal function]. *Nefrologia* 2007;27:168-74.

13589 *Cha RH, Lee CS, Lim YH, Kim H, Lee SH, Yu KS, et al. Clinical usefulness of serum cystatin C and the pertinent estimation of glomerular filtration rate based on cystatin C. *Nephrology (Carlton)*. 2010;15:768-76.

12925 *Chakrabarti S, Chattopadhyay I. Measuring renal function in old age. *Reviews in Clinical Gerontology* 2008;18:257-267.

1807 *Chantrel F, Agin A, Offner M, Koehl C, Moulin B, Hannedouche T. Comparison of cystatin C versus creatinine for detection of mild renal failure. *Clin Nephrol* 2000;54:374-81.

13549 *Chatelut E, Schmitt A, Lansiaux A, Bobin-Dubigeon C, Etienne-Grimaldi M, Boisdrion-Celle M, et al. A universal formula based on cystatin C to perform individual dosing of carboplatin in normal weight, underweight, and obese patients. *Journal of Clinical Oncology* 2009;27:2527.

4891 *Chew JS, Saleem M, Florkowski CM, George PM. Cystatin C-a paradigm of evidence based laboratory medicine. *Clin Biochem Rev* 2008;29:47-62.

4892 *Chew JS, Saleem M, Florkowski CM, George PM. Estimating renal function in oncology patients using cystatin C-based equations. *Clin Oncol (R Coll Radiol)* 2009;21:425-6.

12929 *Choe JY, Park SH, Kim SK. Serum cystatin C is a potential endogenous marker for the estimation of renal function in male gout patients with renal impairment. *J Korean Med Sci*. 2010;25:42-8.

13566 *Choi A, Scherzer R, Bacchetti P, Tien PC, Saag MS, Gibert CL, et al. Cystatin C, albuminuria, and 5-year all-cause mortality in HIV-infected persons. *Am J Kidney Dis* 2010;56:872-82.

4911 *Cholongitas E, Shusang V, Marelli L, Nair D, Thomas M, Patch D, et al. Review article: renal function assessment in cirrhosis - difficulties and alternative measurements. *Aliment Pharmacol Ther* 2007;26:969-78.

13590 *Cholongitas E, Xirouchakis E, Garcovich M, Burroughs AK. Evaluation of renal function in patients with cirrhosis. *J Hepatol*. 2010;53:589.

4918 *Christensson A, Ekberg J, Grubb A, Ekberg H, Lindstrom V, Lilja H. Serum cystatin C is a more sensitive and more accurate marker of glomerular filtration rate than enzymatic measurements of creatinine in renal transplantation. *Nephron Physiol* 2003;94:p19-27.

13591 *Christensson A, Elmståhl S. Estimation of the age-dependent decline of glomerular filtration rate from formulas based on creatinine and cystatin C in the general elderly population. *Nephron Clin Pract*. 2011;117:c40-50.

4919 *Christensson AG, Grubb AO, Nilsson JA, Norrgren K, Sterner G, Sundkvist G. Serum cystatin C advantageous compared with serum creatinine in the detection of mild but not severe diabetic nephropathy. *J Intern Med* 2004;256:510-8.

13550 *Chudleigh RA, Dunseath G, Evans W, Harvey JN, Evans P, Ollerton R, et al. How reliable is estimation of glomerular filtration rate at diagnosis of type 2 diabetes? *Diabetes care* 2007;30:300-305.

- 4923 *Chudleigh RA, Ollerton RL, Dunseath G, Peter R, Harvey JN, Luzio S, et al. Performance of the revised '175' Modification of Diet in Renal Disease equation in patients with type 2 diabetes. *Diabetologia* 2008;51:1714-8.
- 4924 *Chudleigh RA, Ollerton RL, Dunseath G, Peter R, Harvey JN, Luzio S, et al. Use of cystatin C-based estimations of glomerular filtration rate in patients with type 2 diabetes. *Diabetologia* 2009;52:1274-8.
- 13592 *Chung MY, Jun DW, Sung SA. Diagnostic value of cystatin C for predicting acute kidney injury in patients with liver cirrhosis. *Korean J Hepatol.* 2010;16:301-7.
- 12930 *Cimerman N, Brguljan PM, Krasovec M, Suskovic S, Kos J. Serum cystatin C, a potent inhibitor of cysteine proteinases, is elevated in asthmatic patients. *Clin Chim Acta.* 2000;300:83-95.
- 12931 *Cimerman N, Brguljan PM, Krasovec M, Suskovic S, Kos J. Twenty-four hour variations of cystatin C and total cysteine proteinase inhibitory activity in sera from healthy subjects. *Clinica Chimica Acta* 2000;291:89-95.
- 12932 *Claus T, Elitok S, Schmitt R, Luft FC, Kettritz R. Thyroid function and glomerular filtration--a potential for Grave errors. *Nephrol Dial Transplant.* 2005;20:1002-3.
- 12933 *Codognotto M, Piccoli A, Zaninotto M, Mion M, Vertolli U, Tona F, et al. Cystatin C in heart failure is nothing more than a bystander of glomerular filtration. *J Nephrol.* 2007;20:219-27.
- 1081 *Coll E, Botey A, Alvarez L, Poch E, Quinto L, Saurina A, et al. Serum cystatin C as a new marker for noninvasive estimation of glomerular filtration rate and as a marker for early renal impairment. *Am J Kidney Dis* 2000;36:29-34.
- 4977 *Cordeiro VF, Pinheiro DC, Silva GB, Jr., Lima JW, Mota RM, Liborio AB, et al. Comparative study of cystatin C and serum creatinine in the estimative of glomerular filtration rate in children. *Clin Chim Acta* 2008;391:46-50.
- 12935 *Coresh J, Astor B. Decreased kidney function in the elderly: clinical and preclinical, neither benign. *Ann Intern Med.* 2006;145:299-301.
- 4988 *Corrao AM, Lisi G, Di Pasqua G, Guizzardi M, Marino N, Ballone E, et al. Serum cystatin C as a reliable marker of changes in glomerular filtration rate in children with urinary tract malformations. *J Urol* 2006;175:303-9.
- 13393 *Corsonello A, Pedone C, Lattanzio F, Cherubini A, Onder G, Corica F, et al. Chronic kidney disease and 1-year survival in elderly patients discharged from acute care hospitals: a comparison of three glomerular filtration rate equations. *Nephrol Dial Transplant* 2010.
- 14073 *Corsonello A, Pedone C, Lattanzio F, Onder G, Incalzi RA. Association between Glomerular Filtration Rate and Adverse Drug Reactions in Elderly Hospitalized Patients. *Drugs & Aging* 2011;28:379-390.
- 1090 *Croda-Todd M, Juarez E, Hernandez P, Flores G, Rivera G, Bocanegra-Garcia V. Reference intervals for serum cystatin C in healthy Mexican adults. *Clin Chem Lab Med* 2007;45:925-7.
- 12936 *Curhan G. Cystatin C: A marker of renal function or something more? *Clinical Chemistry* 2005;51:293-294.
- 1095 *Dajak M, Ignjatovic S, Jovicic S, Majkic-Singh N. The values of estimated glomerular filtration rate calculated with creatinine and cystatin C based equations in healthy adults. *Clin Lab* 2008;54:153-9.
- 14034 *Damman K, van der Harst P, Smilde TDJ, Voors AA, Navis G, van Veldhuisen DJ, et al. Use of cystatin C levels in estimating renal function and prognosis in patients with chronic systolic heart failure. *Heart* 2011.
- 1833 *Daniel JP, Chantrel F, Offner M, Moulin B, Hannedouche T. Comparison of cystatin C, creatinine and creatinine clearance vs. GFR for detection of renal failure in renal transplant patients. *Ren Fail* 2004;26:253-7.
- 12937 *Davey P, Peaston R, Rose JD, Jackson RA, Wyatt MG. Impact on renal function after endovascular aneurysm repair with uncovered supra-renal fixation assessed by serum cystatin C. *Eur J Vasc Endovasc Surg.* 2008;35:439-45.

- 12939 *de Boer IH, Katz R, Cao JJ, Fried LF, Kestenbaum B, Mukamal K, et al. Cystatin C, albuminuria, and mortality among older adults with diabetes. *Diabetes Care*. 2009;32:1833-8.
- 12940 *de Boer IH, Katz R, Fried LF, Ix JH, Luchsinger J, Sarnak MJ, et al. Obesity and change in estimated GFR among older adults. *Am J Kidney Dis*. 2009;54:1043-51.
- 12941 *de Burbure C, Buchet JP, Leroyer A, Nisse C, Haguenoer JM, Mutti A, et al. Renal and neurologic effects of cadmium, lead, mercury, and arsenic in children: Evidence of early effects and multiple interactions at environmental exposure levels. *Environmental Health Perspectives* 2006;114:584-590.
- 13593 *De Francisco ALM, Fernandez E. The importance of glomerular filtration rate (GFR) estimating equations in the care of hospitalized patients. *European Journal of Internal Medicine* 2011;22:120-121.
- 5061 *de Jong PE, Gansevoort RT. Screening techniques for detecting chronic kidney disease. *Curr Opin Nephrol Hypertens* 2005;14:567-72.
- 13594 *de Jong PE, Gansevoort RT. Is cystatin C the answer to detecting progression in CKD? *J Am Soc Nephrol*. 2011;22:9-11.
- 12942 *Deinum J, Derkx FH. Cystatin for estimation of glomerular filtration rate? *Lancet*. 2000;356:1624-5.
- 1112 *Delanaye P, Cavalier E, Depas G, Chapelle JP, Krzesinski JM. New data on the intraindividual variation of cystatin C. *Nephron Clin Pract* 2008;108:c246-8.
- 5098 *Delanaye P, Cavalier E, Krzesinski JM, Mariat C. Cystatin C-based equations: don't repeat the same errors with analytical considerations. *Nephrol Dial Transplant* 2008;23:1065; author reply 1065-6.
- 5101 *Delanaye P, Cavalier E, Radermecker RP, Paquot N, Depas G, Chapelle JP, et al. Cystatin C or creatinine for detection of stage 3 chronic kidney disease in anorexia nervosa. *Nephron Clin Pract* 2008;110:c158-63.
- 5102 *Delanaye P, Cavalier E, Radermecker RP, Paquot N, Depas G, Chapelle JP, et al. Estimation of GFR by different creatinine- and cystatin-C-based equations in anorexia nervosa. *Clin Nephrol* 2009;71:482-91.
- 5103 *Delanaye P, Cavalier E, Saint-Remy A, Lutteri L, Krzesinski JM. Discrepancies between creatinine-based and cystatin C-based equations in estimating prevalence of stage 3 chronic kidney disease in an elderly population. *Scand J Clin Lab Invest* 2009;69:344-9.
- 5108 *Delanaye P, Lambermont B, Chapelle JP, Gielen J, Gerard P, Rorive G. Plasmatic cystatin C for the estimation of glomerular filtration rate in intensive care units. *Intensive Care Med* 2004;30:980-3.
- 5109 *Delanaye P, Nellessen E, Cavalier E, Depas G, Grosch S, Defraigne JO, et al. Is cystatin C useful for the detection and the estimation of low glomerular filtration rate in heart transplant patients? *Transplantation* 2007;83:641-4.
- 5111 *Delanaye P, Pieroni L, Abshoff C, Lutteri L, Chapelle JP, Krzesinski JM, et al. Analytical study of three cystatin C assays and their impact on cystatin C-based GFR-prediction equations. *Clin Chim Acta* 2008;398:118-24.
- 12943 *Delaney MP, Stevens PE, Al Hasani M, Stowe HJ, Judge C, Lamb EJ. Relationship of serum cystatin C to peritoneal and renal clearance measures in peritoneal dialysis: a cross-sectional study. *Am J Kidney Dis*. 2008;51:278-84.
- 14074 *Delanghe J. Beyond creatinine standardisation. *Clinical Chemistry and Laboratory Medicine* 2011;49:S50.
- 1116 *Delanghe JR. How to establish glomerular filtration rate in children. *Scand J Clin Lab Invest Suppl* 2008;241:46-51.
- 2 *Delanghe JR. How to estimate GFR in children. *Nephrol Dial Transplant* 2009;24:714-6.
- 12944 *Demirtas S, Akan O, Can M, Elmali E, Akan H. Cystatin C can be affected by nonrenal factors: a preliminary study on leukemia. *Clin Biochem*. 2006;39:115-8.

- 5117 *Demirtas S, Bozbas A, Akbay A, Yavuz Y, Karaca L. Diagnostic value of serum cystatin C for evaluation of hepatorenal syndrome. *Clin Chim Acta* 2001;311:81-9.
- 12945 *Den Hollander JG, Wulkan RW, Mantel MJ, Berghout A. Is cystatin C a marker of glomerular filtration rate in thyroid dysfunction? [6]. *Clinical Chemistry* 2003;49:1558-1559.
- 12946 *Dennen P, Parikh CR. Biomarkers of acute kidney injury: Can we replace serum creatinine? *Clinical Nephrology* 2007;68:269-278.
- 13595 *Deo R, Sotoodehnia N, Katz R, Sarnak MJ, Fried LF, Chonchol M, et al. Cystatin C and sudden cardiac death risk in the elderly. *Circ Cardiovasc Qual Outcomes*. 2010;3:159-64.
- 12948 *Dharnidharka VR, Kwon C, Stevens G. Serum cystatin C is superior to serum creatinine as a marker of kidney function: a meta-analysis. *Am J Kidney Dis*. 2002;40:221-6.
- 13596 *Donadio C. Serum and urinary markers of early impairment of GFR in chronic kidney disease patients: diagnostic accuracy of urinary beta-trace protein. *Am J Physiol Renal Physiol*. 2010;299:F1407-23.
- 13597 *Donadio C, Abdelkawy H, Grassi G. Echographic renal dimensions can predict glomerular filtration rate of potential living kidney donors. *Transplant Proc*. 2010;42:1035-9.
- 12949 *Donadio C, Lucchesi A, Ardini M, Giordani R. Cystatin C, beta 2-microglobulin, and retinol-binding protein as indicators of glomerular filtration rate: comparison with plasma creatinine. *J Pharm Biomed Anal*. 2001;24:835-42.
- 12950 *Donadio C, Lucchesi A, Coll E, Botey A, Alvarez L, Piera C, et al. Serum cystatin as a marker of glomerular filtration rate [1] (multiple letters). *American Journal of Kidney Diseases* 2001;37:448-451.
- 12951 *Donahue RP, Stranges S, Rejman K, Rafalson LB, Dmochowski J, Trevisan M. Elevated cystatin C concentration and progression to pre-diabetes: the Western New York study. *Diabetes Care*. 2007;30:1724-9.
- 12954 *Dworkin LD. Serum cystatin C as a marker of glomerular filtration rate. *Current Opinion in Nephrology and Hypertension* 2001;10:551-553.
- 13598 *Eaglestone G, Ellis P. Cystatin C as a marker of renal function in older adults. *Nursing Standard* 2009;24:35-9.
- 12955 *Eggers KM, Dellborg M, Oldgren J, Swahn E, Venge P, Lindahl B. Risk prediction in chest pain patients by biochemical markers including estimates of renal function. *Int J Cardiol*. 2008;128:207-13.
- 12957 *El-Shafey EM, El-Nagar GF, Selim MF, El-Sorogy HA, Sabry AA. Is serum cystatin C an accurate endogenous marker of glomerular filtration rate for detection of early renal impairment in patients with type 2 diabetes mellitus? *Ren Fail*. 2009;31:355-9.
- 12958 *ElShafie AI, Elghazali G, Ronnelid J, Venge P. Cystatin C as a marker of immune complex-associated renal impairment in a Sudanese population with visceral leishmaniasis. *Am J Trop Med Hyg*. 2006;75:864-8.
- 1872 *Emara M, Zahran A, Abd El Hady H, Shoker A. How to best define patients with moderate chronic kidney disease. *Nephron Clin Pract* 2008;110:c195-206.
- 12959 *Epstein M, Leary E, Laskey R, Zuckerman A, Reed D, Lowe W. Atorvastatin does not induce glomerular or tubular dysfunction even at high doses. *J Cardiometab Syndr*. 2007;2:163-7.
- 13599 *Eriksen BO, Mathisen UD, Melsom T, Ingebretsen OC, Jenssen TG, Njolstad I, et al. Cystatin C is not a better estimator of GFR than plasma creatinine in the general population. *Kidney Int*. 2010;78:1305-11.
- 13851 *Eriksen BO, Mathisen UD, Melsom T, Ingebretsen OC, Jenssen TG, Njolstad I, et al. The Role of Cystatin C in Improving GFR Estimation in the General Population. *Am J Kidney Dis* 2011.
- 13852 *Fabbian F, Pala M, Scanelli G, Manzato E, Longhini C, Portaluppi F. Estimation of renal function in patients with eating disorders. *Int J Eat Disord* 2011;44:233-7.

5295 *Fabre EE, Raynaud-Simon A, Golmard JL, Gourgouillon N, Beaudoux JL, Nivet-Antoine V. Interest and limits of glomerular filtration rate (GFR) estimation with formulae using creatinine or cystatin C in the malnourished elderly population. *Arch Gerontol Geriatr* 2009.

12961 *Fadowski JJ, Navas-Acien A, Tellez-Plaza M, Guallar E, Weaver VM, Furth SL. Blood lead level and kidney function in US adolescents: The Third National Health and Nutrition Examination Survey. *Arch Intern Med*. 2010;170:75-82.

13853 *Fadowski JJ, Neu AM, Schwartz GJ, Furth SL. Pediatric GFR estimating equations applied to adolescents in the general population. *Clin J Am Soc Nephrol* 2011;6:1427-35.

3354 *Fanos V, Mussap M, Plebani M, Cataldi L. Cystatin C in paediatric nephrology. Present situation and prospects. *Minerva Pediatr* 1999;51:167-77.

5325 *Fehrman-Ekholm I, Seeberger A, Bjork J, Sterner G. Serum cystatin C: a useful marker of kidney function in very old people. *Scand J Clin Lab Invest* 2009;69:606-11.

12962 *Felicio ML, Andrade RR, Castiglia YM, Silva MA, Vianna PT, Martins AS. Cystatin C and glomerular filtration rate in the cardiac surgery with cardiopulmonary bypass. *Rev Bras Cir Cardiovasc*. 2009;24:305-11.

5331 *Feriozzi S, Germain DP, Di Vito R, Legrand A, Ricci R, Barbey F. Cystatin C as a marker of early changes of renal function in Fabry nephropathy. *J Nephrol* 2007;20:437-43.

13600 *Fesler P, Mimran A. Estimation of Glomerular Filtration Rate: What Are the Pitfalls? *Curr Hypertens Rep*. 2011.

5342 *Filler G, Foster J, Acker A, Lepage N, Akbari A, Ehrich JH. The Cockcroft-Gault formula should not be used in children. *Kidney Int* 2005;67:2321-4.

13601 *Filler G, Gharib M, Casier S, Lodige P, Ehrich JH, Dave S. Prevention of chronic kidney disease in spina bifida. *Int Urol Nephrol*. 2011.

5343 *Filler G, Lepage N. Should the Schwartz formula for estimation of GFR be replaced by cystatin C formula? *Pediatr Nephrol* 2003;18:981-5.

1168 *Filler G, Priem F, Lepage N, Sinha P, Vollmer I, Clark H, et al. Beta-trace protein, cystatin C, beta(2)-microglobulin, and creatinine compared for detecting impaired glomerular filtration rates in children. *Clin Chem* 2002;48:729-36.

1169 *Filler G, Priem F, Vollmer I, Gellermann J, Jung K. Diagnostic sensitivity of serum cystatin for impaired glomerular filtration rate. *Pediatr Nephrol* 1999;13:501-5.

5357 *Finney H, Bates CJ, Price CP. Plasma cystatin C determinations in a healthy elderly population. *Arch Gerontol Geriatr* 1999;29:75-94.

1172 *Finney H, Newman DJ, Price CP. Adult reference ranges for serum cystatin C, creatinine and predicted creatinine clearance. *Ann Clin Biochem* 2000;37 (Pt 1):49-59.

12967 *Finney H, Williams AH, Price CP. Serum cystatin C in patients with myeloma. *Clin Chim Acta*. 2001;309:1-6.

1899 *Fliser D, Ritz E. Serum cystatin C concentration as a marker of renal dysfunction in the elderly. *Am J Kidney Dis* 2001;37:79-83.

13602 *Foley RN, Wang C, Collins AJ. Cystatin C, mortality risk and clinical triage in US adults: threshold values and hierarchical importance. *Nephrol Dial Transplant*. 2010.

12970 *Foley RN, Wang C, Snyder JJ, Collins AJ. Cystatin C Levels in U.S. Adults, 1988-1994 Versus 1999-2002: NHANES. *Clin J Am Soc Nephrol*. 2009.

5390 *Fontserè N, Esteve V, Saurina A, Pou M, Barba N, Deulofeu R, et al. The search for a new marker of renal function in older patients with chronic kidney disease stages 3-4: usefulness of cystatin C-based equations. *Nephron Clin Pract* 2009;112:c164-70.

12971 *Forsblom C, Hiukka A, Leinonen ES, Sundvall J, Groop PH, Taskinen MR. Effects of Long-Term Fenofibrate Treatment on Markers of Renal Function in Type 2 Diabetes: FIELD Helsinki Substudy. *Diabetes Care*. 2009.

5396 *Foster MC, Keyes MJ, Larson MG, Vita JA, Mitchell GF, Meigs JB, et al. Relations of measures of endothelial function and kidney disease: the Framingham Heart Study. *Am J Kidney Dis* 2008;52:859-67.

12972 *Franco MC, Nagasako SS, Machado PG, Nogueira PC, Pestana JO, Sesso R. Cystatin C and renal function in pediatric kidney transplant recipients. *Braz J Med Biol Res*. 2009;42:1225-9.

5412 *Franco MC, Nishida SK, Sesso R. GFR estimated from cystatin C versus creatinine in children born small for gestational age. *Am J Kidney Dis* 2008;51:925-32.

12973 *Fraser JS, Muller AF, Smith DJ, Newman DJ, Lamb EJ. Renal tubular injury is present in acute inflammatory bowel disease prior to the introduction of drug therapy. *Aliment Pharmacol Ther*. 2001;15:1131-7.

13551 *Friedman AN, Strother M, Quinney SK, Hall S, Perkins SM, Brizendine EJ, et al. Measuring the Glomerular Filtration Rate in Obese Individuals without Overt Kidney Disease. *Nephron - Clinical Practice* 2010;116:c224-c234.

5455 *Gabutti L, Ferrari N, Mombelli G, Marone C. Does cystatin C improve the precision of Cockcroft and Gault's creatinine clearance estimation? *J Nephrol* 2004;17:673-8.

12981 *Garcia A, Hermida J, Tutor JC. Estimation of the glomerular filtration rate from serum creatinine and cystatin C with regard to therapeutic digoxin monitoring. *J Clin Pharmacol*. 2007;47:1450-5.

5469 *Garcia Acuna JM, Gonzalez-Babarro E, Grigorian Shamagian L, Pena-Gil C, Vidal Perez R, Lopez-Lago AM, et al. Cystatin C provides more information than other renal function parameters for stratifying risk in patients with acute coronary syndrome. *Rev Esp Cardiol* 2009;62:510-9.

12982 *Garg AX, Salvadori M, Okell JM, Thiessen-Philbrook HR, Suri RS, Filler G, et al. Albuminuria and estimated GFR 5 years after Escherichia coli O157 hemolytic uremic syndrome: an update. *Am J Kidney Dis*. 2008;51:435-44.

5484 *Garlipp CR, Bottini PV, Queiroz RR, Afaz SH. Estimating glomerular filtration rate by measurement of cystatin C serum concentrations: a comparison with chrome-EDTA clearance. *Clin Chim Acta* 2008;393:125-7.

13552 *Gaspari F, Ferrari S, Stucchi N, Centemeri E, Carrara F, Pellegrino M, et al. Performance of different prediction equations for estimating renal function in kidney transplantation. *American Journal of Transplantation* 2004;4:1826-1835.

12984 *Geramizadeh B, Azarpira N, Ayatollahi M, Rais-Jalali GA, Aghdai M, Yaghoobi R, et al. Value of serum cystatin C as a marker of renal function in the early post kidney transplant period. *Saudi J Kidney Dis Transpl*. 2009;20:1015-7.

12985 *Gerbes AL, Gulberg V, Bilzer M, Vogeser M. Evaluation of serum cystatin C concentration as a marker of renal function in patients with cirrhosis of the liver. *Gut*. 2002;50:106-10.

5511 *Gerhardt T, Poge U, Stoffel-Wagner B, Ahrendt M, Wolff M, Spengler U, et al. Estimation of glomerular filtration rates after orthotopic liver transplantation: Evaluation of cystatin C-based equations. *Liver Transpl* 2006;12:1667-72.

13874 *Gerhardt T, Poge U, Stoffel-Wagner B, Palmedo H, Sauerbruch T, Woitas RP. Is beta-trace protein an alternative marker of glomerular filtration rate in liver transplant recipients? *Liver Int* 2011.

- 14075 *Gerhardt T, Pöge U, Stoffel-Wagner B, Palmedo H, Sauerbruch T, Woitas RP. Creatinine-based glomerular filtration rate estimation in patients with liver disease: the new Chronic Kidney Disease Epidemiology Collaboration equation is not better. *European Journal of Gastroenterology & Hepatology* 2011;23:969-973.
- 12986 *Giessing M, Liefeldt L. Getting the measure of renal function: Determination of best predicted creatinine in living-donor kidney renal transplant recipients based on donor kidney function. *Transplantation* 2008;86:513-514.
- 5531 *Giles PD, Rylance PB, Crothers DC. New results from the Modification of Diet in Renal Disease study: the importance of clinical outcomes in test strategies for early chronic kidney disease. *QJM* 2008;101:155-8.
- 12988 *Gokkusu CA, Ozden TA, Gul H, Yildiz A. Relationship between plasma Cystatin C and creatinine in chronic renal diseases and Tx-transplant patients. *Clin Biochem.* 2004;37:94-7.
- 12990 *Gomez Dammeier BH, Karanik E, Gluer S, Jesch NK, Kubler J, Latta K, et al. Anuria during pneumoperitoneum in infants and children: a prospective study. *J Pediatr Surg.* 2005;40:1454-8.
- 13338 *Gossmann J, Wilhelm A, Kachel HG, Jordan J, Sann U, Geiger H, et al. Long-term consequences of live kidney donation follow-up in 93% of living kidney donors in a single transplant center. *American Journal of Transplantation* 2005;5:2417-2424.
- 12992 *Gourishankar S, Courtney M, Jhangri GS, Cembrowski G, Pannu N. Serum cystatin C performs similarly to traditional markers of kidney function in the evaluation of donor kidney function prior to and following unilateral nephrectomy. *Nephrol Dial Transplant.* 2008;23:3004-9.
- 5597 *Grassi G, Abdelkawy H, Barsotti M, Paleologo G, Tregnaghi C, Rizzo G, et al. Living kidney transplantation: evaluation of renal function and morphology of potential donors. *Transplant Proc* 2009;41:1121-4.
- 13553 *Graziani F, Cei S, La Ferla F, Vano M, Gabriele M, Tonetti M. Effects of non-surgical periodontal therapy on the glomerular filtration rate of the kidney: an exploratory trial. *Journal of Clinical Periodontology* 2010;37:638-43.
- 12993 *Greco F, Hamza A, Wagner S, Hoda MR, Inferrera A, Lupo A, et al. Hand-Assisted Laparoscopic Living-Donor Nephrectomy Versus Open Surgery: Evaluation of Surgical Trauma and Late Graft Function in 82 Patients. *Transplantation Proceedings* 2009;41:4039-4043.
- 12994 *Groesbeck D, Kottgen A, Parekh R, Selvin E, Schwartz GJ, Coresh J, et al. Age, gender, and race effects on cystatin C levels in US adolescents. *Clin J Am Soc Nephrol.* 2008;3:1777-85.
- 5611 *Gronroos MH, Jahnukainen T, Irjala K, Harkonen R, Hurme S, Mottonen M, et al. Comparison of glomerular function tests in children with cancer. *Pediatr Nephrol* 2008;23:797-803.
- 13579 *Grubb A. Non-invasive estimation of glomerular filtration rate (GFR). The Lund model: Simultaneous use of cystatin C- and creatinine-based GFR-prediction equations, clinical data and an internal quality check. *Scand J Clin Lab Invest.* 2010;70:65-70.
- 1215 *Grubb A, Bjork J, Lindstrom V, Sterner G, Bondesson P, Nyman U. A cystatin C-based formula without anthropometric variables estimates glomerular filtration rate better than creatinine clearance using the Cockcroft-Gault formula. *Scand J Clin Lab Invest* 2005;65:153-62.
- 13559 *Grubb A, Bjork J, Nyman U, Pollak J, Bengzon J, Ostner G, et al. Cystatin C, a marker for successful aging and glomerular filtration rate, is not influenced by inflammation. *Scand J Clin Lab Invest* 2011.
- 5617 *Grubb A, Nyman U, Bjork J, Lindstrom V, Rippe B, Sterner G, et al. Simple cystatin C-based prediction equations for glomerular filtration rate compared with the modification of diet in renal disease prediction equation for adults and the Schwartz and the Counahan-Barratt prediction equations for children. *Clin Chem* 2005;51:1420-31.
- 5618 *Grubb A, Simonsen O, Sturfelt G, Truedsson L, Thysell H. Serum concentration of cystatin C, factor D and beta 2-microglobulin as a measure of glomerular filtration rate. *Acta Med Scand* 1985;218:499-503.

- 12996 *Gruev T, Chakalarovski K, Stojceva-Taneva O, Grueva A, Trencova K. Effects of glucocorticoid immunosuppression on serum cystatin C levels. *Journal of Medical Biochemistry* 2009;28:191-196.
- 540 *Gualano B, Ugrinowitsch C, Novaes RB, Artioli GG, Shimizu MH, Seguro AC, et al. Effects of creatine supplementation on renal function: a randomized, double-blind, placebo-controlled clinical trial. *Eur J Appl Physiol* 2008;103:33-40.
- 12998 *Gunther A, Burchard GD, Slevogt H, Abel W, Grobusch MP. Renal dysfunction in falciparum--malaria is detected more often when assessed by serum concentration of cystatin C instead of creatinine. *Trop Med Int Health*. 2002;7:931-4.
- 12999 *Haase M, Haase-Fielitz A, Bagshaw SM, Reade MC, Morgera S, Seevenayagam S, et al. Phase II, randomized, controlled trial of high-dose N-acetylcysteine in high-risk cardiac surgery patients. *Crit Care Med*. 2007;35:1324-31.
- 13000 *Haase M, Haase-Fielitz A, Ratnaike S, Reade MC, Bagshaw SM, Morgera S, et al. N-Acetylcysteine does not artifactually lower plasma creatinine concentration. *Nephrol Dial Transplant*. 2008;23:1581-7.
- 13001 *Haase-Fielitz A, Bellomo R, Devarajan P, Story D, Matalanis G, Dragun D, et al. Novel and conventional serum biomarkers predicting acute kidney injury in adult cardiac surgery--a prospective cohort study. *Crit Care Med*. 2009;37:553-60.
- 5666 *Halacova M, Kotaska K, Kukacka J, Vavrova V, Kuzelova M, Ticha J, et al. Serum cystatin C level for better assessment of glomerular filtration rate in cystic fibrosis patients treated by amikacin. *J Clin Pharm Ther* 2008;33:409-17.
- 5671 *Hallan S, Asberg A, Lindberg M, Johnsen H. Validation of the Modification of Diet in Renal Disease formula for estimating GFR with special emphasis on calibration of the serum creatinine assay. *Am J Kidney Dis* 2004;44:84-93.
- 13554 *Hallan SI, Orth SR. The KDOQI 2002 classification of chronic kidney disease: For whom the bell tolls. *Nephrology Dialysis Transplantation* 2010;25:2832-2836.
- 13002 *Hallberg P, Melhus H, Hansson LO, Larsson A. Cystatin C vs creatinine as markers of renal function in patients on digoxin treatment. *Ups J Med Sci*. 2004;109:247-53.
- 5686 *Han H, Dwyer JT, Selhub J, Jacques PF, Houser RF, Park JH, et al. Serum cystatin C is an independent predictor of total homocysteine levels in stable Korean renal transplant recipients with normal serum creatinine. *J Ren Nutr* 2001;11:149-54.
- 13003 *Han L, Bai X, Lin H, Sun X, Chen XM. Lack of independent relationship between age-related kidney function decline and carotid intima-media thickness in a healthy Chinese population. *Nephrol Dial Transplant*. 2010.
- 1230 *Hari P, Bagga A, Mahajan P, Lakshmy R. Effect of malnutrition on serum creatinine and cystatin C levels. *Pediatr Nephrol* 2007;22:1757-61.
- 5697 *Harmoinen A, Lehtimäki T, Korpela M, Turjanmaa V, Saha H. Diagnostic accuracies of plasma creatinine, cystatin C, and glomerular filtration rate calculated by the Cockcroft-Gault and Levey (MDRD) formulas. *Clin Chem* 2003;49:1223-5.
- 1231 *Harmoinen A, Ylinen E, Ala-Houhala M, Janas M, Kaila M, Kouri T. Reference intervals for cystatin C in pre- and full-term infants and children. *Pediatr Nephrol* 2000;15:105-8.
- 5698 *Harmoinen AP, Kouri TT, Wirta OR, Lehtimäki TJ, Rantalaiho V, Turjanmaa VM, et al. Evaluation of plasma cystatin C as a marker for glomerular filtration rate in patients with type 2 diabetes. *Clin Nephrol* 1999;52:363-70.
- 13555 *Harris KPG, Stribling B. Automated estimated GFR reporting: A new tool to promote safer prescribing in patients with chronic kidney disease? *Therapeutics and Clinical Risk Management* 2007;3:969-972.

13007 *Harten J, Crozier JE, McCreath B, Hay A, McMillan DC, McArdle CS, et al. Effect of intraoperative fluid optimisation on renal function in patients undergoing emergency abdominal surgery: a randomised controlled pilot study (ISRCTN 11799696). *Int J Surg*. 2008;6:197-204.

5700 *Hartlev LB, Rotbol CB, Bluhme H, Palshof T, Rehling M. [Monitoring renal function in patients treated with cytostatics]. *Ugeskr Laeger* 2008;170:3664-7.

1950 *Hayashi T, Nitta K, Hatano M, Nakauchi M, Nihei H. The serum cystatin C concentration measured by particle-enhanced immunonephelometry is well correlated with inulin clearance in patients with various types of glomerulonephritis. *Nephron* 1999;82:90-2.

13008 *Hayashi T, Nitta K, Uchida K, Honda K, Kobayashi H, Kawashima A, et al. Clinical assessment of serum cystatin C as a marker of glomerular filtration rate in patients with various renal diseases. *Clinical and Experimental Nephrology* 2000;4:133-136.

5729 *Hazar V, Gungor O, Guven AG, Aydin F, Akbas H, Gungor F, et al. Renal function after hematopoietic stem cell transplantation in children. *Pediatr Blood Cancer* 2009;53:197-202.

13009 *Hei ZQ, Li XY, Shen N, Pang HY, Zhou SL, Guan JQ. Prognostic values of serum cystatin C and beta2 microglobulin, urinary beta2 microglobulin and N-acetyl-beta-D-glucosaminidase in early acute renal failure after liver transplantation. *Chin Med J (Engl)*. 2008;121:1251-6.

13010 *Heilman RL, Mazur MJ. Cystatin C as a more sensitive indicator of diminished glomerular filtration rate. *Liver Transplantation* 2005;11:264-266.

5746 *Heise D, Waeschle RM, Schlobohm J, Wessels J, Quintel M. Utility of cystatin C for assessment of renal function after cardiac surgery. *Nephron Clin Pract* 2009;112:c107-14.

13011 *Hekmat R, Eshraghi H. Comparison of serum cystatin C and creatinine levels to evaluate early renal function after kidney transplantation. *Iranian Journal of Medical Sciences* 2009;34:94-99.

1236 *Helin I, Axenram M, Grubb A. Serum cystatin C as a determinant of glomerular filtration rate in children. *Clin Nephrol* 1998;49:221-5.

5755 *Hellerstein S, Berenbom M, Erwin P, Wilson N, DiMaggio S. The ratio of urinary cystatin C to urinary creatinine for detecting decreased GFR. *Pediatr Nephrol* 2004;19:521-5.

13013 *Herget-Rosenthal S, Pietruck F, Volbracht L, Philipp T, Kribben A. Serum cystatin C--a superior marker of rapidly reduced glomerular filtration after uninephrectomy in kidney donors compared to creatinine. *Clin Nephrol*. 2005;64:41-6.

13014 *Herget-Rosenthal S, Trabold S, Huesing J, Heemann U, Philipp T, Kribben A. Cystatin C--an accurate marker of glomerular filtration rate after renal transplantation? *Transpl Int*. 2000;13:285-9.

13015 *Herget-Rosenthal S, Trabold S, Pietruck F, Holtmann M, Philipp T, Kribben A. Cystatin C: efficacy as screening test for reduced glomerular filtration rate. *Am J Nephrol*. 2000;20:97-102.

5778 *Herget-Rosenthal S, van Wijk JA, Brocker-Preuss M, Bokenkamp A. Increased urinary cystatin C reflects structural and functional renal tubular impairment independent of glomerular filtration rate. *Clin Biochem* 2007;40:946-51.

13016 *Heringlake M, Wernerus M, Grunefeld J, Klaus S, Heinze H, Bechtel M, et al. The metabolic and renal effects of adrenaline and milrinone in patients with myocardial dysfunction after coronary artery bypass grafting. *Crit Care*. 2007;11:R51.

13018 *Hermida J, Romero R, Tutor JC. Relationship between serum cystatin C and creatinine in kidney and liver transplant patients. *Clin Chim Acta*. 2002;316:165-70.

3441 *Hermida J, Tutor JC. Serum cystatin C for the prediction of glomerular filtration rate with regard to the dose adjustment of amikacin, gentamicin, tobramycin, and vancomycin. *Ther Drug Monit* 2006;28:326-31.

- 13019 *Hermida J, Tutor JC. Comparison of estimated glomerular filtration rates from serum creatinine and cystatin C in patients with impaired creatinine production. *Clin Lab.* 2006;52:483-90.
- 5793 *Herrero-Morin JD, Malaga S, Fernandez N, Rey C, Dieguez MA, Solis G, et al. Cystatin C and beta2-microglobulin: markers of glomerular filtration in critically ill children. *Crit Care* 2007;11:R59.
- 5830 *Hjorth L, Wiebe T, Karpman D. Correct evaluation of renal glomerular filtration rate requires clearance assays. *Pediatr Nephrol* 2002;17:847-51.
- 1246 *Hoek FJ, Kemperman FA, Krediet RT. A comparison between cystatin C, plasma creatinine and the Cockcroft and Gault formula for the estimation of glomerular filtration rate. *Nephrol Dial Transplant* 2003;18:2024-31.
- 5842 *Hoek FJ, Korevaar JC, Dekker FW, Boeschoten EW, Krediet RT. Estimation of residual glomerular filtration rate in dialysis patients from the plasma cystatin C level. *Nephrol Dial Transplant* 2007;22:1633-8.
- 13023 *Hoffmann U, Fischereder M, Kruger B, Drobnik W, Kramer BK. The value of N-acetylcysteine in the prevention of radiocontrast agent-induced nephropathy seems questionable. *J Am Soc Nephrol.* 2004;15:407-10.
- 5850 *Hojs R, Bevc S, Antolinc B, Gorenjak M, Puklavec L. Serum cystatin C as an endogenous marker of renal function in the elderly. *Int J Clin Pharmacol Res* 2004;24:49-54.
- 5851 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Serum cystatin C as an endogenous marker of renal function in patients with mild to moderate impairment of kidney function. *Nephrol Dial Transplant* 2006;21:1855-62.
- 5852 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Serum cystatin C-based equation compared to serum creatinine-based equations for estimation of glomerular filtration rate in patients with chronic kidney disease. *Clin Nephrol* 2008;70:10-7.
- 5853 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Serum cystatin C as an endogenous marker of renal function in patients with chronic kidney disease. *Ren Fail* 2008;30:181-6.
- 13505 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Serum cystatin C-based formulas for prediction of glomerular filtration rate in patients with chronic kidney disease. *Nephron - Clinical Practice* 2010;114:c118-c126.
- 13875 *Hojs R, Bevc S, Ekart R, Gorenjak M, Puklavec L. Kidney function estimating equations in patients with chronic kidney disease. *Int J Clin Pract* 2011;65:458-64.
- 13580 *Hoke M, Amighi J, Mlekusch W, Schlager O, Exner M, Sabeti S, et al. Cystatin C and the risk for cardiovascular events in patients with asymptomatic carotid atherosclerosis. *Stroke.* 2010;41:674-9.
- 13025 *Hollevoet K, Bernard D, De Geeter F, Walgraeve N, Van Den Eeckhaut A, Vanholder R, et al. Glomerular filtration rate is a confounder for the measurement of soluble mesothelin in serum. *Clinical Chemistry* 2009;55:1431-1433.
- 5861 *Holmquist P, Torffvit O, Sjoblad S. Metabolic status in diabetes mellitus affects markers for glomerular filtration rate. *Pediatr Nephrol* 2003;18:536-40.
- 5864 *Holweger K, Lipp HP, Dietz K, Hartmann JT, Bokemeyer C. Novel algorithm for more accurate calculation of renal function in adults with cancer. *Ann Pharmacother* 2008;42:1749-57.
- 13027 *Hoppe A, Seronie-Vivien S, Thomas F, Delord JP, Malard L, Canal P, et al. Serum cystatin C is a better marker of topotecan clearance than serum creatinine. *Clin Cancer Res.* 2005;11:3038-44.
- 1986 *Horackova M, Schuck O, Komers R, Charvat J, Teplan V, Kvapil M. Effect of rofecoxib on the glomerular filtration rate, proteinuria and the renin-angiotensin-aldosterone system in elderly subjects with chronic renal impairment. *Int J Clin Pharmacol Ther* 2005;43:413-9.

13857 *Huang SH, Macnab JJ, Sontrop JM, Filler G, Gallo K, Lindsay RM, et al. Performance of the creatinine-based and the cystatin C-based glomerular filtration rate (GFR) estimating equations in a heterogenous sample of patients referred for nuclear GFR testing. *Transl Res* 2011;157:357-67.

13577 *Huang SH, Sharma AP, Yasin A, Lindsay RM, Clark WF, Filler G. Hyperfiltration Affects Accuracy of Creatinine eGFR Measurement. *Clin J Am Soc Nephrol*. 2011;6:274-280.

14076 *Hudson JQ, Nyman HA. Use of estimated glomerular filtration rate for drug dosing in the chronic kidney disease patient. *Current Opinion in Nephrology & Hypertension* 2011;20:482-491.

13031 *Hutchison CA, Cockwell P, Harding S, Mead GP, Bradwell AR, Barnett AH. Quantitative assessment of serum and urinary polyclonal free light chains in patients with type II diabetes: an early marker of diabetic kidney disease? *Expert Opin Ther Targets*. 2008;12:667-76.

13032 *Hutchison CA, Harding S, Hewins P, Mead GP, Townsend J, Bradwell AR, et al. Quantitative assessment of serum and urinary polyclonal free light chains in patients with chronic kidney disease. *Clin J Am Soc Nephrol*. 2008;3:1684-90.

5934 *Hwang SJ, Yang Q, Meigs JB, Pearce EN, Fox CS. A genome-wide association for kidney function and endocrine-related traits in the NHLBI's Framingham Heart Study. *BMC Med Genet* 2007;8 Suppl 1:S10.

13033 *Hynninen MS, Niemi TT, Poyhia R, Raininko EI, Salmenpera MT, Lepantalo MJ, et al. N-acetylcysteine for the prevention of kidney injury in abdominal aortic surgery: a randomized, double-blind, placebo-controlled trial. *Anesth Analg*. 2006;102:1638-45.

13034 *Ichimoto E, Jo K, Kobayashi Y, Inoue T, Nakamura Y, Kuroda N, et al. Prognostic significance of cystatin C in patients with ST-elevation myocardial infarction. *Circ J*. 2009;73:1669-73.

13837 *Iliadis F, Didangelos T, Ntemka A, Makedou A, Moralidis E, Gotzamani-Psarakou A, et al. Glomerular filtration rate estimation in patients with type 2 diabetes: creatinine- or cystatin C-based equations? *Diabetologia* 2011.

5973 *Ishani A, Paudel M, Taylor BC, Barrett-Connor E, Jamal S, Canales M, et al. Renal function and rate of hip bone loss in older men: the Osteoporotic Fractures in Men Study. *Osteoporos Int* 2008;19:1549-56.

3479 *Islekel H, Soylu A, Altun Z, Yis U, Turkmen M, Kavukcu S. Serum and urine cystatin C levels in children with post-pyelonephritic renal scarring: a pilot study. *Int Urol Nephrol* 2007;39:1241-50.

5979 *Issa N, Meyer KH, Arrigain S, Choure G, Fatica RA, Nurko S, et al. Evaluation of creatinine-based estimates of glomerular filtration rate in a large cohort of living kidney donors. *Transplantation* 2008;86:223-30.

13037 *Iwanaga Y, Miyazaki S. Cystatin C in a high-risk combination. *Circ J*. 2009;73:1593-4.

5990 *Ix JH, Chertow GM, Shlipak MG, Brandenburg VM, Ketteler M, Whooley MA. Fetuin-A and kidney function in persons with coronary artery disease--data from the Heart and Soul Study. *Nephrol Dial Transplant* 2006;21:2144-51.

13581 *Ix JH, de Boer IH, Wassel CL, Criqui MH, Shlipak MG, Whooley MA. Urinary creatinine excretion rate and mortality in persons with coronary artery disease: the Heart and Soul Study. *Circulation*. 2010;121:1295-303.

13341 *Ix JH, Katz R, De Boer IH, Kestenbaum BR, Allison MA, Siscovick DS, et al. Association of chronic kidney disease with the spectrum of ankle brachial index the CHS (Cardiovascular Health Study). *J Am Coll Cardiol*. 2009;54:1176-84.

5992 *Ix JH, Shlipak MG, Chertow GM, Whooley MA. Association of cystatin C with mortality, cardiovascular events, and incident heart failure among persons with coronary heart disease: data from the Heart and Soul Study. *Circulation* 2007;115:173-9.

6027 *Jaroszewicz J, Wiercinska-Drapalo A, Lapinski TW, Prokopowicz D, Rogalska M, Parfieniuk A. Does HAART improve renal function? An association between serum cystatin C concentration, HIV viral load and HAART duration. *Antivir Ther* 2006;11:641-5.

- 13876 *Jarvela K, Maaranen P, Harmoinen A, Huhtala H, Sisto T. Cystatin C in diabetics as a marker of mild renal insufficiency after CABG. *Ann Thorac Cardiovasc Surg* 2011;17:277-82.
- 13040 *Jayagopal V, Keevil BG, Atkin SL, Jennings PE, Kilpatrick ES. Paradoxical changes in cystatin C and serum creatinine in patients with hypo- and hyperthyroidism. *Clin Chem*. 2003;49:680-1.
- 13041 *Jenkins AJ, Zhang SX, Rowley KG, Karschimkus CS, Nelson CL, Chung JS, et al. Increased serum pigment epithelium-derived factor is associated with microvascular complications, vascular stiffness and inflammation in Type 1 diabetes. *Diabet Med*. 2007;24:1345-51.
- 13042 *Jenkins MA, Brown DJ, Ierino FL, Ratnaike SI. Cystatin C for estimation of glomerular filtration rate in patients with spinal cord injury. *Ann Clin Biochem*. 2003;40:364-8.
- 13044 *Jernberg T, Lindahl B, James S, Larsson A, Hansson LO, Wallentin L. Cystatin C: a novel predictor of outcome in suspected or confirmed non-ST-elevation acute coronary syndrome. *Circulation*. 2004;110:2342-8.
- 13045 *John GT, Fleming JJ, Talaulikar GS, Selvakumar R, Thomas PP, Jacob CK. Measurement of renal function in kidney donors using serum cystatin C and beta(2)-microglobulin. *Ann Clin Biochem*. 2003;40:656-8.
- 6057 *Johnson TM, 2nd, Sands JM, Ouslander JG. A prospective evaluation of the glomerular filtration rate in older adults with frequent nighttime urination. *J Urol* 2002;167:146-50.
- 13047 *Jonasson T, Ohlin H, Andersson A, Arnadottir M, Hultberg B. Renal function exerts only a minor influence on high plasma homocysteine concentrations in patients with acute coronary syndromes. *Clin Chem Lab Med*. 2002;40:137-42.
- 13048 *Jones CY, Jones CA, Wilson IB, Knox TA, Levey AS, Spiegelman D, et al. Cystatin C and creatinine in an HIV cohort: the nutrition for healthy living study. *Am J Kidney Dis*. 2008;51:914-24.
- 13339 *Jones TE, Peter JV, Field J. Aminoglycoside clearance is a good estimate of creatinine clearance in intensive care unit patients. *Anaesth Intensive Care*. 2009;37:944-52.
- 6065 *Jonsson AS, Flodin M, Hansson LO, Larsson A. Estimated glomerular filtration rate (eGFR_{CystC}) from serum cystatin C shows strong agreement with iohexol clearance in patients with low GFR. *Scand J Clin Lab Invest* 2007;67:801-9.
- 13050 *Jovanovic D, Krstivojevic P, Obradovic I, Durdevic V, Dukanovic L. Serum cystatin C and beta2-microglobulin as markers of glomerular filtration rate. *Ren Fail*. 2003;25:123-33.
- 13051 *Jung K, Jung M. Cystatin C: a promising marker of glomerular filtration rate to replace creatinine. *Nephron*. 1995;70:370-1.
- 13052 *Kalansooriya A, Holbrook I, Jennings P, Whiting PH. Serum cystatin C, enzymuria, tubular proteinuria and early renal insult in type 2 diabetes. *Br J Biomed Sci*. 2007;64:121-3.
- 6132 *Karstila K, Harmoinen AP, Lehtimaki TJ, Korpela MM, Mustonen JT, Saha HH. Measurement of the kidney function in patients with rheumatoid arthritis: plasma cystatin C versus ⁵¹Cr-EDTA clearance. *Nephron Clin Pract* 2008;108:c284-90.
- 13054 *Kato K, Sato N, Yamamoto T, Iwasaki YK, Tanaka K, Mizuno K. Valuable markers for contrast-induced nephropathy in patients undergoing cardiac catheterization. *Circ J*. 2008;72:1499-505.
- 13055 *Kazama JJ, Kutsuwada K, Ataka K, Maruyama H, Gejyo F. Serum cystatin C reliably detects renal dysfunction in patients with various renal diseases. *Nephron*. 2002;91:13-20.
- 13582 *Kees MG, Hilpert JW, Gnewuch C, Kees F, Voegeler S. Clearance of vancomycin during continuous infusion in Intensive Care Unit patients: correlation with measured and estimated creatinine clearance and serum cystatin C. *Int J Antimicrob Agents*. 2010;36:545-8.

- 6150 *Keller C, Katz R, Cushman M, Fried LF, Shlipak M. Association of kidney function with inflammatory and procoagulant markers in a diverse cohort: a cross-sectional analysis from the Multi-Ethnic Study of Atherosclerosis (MESA). *BMC Nephrol* 2008;9:9.
- 13056 *Keller CR, Odden MC, Fried LF, Newman AB, Angleman S, Green CA, et al. Kidney function and markers of inflammation in elderly persons without chronic kidney disease: the health, aging, and body composition study. *Kidney Int.* 2007;71:239-44.
- 13057 *Keller T, Messow CM, Lubos E, Nicaud V, Wild PS, Rupprecht HJ, et al. Cystatin C and cardiovascular mortality in patients with coronary artery disease and normal or mildly reduced kidney function: results from the AtheroGene study. *Eur Heart J.* 2009;30:314-20.
- 13058 *Kestenbaum B, Rudser KD, de Boer IH, Peralta CA, Fried LF, Shlipak MG, et al. Differences in kidney function and incident hypertension: the multi-ethnic study of atherosclerosis. *Ann Intern Med.* 2008;148:501-8.
- 13059 *Keyes R, Bagshaw SM. Early diagnosis of acute kidney injury in critically ill patients. *Expert Review of Molecular Diagnostics* 2008;8:455-464.
- 13060 *Kiefer D. In the news Cystatin C valuable in detecting kidney dysfunction. *Life Extension* 2006;12:14.
- 13061 *Kilic T, Oner G, Ural E, Yumuk Z, Sahin T, Bildirici U, et al. Comparison of the long-term prognostic value of Cystatin C to other indicators of renal function, markers of inflammation and systolic dysfunction among patients with acute coronary syndrome. *Atherosclerosis.* 2009;207:552-8.
- 6185 *Kilpatrick ES, Keevil BG, Addison GM. Does adjustment of GFR to extracellular fluid volume improve the clinical utility of cystatin C? *Arch Dis Child* 2000;82:499-502.
- 13062 *Kim BJ, Sung KC, Kim BS, Kang JH, Lee KB, Kim H, et al. Effect of N-Acetylcysteine on cystatin C-Based renal function after Elective coronary angiography (ENABLE Study): A prospective, randomized trial. *Int J Cardiol.* 2008.
- 13799 *Kim DJ, Kang HS, Choi HS, Cho HJ, Kim ES, Keum B, et al. Serum cystatin C level is a useful marker for the evaluation of renal function in patients with cirrhotic ascites and normal serum creatinine levels. *Korean J Hepatol* 2011;17:130-8.
- 13556 *Kim WR, Leise M, Pedersen R, Therneau T, Rosen C, Pungpapong S. Accuracy of eGFR_{cys} in measuring renal function in liver transplant recipients. *Hepatology International* 2010;4:28.
- 13063 *Kimmel M, Butscheid M, Brenner S, Kuhlmann U, Klotz U, Alschner DM. Improved estimation of glomerular filtration rate by serum cystatin C in preventing contrast induced nephropathy by N-acetylcysteine or zinc-- preliminary results. *Nephrol Dial Transplant.* 2008;23:1241-5.
- 13574 *Kiyosue A, Hirata Y, Ando J, Fujita H, Morita T, Takahashi M, et al. Plasma cystatin C concentration reflects the severity of coronary artery disease in patients without chronic kidney disease. *Circ J.* 2010;74:2441-7.
- 6222 *Kleber M, Cybulla M, Bauchmuller K, Ihorst G, Koch B, Engelhardt M. Monitoring of renal function in cancer patients: an ongoing challenge for clinical practice. *Ann Oncol* 2007;18:950-8.
- 63 *Kleinert J, Lorenz M, Hauser AC, Becherer A, Staudenherz A, Fodinger M, et al. Measurement of renal function in patients with Fabry disease. *Acta Paediatr Suppl* 2005;94:19-23; discussion 9-10.
- 13065 *Knight EL, Verhave JC, Spiegelman D, Hillege HL, de Zeeuw D, Curhan GC, et al. Factors influencing serum cystatin C levels other than renal function and the impact on renal function measurement. *Kidney Int.* 2004;65:1416-21.
- 13066 *Kocak H, Oner-Iyidogan Y, Gurdol F, Kocak T, Esin D. The relation between serum MDA and cystatin C levels in chronic spinal cord injury patients. *Clin Biochem.* 2005;38:1034-7.
- 13067 *Kocak H, Oner-Iyidogan Y, Gurdol F, Kocak T, Nane I, Genc S. Cystatin-C and creatinine as indices of glomerular filtration rate in the immediate follow-up of renal transplant patients. *Clin Exp Med.* 2005;5:14-9.

- 2035 *Koch M, Beien A, Fussha Ller A, Zitta S, Haastert B, Trapp R. Impact of age, body mass index, insulin resistance and proteinuria on the kidney function in obese patients with Type 2 diabetes and renal insufficiency. *Clin Nephrol* 2008;69:10-7.
- 13068 *Kociol R, Rogers J, Shaw A. Organ cross talk in the critically ill: The heart and kidney. *Blood Purification* 2009;27:311-320.
- 13069 *Koenig W. Is elevated cystatin C a predictor of cardiovascular risk in elderly people without chronic kidney disease? Commentary. *Nature Clinical Practice Cardiovascular Medicine* 2007;4:76-77.
- 13070 *Koenig W, Twardella D, Brenner H, Rothenbacher D. Plasma concentrations of cystatin C in patients with coronary heart disease and risk for secondary cardiovascular events: more than simply a marker of glomerular filtration rate. *Clin Chem*. 2005;51:321-7.
- 13072 *Kravaritou M, Thanopoulou A, Karamanos B, Kofinis A, Noutsou M, Spanou E, et al. Evidence that even "normal" albuminuria may denote incipient GFR reduction in patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract*. 2009;85:317-21.
- 13073 *Krawczuk-Rybak M, Kuzmicz M, Wysocka J. Renal function during and after treatment for acute lymphoblastic leukemia in children. *Pediatr Nephrol*. 2005;20:782-5.
- 13074 *Krediet RT. How to preserve residual renal function in patients with chronic kidney disease and on dialysis? *Nephrology Dialysis Transplantation* 2006;21:ii42-ii46.
- 6285 *Krieser D, Rosenberg AR, Kainer G, Naidoo D. The relationship between serum creatinine, serum cystatin C and glomerular filtration rate in pediatric renal transplant recipients: a pilot study. *Pediatr Transplant* 2002;6:392-5.
- 1321 *Kristensen K, Lindstrom V, Schmidt C, Blirup-Jensen S, Grubb A, Wide-Swensson D, et al. Temporal changes of the plasma levels of cystatin C, beta-trace protein, beta2-microglobulin, urate and creatinine during pregnancy indicate continuous alterations in the renal filtration process. *Scand J Clin Lab Invest* 2007;67:612-8.
- 13075 *Kristensen K, Wide-Swensson D, Schmidt C, Blirup-Jensen S, Lindstrom V, Strevens H, et al. Cystatin C, beta-2-microglobulin and beta-trace protein in pre-eclampsia. *Acta Obstet Gynecol Scand*. 2007;86:921-6.
- 13076 *Krum H, lyngkaran P, Lekawanvijit S. Pharmacologic management of the cardiorenal syndrome in heart failure. *Current Heart Failure Reports* 2009;6:105-111.
- 13557 *Kukla A, El-Shahawi Y, Leister E, Kasiske B, Mauer M, Matas A, et al. GFR-estimating models in kidney transplant recipients on a steroid-free regimen. In: *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*; 2010. p 1653-61.
- 13507 *Kwong YT, Stevens LA, Selvin E, Zhang YL, Greene T, Van Lente F, et al. Imprecision of urinary iothalamate clearance as a gold-standard measure of GFR decreases the diagnostic accuracy of kidney function estimating equations. In: *American journal of kidney diseases : the official journal of the National Kidney Foundation*; 2010. p 39-49.
- 1330 *Kyhse-Andersen J, Schmidt C, Nordin G, Andersson B, Nilsson-Ehle P, Lindstrom V, et al. Serum cystatin C, determined by a rapid, automated particle-enhanced turbidimetric method, is a better marker than serum creatinine for glomerular filtration rate. *Clin Chem* 1994;40:1921-6.
- 13079 *Laisalmi-Kokki M, Pesonen E, Kokki H, Valta P, Pitkanen M, Teppo AM, et al. Potentially detrimental effects of N-acetylcysteine on renal function in knee arthroplasty. *Free Radic Res*. 2009;43:691-6.
- 13080 *Lamb E. Assessment of kidney function in adults. *Medicine* 2007;35:359-364.
- 6343 *Lamb EJ, O'Riordan SE, Delaney MP. Kidney function in older people: pathology, assessment and management. *Clin Chim Acta* 2003;334:25-40.
- 6345 *Lamb EJ, Tomson CR, Roderick PJ, Clinical Sciences Reviews Committee of the Association for Clinical B. Estimating kidney function in adults using formulae. *Ann Clin Biochem* 2005;42:321-45.

- 6347 *Lamb EJ, Webb MC, Simpson DE, Coakley AJ, Newman DJ, O'Riordan SE. Estimation of glomerular filtration rate in older patients with chronic renal insufficiency: is the modification of diet in renal disease formula an improvement? *J Am Geriatr Soc* 2003;51:1012-7.
- 13081 *Lambermont B, D'Orio V. Cystatin C blood level as a risk factor for death after heart surgery. *Eur Heart J*. 2007;28:2818; author reply 2818-9.
- 13082 *Lameire N. Contrast-induced nephropathy in the critically-ill patient: Focus on emergency screening and prevention. *Acta Clinica Belgica* 2007;62:346-352.
- 13083 *Lameire N, Van Biesen W, Hoste E, Vanholder R. The prevention of acute kidney injury: An in-depth narrative review Part 1: Volume resuscitation and avoidance of drug- and nephrotoxin-induced AKI. *NDT Plus* 2008;1:392-402.
- 6357 *Lane BR, Poggio ED, Herts BR, Novick AC, Campbell SC. Renal function assessment in the era of chronic kidney disease: renewed emphasis on renal function centered patient care. *J Urol* 2009;182:435-43; discussion 443-4.
- 13084 *Lang JD, Chaiwat O. Clinical use of hydroxyethylstarch: Colloidal confusion or clarity? *Critical Care Medicine* 2007;35:2864-2866.
- 6362 *Lankisch P, Wessalowski R, Maisonneuve P, Haghgu M, Hermsen D, Kramm CM. Serum cystatin C is a suitable marker for routine monitoring of renal function in pediatric cancer patients, especially of very young age. *Pediatr Blood Cancer* 2006;46:767-72.
- 13085 *Lapinski TW, Parfieniuk A, Rogalska-Plonska M, Czajkowska J, Flisiak R. Prevalence of cryoglobulinaemia in hepatitis C virus- and hepatitis C virus/human immunodeficiency virus-infected individuals: implications for renal function. *Liver Int*. 2009;29:1158-61.
- 6365 *Larsson A. Cystatin C: an emerging glomerular filtration rate marker. *Scand J Clin Lab Invest* 2005;65:89-91.
- 13086 *Larsson A. Author response to: Reference values for clinical chemistry tests during normal pregnancy. *BJOG: An International Journal of Obstetrics and Gynaecology* 2008;115:1716-1717.
- 644 *Larsson A, Akerstedt T, Hansson LO, Axelsson J. Circadian variability of cystatin C, creatinine, and glomerular filtration rate (GFR) in healthy men during normal sleep and after an acute shift of sleep. *Chronobiol Int* 2008;25:1047-61.
- 5 *Larsson A, Flodin M, Hansson LO, Carlsson L. Patient selection has a strong impact on cystatin C and Modification of Diet in Renal Disease (MDRD) estimated glomerular filtration rate. *Clin Biochem* 2008;41:1355-61.
- 13088 *Larsson A, Helmersson J, Hansson LO, Basu S. Increased serum cystatin C is associated with increased mortality in elderly men. *Scand J Clin Lab Invest*. 2005;65:301-5.
- 6368 *Larsson A, Malm J, Grubb A, Hansson LO. Calculation of glomerular filtration rate expressed in mL/min from plasma cystatin C values in mg/L. *Scand J Clin Lab Invest* 2004;64:25-30.
- 13089 *Lassus J, Harjola VP, Sund R, Siirila-Waris K, Melin J, Peuhkurinen K, et al. Prognostic value of cystatin C in acute heart failure in relation to other markers of renal function and NT-proBNP. *Eur Heart J*. 2007;28:1841-7.
- 1340 *Laterza OF, Price CP, Scott MG. Cystatin C: an improved estimator of glomerular filtration rate? *Clin Chem* 2002;48:699-707.
- 133 *Launay-Vacher V, Chatelut E, Lichtman SM, Wildiers H, Steer C, Aapro M, et al. Renal insufficiency in elderly cancer patients: International Society of Geriatric Oncology clinical practice recommendations. *Ann Oncol* 2007;18:1314-21.

- 6388 *Le Bricon T, Leblanc I, Benlakehal M, Gay-Bellile C, Erlich D, Boudaoud S. Evaluation of renal function in intensive care: plasma cystatin C vs. creatinine and derived glomerular filtration rate estimates. *Clin Chem Lab Med* 2005;43:953-7.
- 6389 *Le Bricon T, Thervet E, Benlakehal M, Bousquet B, Legendre C, Erlich D. Changes in plasma cystatin C after renal transplantation and acute rejection in adults. *Clin Chem* 1999;45:2243-9.
- 13090 *Le Bricon T, Thervet E, Froissart M, Benlakehal M, Bousquet B, Legendre C, et al. Plasma cystatin C is superior to 24-h creatinine clearance and plasma creatinine for estimation of glomerular filtration rate 3 months after kidney transplantation. *Clin Chem*. 2000;46:1206-7.
- 13091 *Leach TD, Kitiyakara C, Price CP, Stevens JM, Newman DJ. Prognostic significance of serum cystatin c concentrations in renal transplant recipients: 5-year follow-up. *Transplant Proc*. 2002;34:1152-8.
- 6401 *Ledger S. Laboratory assessment of kidney function. *CANNT J* 2006;16:34-7; quiz 38-9.
- 13092 *Ledoux D, Monchi M, Chapelle JP, Damas P. Cystatin C blood level as a risk factor for death after heart surgery. *Eur Heart J*. 2007;28:1848-53.
- 13093 *Lee BW, Ihm SH, Choi MG, Yoo HJ. The comparison of cystatin C and creatinine as an accurate serum marker in the prediction of type 2 diabetic nephropathy. *Diabetes Res Clin Pract*. 2007;78:428-34.
- 13094 *Lee SH, Park SA, Ko SH, Yim HW, Ahn YB, Yoon KH, et al. Insulin resistance and inflammation may have an additional role in the link between cystatin C and cardiovascular disease in type 2 diabetes mellitus patients. *Metabolism*. 2009.
- 6413 *Lee SM, Kim JH, Lee YM, Lee JS, Kim HD. Evaluation of renal function in children with mitochondrial respiratory chain complex defect: usefulness of cystatin C. *Acta Paediatr* 2009;98:1014-8.
- 13095 *Leung N. A tale of two markers. *Liver Transpl*. 2002;8:600-2.
- 13096 *Lewerin C, Ljungman S, Nilsson-Ehle H. Glomerular filtration rate as measured by serum cystatin C is an important determinant of plasma homocysteine and serum methylmalonic acid in the elderly. *J Intern Med*. 2007;261:65-73.
- 13097 *Levin A. Cystatin C, serum creatinine, and estimates of kidney function: searching for better measures of kidney function and cardiovascular risk. *Ann Intern Med*. 2005;142:586-8.
- 6450 *Lewis J, Agodoa L, Cheek D, Greene T, Middleton J, O'Connor D, et al. Comparison of cross-sectional renal function measurements in African Americans with hypertensive nephrosclerosis and of primary formulas to estimate glomerular filtration rate. *Am J Kidney Dis* 2001;38:744-53.
- 13098 *Li FK, Ho SKN, Yip TPS, Tse KC, Chan TM, Lai KN. Cystatin C assay for the detection of renal dysfunction in Chinese renal transplant recipients. *Clinica Chimica Acta* 2002;322:133-137.
- 13394 *Li HX, Xu GB, Wang XJ, Zhang XC, Yang JM. Diagnostic accuracy of various glomerular filtration rates estimating equations in patients with chronic kidney disease and diabetes. *Chinese Medical Journal* 2010;123:745-751.
- 13100 *Lindstedt G. Multifactorial regulation of plasma natriuretic peptides calls for supplementary measurements. *Scand Cardiovasc J*. 2001;35:355-8.
- 13101 *Lindstrom K, Kindgren L, Zafirova T, Frisenette-Fich C. [Adverse drug effects among the elderly can be reduced]. *Lakartidningen*. 2007;104:242-4.
- 6504 *Ling Q, Xu X, Li JJ, Chen J, Shen JW, Zheng SS. Alternative definition of acute kidney injury following liver transplantation: based on serum creatinine and cystatin C levels. *Transplant Proc* 2007;39:3257-60.
- 6511 *Linzbach S, Samigullin A, Yilmaz S, Tsioga M, Zeiher AM, Spyridopoulos I. Role of N-terminal pro-brain natriuretic peptide and cystatin C to estimate renal function in patients with and without heart failure. *Am J Cardiol* 2009;103:1128-33.

13658 *Lippi G, Franchini M, Targher G. Detection of chronic kidney disease in hospitalized patients: Is one estimating glomerular filtration rate equation better than another? *European Journal of Internal Medicine* 2011;22:119-120.

13103 *Lippi G, Salvagno GL, Montagnana M, Targher G, Franchini M, Guidi GC. No correlation between lipoprotein(a) and biochemical markers of renal function in the general population. *Arch Pathol Lab Med*. 2008;132:1436-8.

13104 *Liu J, Ma L, Yang J, Ren A, Sun Z, Yan G, et al. Increased serum cathepsin S in patients with atherosclerosis and diabetes. *Atherosclerosis*. 2006;186:411-9.

6556 *Louvar DW, Rogers TB, Bailey RF, Matas AJ, Ibrahim HN. Cystatin C is not superior to creatinine-based models in estimating glomerular filtration rate in former kidney donors. *Transplantation* 2007;84:1112-7.

13106 *Lucchese B. Diagnosis: Cystatin C is not an unbiased marker of GFR. *Nature Reviews Nephrology* 2009;5:305.

275 *Lufft V, Hoogestraat-Lufft L, Fels LM, Egbeyong-Baiyee D, Olbricht CJ, Galanski M. Angiography for renal artery stenosis: no additional impairment of renal function by angioplasty. *Eur Radiol* 2002;12:804-9.

13107 *Lupovitch A. More accurate alternatives to serum creatinine for evaluating glomerular filtration rate. *Clin Chem*. 2002;48:2297-8; author reply 2298.

1369 *Ma YC, Zuo L, Chen JH, Luo Q, Yu XQ, Li Y, et al. Improved GFR estimation by combined creatinine and cystatin C measurements. *Kidney Int* 2007;72:1535-42.

6588 *Maahs DM, Ogden LG, Kretowski A, Snell-Bergeon JK, Kinney GL, Berl T, et al. Serum cystatin C predicts progression of subclinical coronary atherosclerosis in individuals with type 1 diabetes. *Diabetes* 2007;56:2774-9.

13108 *Maahs DM, Snell-Bergeon JK, Hokanson JE, Kinney GL, Berl T, Rewers M, et al. Relationship between cystatin C and coronary artery atherosclerosis progression differs by type 1 diabetes. *Diabetes Technol Ther*. 2010;12:25-33.

13109 *Macdonald J, Marcora S, Lemmey A, Jibani M. In Reply. *American Journal of Kidney Diseases* 2007;49:561.

6598 *Macisaac RJ, Tsalamandris C, Thomas MC, Premaratne E, Panagiotopoulos S, Smith TJ, et al. Estimating glomerular filtration rate in diabetes: a comparison of cystatin-C- and creatinine-based methods. *Diabetologia* 2006;49:1686-9.

1371 *Macisaac RJ, Tsalamandris C, Thomas MC, Premaratne E, Panagiotopoulos S, Smith TJ, et al. The accuracy of cystatin C and commonly used creatinine-based methods for detecting moderate and mild chronic kidney disease in diabetes. *Diabet Med* 2007;24:443-8.

13110 *Madero M, Sarnak MJ. Association of cystatin C with adverse outcomes. *Curr Opin Nephrol Hypertens*. 2009;18:258-63.

6606 *Madero M, Sarnak MJ, Stevens LA. Serum cystatin C as a marker of glomerular filtration rate. *Curr Opin Nephrol Hypertens* 2006;15:610-6.

13111 *Madero M, Wassel CL, Peralta CA, Najjar SS, Sutton-Tyrrell K, Fried L, et al. Cystatin C associates with arterial stiffness in older adults. *Journal of the American Society of Nephrology* 2009;20:1086-1093.

14077 *Maeda Y, Araki Y, Uno T, Kuyama T, Nishigaki K. Estimated glomerular filtration rate in peritoneal dialysis practice. *Clinical and Experimental Nephrology* 2011;15:611-612.

1373 *Mafham MM, Niculescu-Duvaz I, Barron J, Emberson JR, Dockrell ME, Landray MJ, et al. A practical method of measuring glomerular filtration rate by iohexol clearance using dried capillary blood spots. *Nephron Clin Pract* 2007;106:c104-12.

13112 *Mahon P, Shorten G. Perioperative acute renal failure. *Curr Opin Anaesthesiol*. 2006;19:332-8.

- 2087 *Maillard N, Mariat C, Bonneau C, Mehdi M, Thibaudin L, Laporte S, et al. Cystatin C-based equations in renal transplantation: moving toward a better glomerular filtration rate prediction? *Transplantation* 2008;85:1855-8.
- 13397 *Maillard N, Mehdi M, Thibaudin L, Berthoux F, Alamartine E, Mariat C. Creatinine-based GFR predicting equations in renal transplantation: Reassessing the tubular secretion effect. *Nephrology Dialysis Transplantation* 2010;25:3076-3082.
- 13114 *Maisei AS. Cardiovascular and renal surrogate markers in the clinical management of hypertension. *Cardiovascular Drugs and Therapy* 2009;23:317-326.
- 3624 *Malamitsi-Puchner A, Briana DD, Kontara L, Boutsikou M, Baka S, Hassiakos D, et al. Serum cystatin C in pregnancies with normal and restricted fetal growth. *Reprod Sci* 2007;14:37-42.
- 6635 *Malyszko J, Bachorzewska-Gajewska H, Malyszko JS, Pawlak K, Dobrzycki S. Serum neutrophil gelatinase-associated lipocalin as a marker of renal function in hypertensive and normotensive patients with coronary artery disease. *Nephrology (Carlton)* 2008;13:153-6.
- 13115 *Manetti L, Genovesi M, Pardini E, Grasso L, Lupi I, Morselli LL, et al. Early effects of methylprednisolone infusion on serum cystatin C in patients with severe Graves' ophthalmopathy [2]. *Clinica Chimica Acta* 2005;356:227-228.
- 13116 *Mangge H, Liebmann P, Tanil H, Herrmann J, Wagner C, Gallistl S, et al. Cystatin C, an early indicator for incipient renal disease in rheumatoid arthritis. *Clin Chim Acta*. 2000;300:195-202.
- 13117 *Mann JFE. What's new in hypertension 2008? *Nephrology Dialysis Transplantation* 2009;24:38-42.
- 13578 *Maple-Brown LJ, Lawton PD, Hughes JT, Sharma SK, Jones GR, Ellis AG, et al. Study Protocol--accurate assessment of kidney function in Indigenous Australians: aims and methods of the eGFR study. *BMC Public Health*. 2010;10:80.
- 13637 *Mariat C, Alamartine E, Afiani A, Thibaudin L, Laurent B, Berthoux P, et al. Predicting glomerular filtration rate in kidney transplantation: Are the K/DOQI guidelines applicable? *American Journal of Transplantation* 2005;5:2698-2703.
- 13118 *Markou N, Kanakaki M. Sleep-disordered breathing and reductions in renal function: Are they associated in the elderly? *Sleep Medicine* 2008;9:598-600.
- 6666 *Marouf R, Mojiminiyi O, Abdella N, Kortom M, Al Wazzan H. Comparison of renal function markers in Kuwaiti patients with sickle cell disease. *J Clin Pathol* 2006;59:345-51.
- 2099 *Martini S, Prevot A, Mosig D, Werner D, van Melle G, Guignard JP. Glomerular filtration rate: measure creatinine and height rather than cystatin C! *Acta Paediatr* 2003;92:1052-7.
- 13119 *Massey D. Commentary: clinical diagnostic use of cystatin C. *J Clin Lab Anal*. 2004;18:55-60.
- 13120 *Matthews K. Cystatin-C-based estimation of GFR in patients with liver cirrhosis. *Nature Clinical Practice Nephrology* 2006;2:239.
- 6706 *Mauss S, Berger F, Kuschak D, Henke J, Hegener P, Wolf E, et al. Cystatin C as a marker of renal function is affected by HIV replication leading to an underestimation of kidney function in HIV patients. *Antivir Ther* 2008;13:1091-5.
- 13121 *Mauss S, Berger F, Schmutz G. Antiretroviral therapy with tenofovir is associated with mild renal dysfunction. *AIDS*. 2005;19:93-5.
- 13122 *Mazul-Sunko B, Zarkovic N, Vrkic N, Antoljak N, Bekavac Beslin M, Nikolic Heitzler V, et al. Proatrial natriuretic peptide (1-98), but not cystatin C, is predictive for occurrence of acute renal insufficiency in critically ill septic patients. *Nephron Clin Pract*. 2004;97:c103-7.

- 13123 *McCullough PA. Multimodality Prevention of Contrast-Induced Acute Kidney Injury. *American Journal of Kidney Diseases* 2008;51:169-172.
- 13124 *McCullough PA, Khan M, James J. Serum cystatin C in the estimation of glomerular filtration on chronic angiotensin-converting enzyme inhibitor therapy: An illustrative case report. *Journal of Clinical Hypertension* 2009;11:651-655.
- 13125 *McDougal WS. Early detection of acute renal failure by serum cystatin C. *J Urol.* 2005;174:1024-5.
- 687 *McManus D, Shlipak M, Ix JH, Ali S, Whooley MA. Association of cystatin C with poor exercise capacity and heart rate recovery: data from the heart and soul study. *Am J Kidney Dis* 2007;49:365-72.
- 13126 *McManus DD, Corteville DC, Shlipak MG, Whooley MA, Ix JH. Relation of kidney function and albuminuria with atrial fibrillation (from the Heart and Soul Study). *Am J Cardiol.* 2009;104:1551-5.
- 13127 *McMurray MD, Trivax JE, McCullough PA. Editorial: Serum cystatin C, renal filtration function, and left ventricular remodeling. *Circulation: Heart Failure* 2009;2:86-89.
- 3655 *McNamara NV, Chen R, Janu MR, Bwititi P, Car G, Seibel M. Early renal failure detection by cystatin C in Type 2 diabetes mellitus: varying patterns of renal analyte expression. *Pathology* 2009;41:269-75.
- 13128 *Meier P, Froidevaux C, Dayer E, Blanc E. Cystatin C concentration and glomerular filtration rate. *Lancet.* 2001;357:634-5.
- 14078 *Melgosa M, Cabrera M, Salas S, Espinosa L, Quero J, Navarro M. Cystatin C values during the first year of life of preterms with birth weight less than 1500 grams. *Pediatric Nephrology* 2011;26:1355.
- 6742 *Mendiluce A, Bustamante J, Martin D, Santos M, Bustamante R, Pascual P, et al. Cystatin C as a marker of renal function in kidney transplant patients. *Transplant Proc* 2005;37:3844-7.
- 13129 *Menon V, Sarnak MJ, Shlipak MG. In response. *Annals of Internal Medicine* 2008;148:324.
- 6743 *Menon V, Shlipak MG, Wang X, Coresh J, Greene T, Stevens L, et al. Cystatin C as a risk factor for outcomes in chronic kidney disease. *Ann Intern Med* 2007;147:19-27.
- 13130 *Mercieri M, Mercieri A, Tritapepe L, Ruggeri M, Arcioni R, Repetto M, et al. High-dose aprotinin with gentamicin-vancomycin antibiotic prophylaxis increases blood concentrations of creatinine and cystatin C in patients undergoing coronary artery bypass grafting. *Br J Anaesth.* 1999;82:531-6.
- 13575 *Metzger J, Kirsch T, Schiffer E, Ulger P, Menten E, Brand K, et al. Urinary excretion of twenty peptides forms an early and accurate diagnostic pattern of acute kidney injury. *Kidney Int.* 2010;78:1252-62.
- 13132 *Middleton RJ, Foley RN, Hegarty J, Cheung CM, McElduff P, Gibson JM, et al. Reply [8]. *Nephrology Dialysis Transplantation* 2006;21:1733-1734.
- 3668 *Miklovicova D, Cervenova O, Cernianska A, Jancovicova Z, Dedik L, Vasilenkova A. Long-term follow-up of renal function in patients after surgery for obstructive uropathy. *Pediatr Nephrol* 2008;23:937-45.
- 13133 *Miller WG. Estimating glomerular filtration rate. *Clin Chem Lab Med.* 2009;47:1017-9.
- 13134 *Miller WG. Estimating equations for glomerular filtration rate in children: Laboratory considerations. *Clinical Chemistry* 2009;55:402-403.
- 6780 *Mingels A, Jacobs L, Kleijnen V, Wodzig W, Dieijen-Visser M. Cystatin C a marker for renal function after exercise. *Int J Sports Med* 2009;30:668-71.
- 13135 *Mingels AMA, Jacobs LHJ, Michielsen ECHJ, Swaanenburg JCJM, Wodzig WKWH, Van Dieijen-Visser MP. Cardiac and renal markers: Reference population versus marathon runners. *Nederlands Tijdschrift voor Klinische Chemie en Laboratoriumgeneeskunde* 2008;33:188-190.
- 13136 *Mitsnefes M, Kimbal T, Kartal J, Kathman T, Mishra J, Devarajan P. Serum cystatin C and left ventricular diastolic dysfunction in children with chronic kidney disease. *Pediatr Nephrol.* 2006;21:1293-8.

- 13137 *Mitsnefes MM, Khoury PR, Devarajan P. Beware of subgroup analysis. *Pediatric Nephrology* 2008;23:1191-1192.
- 13138 *Miyagawa Y, Takemura N, Hirose H. Evaluation of the measurement of serum cystatin C by an enzyme-linked immunosorbent assay for humans as a marker of the glomerular filtration rate in dogs. *J Vet Med Sci.* 2009;71:1169-76.
- 1402 *Mojiminiyi OA, Abdella N. Evaluation of cystatin C and beta-2 microglobulin as markers of renal function in patients with type 2 diabetes mellitus. *J Diabetes Complications* 2003;17:160-8.
- 13139 *Mojiminiyi OA, Abdella N, George S. Evaluation of serum cystatin C and chromogranin A as markers of nephropathy in patients with type 2 diabetes mellitus. *Scand J Clin Lab Invest.* 2000;60:483-9.
- 13140 *Mojiminiyi OA, Abdella NA, Zaki MY, El Gebely SA, Mohamedi HM, Aldhahi WA. Prevalence and associations of low plasma erythropoietin in patients with Type 2 diabetes mellitus. *Diabet Med.* 2006;23:839-44.
- 13141 *Mojiminiyi OA, Al Shayji I, George S. Serum cystatin C could replace serum creatinine as an indicator of glomerular filtration rate in routine practice. *European Journal of Laboratory Medicine* 1998;6:98-101.
- 13142 *Mojiminiyi OA, George S. Serum chromogranin A and its relationship to endogenous markers of glomerular filtration rate. *Ann Clin Biochem.* 1999;36 (Pt 2):247-9.
- 6810 *Mojiminiyi OA, Marouf R, Abdella N, Kortom M, Abdul-Razzak R. Serum concentration of cystatin C is not affected by cellular proliferation in patients with proliferative haematological disorders. *Ann Clin Biochem* 2002;39:308-10.
- 6815 *Momeni M, Baele P, Jacquet L, Mourad M, Waterloos H, Wallemacq P. Cystatin C in cardiac surgery. *Acta Anaesthesiol Belg* 2007;58:107-12.
- 140 *Monteiro MC, Alonso G, Ajzen H, Pereira AB. Assessment of glomerular filtration rate utilizing subcutaneously injected 51Cr-EDTA. *Braz J Med Biol Res* 1994;27:2557-64.
- 2124 *Montini G, Amici G, Milan S, Mussap M, Naturale M, Ratsch IM, et al. Middle molecule and small protein removal in children on peritoneal dialysis. *Kidney Int* 2002;61:1153-9.
- 13143 *Moodley J, Gangaram R, Khanyile R, Ojwang PJ. Serum cystatin C for assessment of glomerular filtration rate in hypertensive disorders of pregnancy. *Hypertens Pregnancy.* 2004;23:309-17.
- 13144 *Moran A, Katz R, Jenny NS, Astor B, Bluemke DA, Lima JA, et al. Left ventricular hypertrophy in mild and moderate reduction in kidney function determined using cardiac magnetic resonance imaging and cystatin C: the multi-ethnic study of atherosclerosis (MESA). *Am J Kidney Dis.* 2008;52:839-48.
- 6836 *Morgan C, Senthilselvan A, Bamforth F, Hoskinson M, Gowrishankar M. Correlation between cystatin C- and renal scan-determined glomerular filtration rate in children with spina bifida. *Pediatr Nephrol* 2008;23:329-32.
- 13145 *Mouton R, Holder K. Laboratory tests of renal function. *Anaesthesia and Intensive Care Medicine* 2006;7:240-243.
- 6865 *Mueller TF, Raeder J, Oettl K, Zitta S, Klausmann G, Estelberger W, et al. Cystatin C does not detect acute changes in glomerular filtration rate in early diabetic nephropathy. *Ren Fail* 2008;30:21-9.
- 13147 *Mulay A, Biyani M, Akbari A. Cystatin C and residual renal function in patients on peritoneal dialysis. *Am J Kidney Dis.* 2008;52:194-5; author reply 195-6.
- 13148 *Muller F, Bernard MA, Benkirane A, Ngo S, Lortat-Jacob S, Oury JF, et al. Fetal urine cystatin C as a predictor of postnatal renal function in bilateral uropathies. *Clin Chem.* 1999;45:2292-3.
- 6876 *Muntner P, Vupputuri S, Coresh J, Uribarri J, Fox CS. Metabolic abnormalities are present in adults with elevated serum cystatin C. *Kidney Int* 2009;76:81-8.

- 13149 *Murray AM. Editorial: The brain and the kidney connection: A model of accelerated vascular cognitive impairment. *Neurology* 2009;73:916-917.
- 144 *Mussap M, Dalla Vestra M, Fioretto P, Saller A, Varagnolo M, Nosadini R, et al. Cystatin C is a more sensitive marker than creatinine for the estimation of GFR in type 2 diabetic patients. *Kidney Int* 2002;61:1453-61.
- 13150 *Mussap M, Fanos V, Pizzini C, Marcolongo A, Chiaffoni G, Plebani M. Predictive value of amniotic fluid cystatin C levels for the early identification of fetuses with obstructive uropathies. *BJOG*. 2002;109:778-83.
- 14079 *Nair DR, Mehta S, Mikhailidis DP. Assessing renal function - Searching for the perfect marker continues! *Archives of Medical Science* 2011;7:565-567.
- 1422 *Nakai K, Kikuchi M, Fujimoto K, Kaneko Y, Omori S, Suwabe A. Serum levels of cystatin C in patients with malignancy. *Clin Exp Nephrol* 2008;12:132-9.
- 13151 *Nakamura T, Ioroi T, Sakaeda T, Horinouchi M, Hayashi N, Saito K, et al. Serum cystatin C levels to predict serum concentration of digoxin in Japanese patients. *Int J Med Sci*. 2006;3:92-6.
- 13152 *Naour N, Fellahi S, Renucci JF, Poitou C, Rouault C, Basdevant A, et al. Potential contribution of adipose tissue to elevated serum cystatin C in human obesity. *Obesity (Silver Spring)*. 2009;17:2121-6.
- 6933 *Naruse H, Ishii J, Kawai T, Hattori K, Ishikawa M, Okumura M, et al. Cystatin C in acute heart failure without advanced renal impairment. *Am J Med* 2009;122:566-73.
- 6936 *Narvaez-Sanchez R, Gonzalez L, Salamanca A, Silva M, Rios D, Arevalo S, et al. Cystatin C could be a replacement to serum creatinine for diagnosing and monitoring kidney function in children. *Clin Biochem* 2008;41:498-503.
- 13153 *Naud JF, Leblanc M. Biomarkers in acute kidney injury. *Biomarker Insights* 2008;2008:115-125.
- 13154 *Nelson CL, Karschimkus CS, Dragicevic G, Packham DK, Wilson AM, O'Neal D, et al. Systemic and vascular inflammation is elevated in early IgA and type 1 diabetic nephropathies and relates to vascular disease risk factors and renal function. *Nephrol Dial Transplant*. 2005;20:2420-6.
- 3715 *Newman DJ. Cystatin C. *Ann Clin Biochem* 2002;39:89-104.
- 13155 *Newman DJ. Cystatin C: what more do we need to know? *Nephron Clin Pract*. 2003;93:c122-3.
- 13156 *Newman DJ, Thakkar H, Edwards RG, Wilkie M, White T, Grubb AO, et al. Serum cystatin C: a replacement for creatinine as a biochemical marker of GFR. *Kidney Int Suppl*. 1994;47:S20-1.
- 13157 *Nguyen MT, Devarajan P. Biomarkers for the early detection of acute kidney injury. *Pediatr Nephrol*. 2008;23:2151-7.
- 13158 *Niccoli G, Conte M, Della Bona R, Altamura L, Siviglia M, Dato I, et al. Cystatin C is associated with an increased coronary atherosclerotic burden and a stable plaque phenotype in patients with ischemic heart disease and normal glomerular filtration rate. *Atherosclerosis*. 2008;198:373-80.
- 13159 *Nickolas TL, Barasch J, Devarajan P. Biomarkers in acute and chronic kidney disease. *Current Opinion in Nephrology and Hypertension* 2008;17:127-132.
- 13160 *Nielsen EI, Sandstrom M, Honore PH, Ewald U, Friberg LE. Developmental pharmacokinetics of gentamicin in preterm and term neonates: population modelling of a prospective study. *Clin Pharmacokinet*. 2009;48:253-63.
- 2144 *Nitta K, Hayashi T, Uchida K, Honda K, Tsukada M, Sekine S, et al. Serum cystatin C concentration as a marker of glomerular filtration rate in patients with various renal diseases. *Intern Med* 2002;41:931-5.
- 13162 *Noble DW. Aminoglycosides--engineering important patient-centered outcomes with antibiotics. *Crit Care Med*. 2009;37:2855-6.

- 1442 *Norlund L, Fex G, Lanke J, Von Schenck H, Nilsson JE, Leksell H, et al. Reference intervals for the glomerular filtration rate and cell-proliferation markers: serum cystatin C and serum beta 2-microglobulin/cystatin C-ratio. *Scand J Clin Lab Invest* 1997;57:463-70.
- 13163 *Norlund L, Grubb A, Fex G, Leksell H, Nilsson JE, Schenck H, et al. The increase of plasma homocysteine concentrations with age is partly due to the deterioration of renal function as determined by plasma cystatin C. *Clin Chem Lab Med*. 1998;36:175-8.
- 13164 *Nunez JFM. The normal ageing kidney - Morphology and physiology. *Reviews in Clinical Gerontology* 2008;18:175-197.
- 12853 *Nyman U, Bjork J, Lindstrom V, Grubb A. The Lund-Malmo creatinine-based glomerular filtration rate prediction equation for adults also performs well in children. *Scand J Clin Lab Invest* 2008:1-9.
- 7012 *Nyman U, Grubb A, Sterner G, Bjork J. Different equations to combine creatinine and cystatin C to predict GFR. Arithmetic mean of existing equations performs as well as complex combinations. *Scand J Clin Lab Invest* 2009;69:619-27.
- 13165 *Oberbauer R, Nenov V, Weidekamm C, Haas M, Szekeres T, Mayer G. Reduction in mean glomerular pore size coincides with the development of large shunt pores in patients with diabetic nephropathy. *Experimental Nephrology* 2001;9:49-53.
- 13166 *Obrenovic R, Petrovic D, Majkic-Singh N, Trbojevic J, Stojimirovic B. Influence of proteinuria on cystatin C serum concentration in patients with primary glomerulonephritis. *Jugoslovenska Medicinska Biohemija* 2006;25:21-25.
- 13167 *Odden MC, Chertow GM, Fried LF, Newman AB, Connelly S, Angleman S, et al. Cystatin C and measures of physical function in elderly adults: the Health, Aging, and Body Composition (HABC) Study. *Am J Epidemiol*. 2006;164:1180-9.
- 13168 *Odden MC, Scherzer R, Bacchetti P, Szczech LA, Sidney S, Grunfeld C, et al. Cystatin C level as a marker of kidney function in human immunodeficiency virus infection: the FRAM study. *Arch Intern Med*. 2007;167:2213-9.
- 13169 *Ogawa Y, Goto T, Tamasawa N, Matsui J, Tando Y, Sugimoto K, et al. Serum cystatin C in diabetic patients. Not only an indicator for renal dysfunction in patients with overt nephropathy but also a predictor for cardiovascular events in patients without nephropathy. *Diabetes Res Clin Pract*. 2008;79:357-61.
- 13171 *O'Hare AM, Newman AB, Katz R, Fried LF, Stehman-Breen CO, Seliger SL, et al. Cystatin C and incident peripheral arterial disease events in the elderly: results from the Cardiovascular Health Study. *Arch Intern Med*. 2005;165:2666-70.
- 7041 *Okamoto G, Sakamoto T, Kimura M, Ukishima Y, Sonoda A, Mori N, et al. Serum cystatin C as a better marker of vancomycin clearance than serum creatinine in elderly patients. *Clin Biochem* 2007;40:485-90.
- 13618 *Okura T, Jotoku M, Irita J, Enomoto D, Nagao T, Desilva VR, et al. Association between cystatin C and inflammation in patients with essential hypertension. *Clin Exp Nephrol*. 2010;14:584-8.
- 13619 *Okura T, Kurata M, Irita J, Enomoto D, Jotoku M, Nagao T, et al. Renal resistance index is a marker of future renal dysfunction in patients with essential hypertension. *J Nephrol*. 2010;23:175-80.
- 13173 *Olivieri O, Bassi A, Pizzolo F, Tinazzi E, Corrocher R. Cystatin C versus creatinine in renovascular disease. *Clinical Chemistry* 2002;48:2256-2259.
- 13174 *Olof H, Peter B. *Lakartidningen* 2009;106:420-421.
- 13401 *Olsson CL, Rippe B, Bendz H. A comparison of plasma cystatin C and plasma creatinine for the screening of renal function in lithium-treated patients. *Clinical nephrology* 2010;74:132-140.
- 7069 *O'Riordan S, Ouldred E, Brice S, Jackson SH, Swift CG. Serum cystatin C is not a better marker of creatinine or digoxin clearance than serum creatinine. *Br J Clin Pharmacol* 2002;53:398-402.

1450 *O'Riordan SE, Webb MC, Stowe HJ, Simpson DE, Kandarpa M, Coakley AJ, et al. Cystatin C improves the detection of mild renal dysfunction in older patients. *Ann Clin Biochem* 2003;40:648-55.

1451 *Orlando R, Mussap M, Plebani M, Piccoli P, De Martin S, Floreani M, et al. Diagnostic value of plasma cystatin C as a glomerular filtration marker in decompensated liver cirrhosis. *Clin Chem* 2002;48:850-8.

13402 *Orskov B, Borresen ML, Feldt-Rasmussen B, Ostergaard O, Laursen I, Strandgaard S. Estimating glomerular filtration rate using the new CKD-EPI equation and other equations in patients with autosomal dominant polycystic kidney disease. *American journal of nephrology* 2010;31:53-57.

7080 *Ortiz F, Harmoinen A, Paavonen T, Koskinen P, Gronhagen-Riska C, Honkanen E. Is Cystatin C more sensitive than creatinine in detecting early chronic allograft nephropathy? *Clin Nephrol* 2008;70:18-25.

7089 *Otsuka T, Tanaka A, Suemaru K, Inoue T, Nishimiya T, Murase M, et al. Evaluation of the clinical application of cystatin C, a new marker of the glomerular filtration rate, for the initial dose-setting of arbekacin. *J Clin Pharm Ther* 2008;33:227-35.

7100 *Ozer BA, Dursun B, Baykal A, Gultekin M, Suleymanlar G. Can cystatin C be a better marker for the early detection of renal damage in primary hypertensive patients? *Ren Fail* 2005;27:247-53.

13177 *Page MK, Bukki J, Luppa P, Neumeier D. Clinical value of cystatin C determination. *Clin Chim Acta*. 2000;297:67-72.

7114 *Palatini P, Benetti E, Zanier A, Santonastaso M, Mazzer A, Cozzio S, et al. Cystatin C as Predictor of Microalbuminuria in the Early Stage of Hypertension. *Nephron Clin Pract* 2009;113:c309-c314.

14080 *Papandreou C, Poulaki E, Pringos N, Benou N, Lykoka E, Piperaki K, et al. Cystatin C as a biomarker for glomerular filtration rate in cancer patients. *Clinical Chemistry and Laboratory Medicine* 2011;49:S263.

13620 *Papassotiriou I, Margeli A, Hantzi E, Delaporta P, Sergouniotti A, Goussetis E, et al. Cystatin C levels in patients with (beta)-thalassemia during deferasirox treatment. *Blood Cells, Molecules, and Diseases* 2010;44:152-155.

14081 *Pappy R, Abu-Fadel MS. Additional data supporting the use of statins to prevent contrast induced nephropathy. *Int J Cardiol* 2011;152:265-266.

13178 *Parikh CR, Devarajan P. New biomarkers of acute kidney injury. *Critical Care Medicine* 2008;36:S159-S165.

758 *Parikh NI, Hwang SJ, Yang Q, Larson MG, Guo CY, Robins SJ, et al. Clinical correlates and heritability of cystatin C (from the Framingham Offspring Study). *Am J Cardiol* 2008;102:1194-8.

13181 *Paskalev E, Lambreva L, Simeonov P, Koicheva N, Beleva B, Genova M, et al. Serum cystatin C in renal transplant patients. *Clin Chim Acta*. 2001;310:53-6.

13182 *Patel PC, Ayers CR, Murphy SA, Peshock R, Khera A, de Lemos JA, et al. Association of cystatin C with left ventricular structure and function: the Dallas Heart Study. *Circ Heart Fail*. 2009;2:98-104.

13186 *Peiris H, Chandrasena LG, Lanerolle RD. Serum cystatin C as a marker to identify patients with moderately impaired renal function. *Indian Journal of Clinical Biochemistry* 2008;23:163-166.

13188 *Peralta CA, Katz R, Madero M, Sarnak M, Kramer H, Criqui MH, et al. The differential association of kidney dysfunction with small and large arterial elasticity: the multiethnic study of atherosclerosis. *Am J Epidemiol*. 2009;169:740-8.

13565 *Peralta CA, Katz R, Sarnak MJ, Ix J, Fried LF, De Boer I, et al. Cystatin C identifies chronic kidney disease patients at higher risk for complications. *J Am Soc Nephrol* 2011;22:147-55.

13692 *Peralta CA, Shlipak MG, Judd S, Cushman M, McClellan W, Zakai NA, et al. Detection of Chronic Kidney Disease With Creatinine, Cystatin C, and Urine Albumin-to-Creatinine Ratio and Association With Progression to End-Stage Renal Disease and Mortality. *JAMA* 2011;305:1545-1552.

- 13189 *Peralta CA, Whooley MA, Ix JH, Shlipak MG. Kidney function and systolic blood pressure new insights from cystatin C: data from the Heart and Soul Study. *Am J Hypertens*. 2006;19:939-46.
- 13191 *Perianayagam MC, Seabra VF, Tighiouart H, Liangos O, Jaber BL. Serum cystatin C for prediction of dialysis requirement or death in acute kidney injury: a comparative study. *Am J Kidney Dis*. 2009;54:1025-33.
- 13193 *Perkins BA, Nelson RG, Krolewski AS. Cystatin C and the risk of death. *N Engl J Med*. 2005;353:842-4; author reply 842-4.
- 7175 *Perkins BA, Nelson RG, Ostrander BE, Blouch KL, Krolewski AS, Myers BD, et al. Detection of renal function decline in patients with diabetes and normal or elevated GFR by serial measurements of serum cystatin C concentration: results of a 4-year follow-up study. *J Am Soc Nephrol* 2005;16:1404-12.
- 7177 *Perlemoine C, Beauvieux MC, Rigalleau V, Baillet L, Barthes N, Derache P, et al. Interest of cystatin C in screening diabetic patients for early impairment of renal function. *Metabolism* 2003;52:1258-64.
- 7198 *Pham-Huy A, Leonard M, Lepage N, Halton J, Filler G. Measuring glomerular filtration rate with cystatin C and beta-trace protein in children with spina bifida. *J Urol* 2003;169:2312-5.
- 13198 *Pinto SWL, Sesso R, Vasconcelos E, Watanabe YJ, Pansute AM. Cystatin C is not more sensitive than creatinine for detecting early renal impairment in patients with diabetes. *American Journal of Kidney Diseases* 2001;38:310-316.
- 13199 *Piwowar A, Knapik-Kordecka M, Buczynska H, Warwas M. Plasma cystatin C concentration in non-insulin-dependent diabetes mellitus: relation with nephropathy. *Arch Immunol Ther Exp (Warsz)*. 1999;47:327-31.
- 7219 *Pizzarelli F, Lauretani F, Bandinelli S, Windham GB, Corsi AM, Giannelli SV, et al. Predictivity of survival according to different equations for estimating renal function in community-dwelling elderly subjects. *Nephrol Dial Transplant* 2009;24:1197-205.
- 2186 *Plebani M, Dall'Amico R, Mussap M, Montini G, Ruzzante N, Marsilio R, et al. Is serum cystatin C a sensitive marker of glomerular filtration rate (GFR)? A preliminary study on renal transplant patients. *Ren Fail* 1998;20:303-9.
- 7226 *Podracka L, Feber J, Lepage N, Filler G. Intra-individual variation of cystatin C and creatinine in pediatric solid organ transplant recipients. *Pediatr Transplant* 2005;9:28-32.
- 2190 *Poge U, Gerhardt T, Stoffel-Wagner B, Klehr HU, Sauerbruch T, Woitas RP. Calculation of glomerular filtration rate based on cystatin C in cirrhotic patients. *Nephrol Dial Transplant* 2006;21:660-4.
- 7228 *Poge U, Gerhardt T, Stoffel-Wagner B, Palmedo H, Klehr HU, Sauerbruch T, et al. Cystatin C-based calculation of glomerular filtration rate in kidney transplant recipients. *Kidney Int* 2006;70:204-10.
- 7229 *Poge U, Gerhardt T, Stoffel-Wagner B, Palmedo H, Klehr HU, Sauerbruch T, et al. Prediction of glomerular filtration rate in renal transplant recipients: cystatin C or modification of diet in renal disease equation? *Clin Transplant* 2006;20:200-5.
- 7234 *Poge U, Stoschus B, Stoffel-Wagner B, Gerhardt T, Klehr HU, Sauerbruch T, et al. Cystatin C as an endogenous marker of glomerular filtration rate in renal transplant patients. *Kidney Blood Press Res* 2003;26:55-60.
- 7243 *Poletti PA, Saudan P, Platon A, Mermillod B, Sautter AM, Vermeulen B, et al. I.v. N-acetylcysteine and emergency CT: use of serum creatinine and cystatin C as markers of radiocontrast nephrotoxicity. *AJR Am J Roentgenol* 2007;189:687-92.
- 13201 *Poniatowski B, Malyszko J, Bachorzewska-Gajewska H, Malyszko JS, Dobrzycki S. Serum neutrophil gelatinase-associated lipocalin as a marker of renal function in patients with chronic heart failure and coronary artery disease. *Kidney Blood Press Res*. 2009;32:77-80.

- 7285 *Premaratne E, MacIsaac RJ, Finch S, Panagiotopoulos S, Ekinci E, Jerums G. Serial measurements of cystatin C are more accurate than creatinine-based methods in detecting declining renal function in type 1 diabetes. *Diabetes Care* 2008;31:971-3.
- 7301 *Pucci L, Triscornia S, Lucchesi D, Fotino C, Pellegrini G, Pardini E, et al. Cystatin C and estimates of renal function: searching for a better measure of kidney function in diabetic patients. *Clin Chem* 2007;53:480-8.
- 13200 *Pöge U, Gerhardt T, Bokenkamp A, Stoffel-Wagner B, Klehr HU, Sauerbruch T, et al. Time course of low molecular weight proteins in the early kidney transplantation period--influence of corticosteroids. *Nephrol Dial Transplant*. 2004;19:2858-63.
- 13576 *Qutb A, Syed G, Tamim HM, Al Jondeby M, Jaradat M, Tamimi W, et al. Cystatin C-based formula is superior to MDRD, Cockcroft-Gault and Nankivell formulae in estimating the glomerular filtration rate in renal allografts. *Exp Clin Transplant*. 2009;7:197-202.
- 7338 *Ramel A, Jonsson PV, Björnsson S, Thorsdóttir I. Differences in the glomerular filtration rate calculated by two creatinine-based and three cystatin-C-based formulae in hospitalized elderly patients. *Nephron Clin Pract* 2008;108:c16-22.
- 13206 *Reese PP, Feldman HI. More evidence that cystatin C predicts mortality better than creatinine. *J Am Soc Nephrol*. 2009;20:2088-90.
- 13207 *Rehman T, Fought J, Solomon R. N-acetylcysteine effect on serum creatinine and cystatin C levels in CKD patients. *Clin J Am Soc Nephrol*. 2008;3:1610-4.
- 13208 *Reinhard M, Erlandsen EJ, Randers E. Biological variation of cystatin C and creatinine. *Scand J Clin Lab Invest*. 2009;69:831-6.
- 13209 *Ricci Z, Ronco C. Kidney diseases beyond nephrology: Intensive care. *Nephrology Dialysis Transplantation* 2008;23:820-826.
- 13210 *Ricci Z, Ronco C. Today's approach to the critically ill patient with acute kidney injury. *Blood Purification* 2009;27:127-134.
- 13211 *Rickli H, Benou K, Ammann P, Fehr T, Brunner-La Rocca HP, Petridis H, et al. Time course of serial cystatin C levels in comparison with serum creatinine after application of radiocontrast media. *Clin Nephrol*. 2004;61:98-102.
- 13212 *Rifkin DE, Katz R, Chonchol M, Fried LF, Cao J, de Boer IH, et al. Albuminuria, impaired kidney function and cardiovascular outcomes or mortality in the elderly. *Nephrol Dial Transplant*. 2009.
- 13213 *Rifkin DE, Shlipak MG, Katz R, Fried LF, Siscovick D, Chonchol M, et al. Rapid kidney function decline and mortality risk in older adults. *Arch Intern Med*. 2008;168:2212-8.
- 13214 *Rigalleau V, Beauvieux MC, Lasseur C, Chauveau P, Raffaitin C, Perlemoine C, et al. The combination of cystatin C and serum creatinine improves the monitoring of kidney function in patients with diabetes and chronic kidney disease. *Clin Chem*. 2007;53:1988-9.
- 7414 *Rigalleau V, Beauvieux MC, Le Moigne F, Lasseur C, Chauveau P, Raffaitin C, et al. Cystatin C improves the diagnosis and stratification of chronic kidney disease, and the estimation of glomerular filtration rate in diabetes. *Diabetes Metab* 2008;34:482-9.
- 13215 *Rigalleau V, Raffaitin C, Gin H, Beauvieux MC, Le Moigne F, Lasseur C, et al. Accounting for Body Composition does not Improve Cystatin C Estimation of GFR in Diabetic Subjects with CKD. *American Journal of Kidney Diseases* 2007;49:560.
- 7425 *Risch L, Blumberg A, Huber A. Rapid and accurate assessment of glomerular filtration rate in patients with renal transplants using serum cystatin C. *Nephrol Dial Transplant* 1999;14:1991-6.
- 7426 *Risch L, Blumberg A, Huber AR. Assessment of renal function in renal transplant patients using cystatin C. A comparison to other renal function markers and estimates. *Ren Fail* 2001;23:439-48.

- 13216 *Risch L, Huber AR. Glucocorticoids and increased serum cystatin C concentrations. *Clinica Chimica Acta* 2002;320:133-134.
- 7428 *Risch L, Huber AR. Assessing glomerular filtration rate in renal transplant recipients by estimates derived from serum measurements of creatinine and cystatin C. *Clin Chim Acta* 2005;356:204-11.
- 13218 *Risch L, Saely C, Reist U, Reist K, Hefti M, Huber AR. Course of glomerular filtration rate markers in patients receiving high-dose glucocorticoids following subarachnoidal hemorrhage [3]. *Clinica Chimica Acta* 2005;360:205-207.
- 13621 *Ristikankare A, Poyhia R, Kuitunen A, Skrifvars M, Hammainen P, Salmenpera M, et al. Serum cystatin C in elderly cardiac surgery patients. *Ann Thorac Surg.* 2010;89:689-94.
- 13813 *Ristiniemi N, Savage C, Bruun L, Pettersson K, Lilja H, Christensson A. Evaluation of a new immunoassay for cystatin C, based on a double monoclonal principle, in men with normal and impaired renal function. *Nephrol Dial Transplant* 2011.
- 7452 *Rodondi N, Yerly P, Gabriel A, Riesen WF, Burnier M, Paccaud F, et al. Microalbuminuria, but not cystatin C, is associated with carotid atherosclerosis in middle-aged adults. *Nephrol Dial Transplant* 2007;22:1107-14.
- 203 *Rodrigo E, de Francisco AL, Escallada R, Ruiz JC, Fresnedo GF, Pinera C, et al. Measurement of renal function in pre-ESRD patients. *Kidney Int Suppl* 2002:11-7.
- 13663 *Rognant N, Bacchetta J, Dubourg L, Ahmed SN, Radenne S, Dumortier J, et al. What is the best alternative to inulin clearance to estimate GFR in patients with decompensated alcoholic cirrhosis? *Nephrol Dial Transplant.* 2010;25:3569-75.
- 13542 *Rombach SM, Baas MC, ten Berge IJ, Krediet RT, Bemelman FJ, Hollak CE. The value of estimated GFR in comparison to measured GFR for the assessment of renal function in adult patients with Fabry disease. *Nephrol Dial Transplant* 2010;25:2549-56.
- 2244 *Roos JF, Doust J, Tett SE, Kirkpatrick CM. Diagnostic accuracy of cystatin C compared to serum creatinine for the estimation of renal dysfunction in adults and children--a meta-analysis. *Clin Biochem* 2007;40:383-91.
- 13224 *Ros S, Bajo A, del Peso G, Garcia de Miguel A, Santacruz S, Fernandez E, et al. Cystatin C as marker of residual renal function in patients on peritoneal dialysis: relation with parameters of peritoneal function. *J Nephrol.* 2007;20:468-73.
- 13225 *Rosner MH. Urinary biomarkers as a diagnostic tool for acute kidney injury. *Biomarkers in Medicine* 2008;2:55-66.
- 13226 *Royackers AANM, van Suijlen JDE, Hofstra LS, Kuiper MA, Bouman CSC, Spronk PE, et al. Serum cystatin C-A useful endogenous marker of renal function in intensive care unit patients at risk for or with acute renal failure? *Current Medicinal Chemistry* 2007;14:2314-2317.
- 7537 *Rule AD. Understanding estimated glomerular filtration rate: implications for identifying chronic kidney disease. *Curr Opin Nephrol Hypertens* 2007;16:242-9.
- 7539 *Rule AD, Bergstralh EJ, Slezak JM, Bergert J, Larson TS. Glomerular filtration rate estimated by cystatin C among different clinical presentations. *Kidney Int* 2006;69:399-405.
- 10 *Saeed A, Svensson MK, Herlitz H, Samuelsson O. [Determination of cystatin C. Not a reliable way to estimate renal function--and rather expensive!]. *Lakartidningen.* 2008;105:1588-9.
- 13231 *Sambasivan AS, Lepage N, Filler G. Cystatin C inpatient variability in children with chronic kidney disease is less than serum creatinine. *Clin Chem.* 2005;51:2215-6.
- 7576 *Samyn M, Cheeseman P, Bevis L, Taylor R, Samaroo B, Buxton-Thomas M, et al. Cystatin C, an easy and reliable marker for assessment of renal dysfunction in children with liver disease and after liver transplantation. *Liver Transpl* 2005;11:344-9.

- 13232 *Sarasua SM, Mueller P, Kathman S, Campagna D, Uddin MS, White MC. Confirming the utility of four kidney biomarker tests in a longitudinal follow-up study. *Ren Fail.* 2003;25:797-817.
- 13233 *Sarkar PD, Rajeshwari G, Shivaprakash TM. Cystatin C - A novel marker of glomerular filtration: A review. *Indian Journal of Clinical Biochemistry* 2005;20:139-144.
- 13234 *Sarnak MJ, Katz R, Fried LF, Siscovick D, Kestenbaum B, Seliger S, et al. Cystatin C and aging success. *Arch Intern Med.* 2008;168:147-53.
- 13235 *Sarnak MJ, Katz R, Stehman-Breen CO, Fried LF, Jenny NS, Psaty BM, et al. Cystatin C concentration as a risk factor for heart failure in older adults. *Ann Intern Med.* 2005;142:497-505.
- 13622 *Sato H, Kuroda T, Tanabe N, Ajiro J, Wada Y, Murakami S, et al. Cystatin C is a sensitive marker for detecting a reduced glomerular filtration rate when assessing chronic kidney disease in patients with rheumatoid arthritis and secondary amyloidosis. *Scand J Rheumatol.* 2010;39:33-7.
- 7633 *Schmitt A, Gladiëff L, Lansiaux A, Bobin-Dubigeon C, Etienne-Grimaldi MC, Boisdron-Celle M, et al. A universal formula based on cystatin C to perform individual dosing of carboplatin in normal weight, underweight, and obese patients. *Clin Cancer Res* 2009;15:3633-9.
- 13241 *Schmitt R, Bachmann S. Impact of thyroid dysfunction on serum cystatin C [3]. *Kidney International* 2003;64:1139-1140.
- 2283 *Schuck O, Teplan V, Sibova J, Stolova M. Predicting the glomerular filtration rate from serum creatinine, serum cystatin C and the Cockcroft and Gault formula with regard to drug dosage adjustment. *Int J Clin Pharmacol Ther* 2004;42:93-7.
- 13615 *Schutzer KM, Svensson MK, Zetterstrand S, Eriksson UG, Wahlander K. Reversible elevations of serum creatinine levels but no effect on glomerular filtration during treatment with the direct thrombin inhibitor AZD0837. *Eur J Clin Pharmacol.* 2010;66:903-10.
- 1551 *Schück O, Teplan V, Jabor A, Stolova M, Skibova J. Glomerular filtration rate estimation in patients with advanced chronic renal insufficiency based on serum cystatin C levels. *Nephron Clin Pract* 2003;93:c146-51.
- 13248 *Seco ML, Rus A, Sierra M, Caballero M, Borque L. Determination of serum cystatin C in patients with essential hypertension. *Nephron.* 1999;81:446-7.
- 13795 *Segarra A, de la Torre J, Ramos N, Quiroz A, Garjau M, Torres I, et al. Assessing Glomerular Filtration Rate in Hospitalized Patients: A Comparison Between CKD-EPI and Four Cystatin C-Based Equations. *Clin J Am Soc Nephrol* 2011;6:2411-20.
- 13249 *Seliger SL, Longstreth WT, Jr., Katz R, Manolio T, Fried LF, Shlipak M, et al. Cystatin C and subclinical brain infarction. *J Am Soc Nephrol.* 2005;16:3721-7.
- 13250 *Seo YS, Jung ES, An H, Kim JH, Jung YK, Yim HJ, et al. Serum cystatin C level is a good prognostic marker in patients with cirrhotic ascites and normal serum creatinine levels. *Liver Int.* 2009;29:1521-7.
- 7711 *Seronie-Vivien S, Delanaye P, Pieroni L, Mariat C, Froissart M, Cristol JP, et al. Cystatin C: current position and future prospects. *Clin Chem Lab Med* 2008;46:1664-86.
- 7715 *Seronie-Vivien S, Toullec S, Malard L, Thomas F, Durrand V, Chatelut E. Contribution of the MDRD equation and of cystatin C for renal function estimates in cancer patients. *Med Oncol* 2006;23:63-73.
- 7716 *Servais A, Giral P, Bernard M, Bruckert E, Deray G, Isnard Bagnis C. Is serum cystatin-C a reliable marker for metabolic syndrome? *Am J Med* 2008;121:426-32.
- 13616 *Shankar A, Lee KE, Klein BE, Muntner P, Brazy PC, Cruickshanks KJ, et al. Estimating glomerular filtration rate in a population-based study. *Vasc Health Risk Manag.* 2010;6:619-27.
- 7729 *Sharma AP, Kathiravelu A, Nadarajah R, Yasin A, Filler G. Body mass does not have a clinically relevant effect on cystatin C eGFR in children. *Nephrol Dial Transplant* 2009;24:470-4.

- 13617 *Shastri S, Katz R, Shlipak MG, Kestenbaum B, Peralta CA, Kramer H, et al. Cystatin C and Albuminuria as Risk Factors for Development of CKD Stage 3: The Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Kidney Dis.* 2011.
- 13254 *Shimizu-Tokiwa A, Kobata M, Ito H, Kobayashi N, Shou I, Funabiki K, et al. Serum cystatin C is a more sensitive marker of glomerular function than serum creatinine. *Nephron.* 2002;92:224-6.
- 13255 *Shlipak MG, Katz R, Cushman M, Sarnak MJ, Stehman-Breen C, Psaty BM, et al. Cystatin-C and inflammatory markers in the ambulatory elderly. *Am J Med.* 2005;118:1416.
- 13256 *Shlipak MG, Katz R, Fried LF, Jenny NS, Stehman-Breen CO, Newman AB, et al. Cystatin-C and mortality in elderly persons with heart failure. *J Am Coll Cardiol.* 2005;45:268-71.
- 13257 *Shlipak MG, Katz R, Kestenbaum B, Fried LF, Newman AB, Siscovick DS, et al. Rate of kidney function decline in older adults: a comparison using creatinine and cystatin C. *Am J Nephrol.* 2009;30:171-8.
- 7758 *Shlipak MG, Katz R, Kestenbaum B, Fried LF, Siscovick D, Sarnak MJ. Clinical and subclinical cardiovascular disease and kidney function decline in the elderly. *Atherosclerosis* 2009;204:298-303.
- 13258 *Shlipak MG, Katz R, Kestenbaum B, Siscovick D, Fried L, Newman A, et al. Rapid decline of kidney function increases cardiovascular risk in the elderly. *J Am Soc Nephrol.* 2009;20:2625-30.
- 7759 *Shlipak MG, Katz R, Sarnak MJ, Fried LF, Newman AB, Stehman-Breen C, et al. Cystatin C and prognosis for cardiovascular and kidney outcomes in elderly persons without chronic kidney disease. *Ann Intern Med* 2006;145:237-46.
- 13260 *Shlipak MG, Sarnak MJ, Katz R, Fried LF, Seliger SL, Newman AB, et al. Cystatin C and the risk of death and cardiovascular events among elderly persons. *N Engl J Med.* 2005;352:2049-60.
- 13864 *Silveiro SP, Araujo GN, Ferreira MN, Souza FD, Yamaguchi HM, Camargo EG. Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) Equation Pronouncedly Underestimates Glomerular Filtration Rate in Type 2 Diabetes. *Diabetes Care* 2011.
- 7783 *Simonsen O, Grubb A, Thysell H. The blood serum concentration of cystatin C (gamma-trace) as a measure of the glomerular filtration rate. *Scand J Clin Lab Invest* 1985;45:97-101.
- 7785 *Singh D, Whooley MA, Ix JH, Ali S, Shlipak MG. Association of cystatin C and estimated GFR with inflammatory biomarkers: the Heart and Soul Study. *Nephrol Dial Transplant* 2007;22:1087-92.
- 7792 *Sjöström P, Tidman M, Jones I. Determination of the production rate and non-renal clearance of cystatin C and estimation of the glomerular filtration rate from the serum concentration of cystatin C in humans. *Scand J Clin Lab Invest* 2005;65:111-24.
- 7796 *Skarby T, Jonsson P, Hjorth L, Behrentz M, Bjork O, Forestier E, et al. High-dose methotrexate: on the relationship of methotrexate elimination time vs renal function and serum methotrexate levels in 1164 courses in 264 Swedish children with acute lymphoblastic leukaemia (ALL). *Cancer Chemother Pharmacol* 2003;51:311-20.
- 13265 *Smith GL, Lichtman JH, Bracken MB, Shlipak MG, Phillips CO, DiCapua P, et al. Renal impairment and outcomes in heart failure: systematic review and meta-analysis. *J Am Coll Cardiol.* 2006;47:1987-96.
- 7827 *Soares AA, Eyff TF, Campani RB, Ritter L, Camargo JL, Silveiro SP. Glomerular filtration rate measurement and prediction equations. *Clin Chem Lab Med* 2009;47:1023-32.
- 13268 *Solichova P, Novackova L, Ochmanova R, Stejskal D. Assessment of serum beta-trace protein (BTP) measurement in the prediction of glomerular filtration rate. Comparison with serum cystatin C. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub.* 2006;150:83-4.
- 13604 *Soulsby N, Greville H, Coulthard K, Doecke C. What is the best method for measuring renal function in adults and children with cystic fibrosis? *J Cyst Fibros.* 2010;9:124-9.

13605 *Spanaus KS, Kollerits B, Ritz E, Hersberger M, Kronenberg F, von Eckardstein A. Serum creatinine, cystatin C, and beta-trace protein in diagnostic staging and predicting progression of primary nondiabetic chronic kidney disease. *Clin Chem*. 2010;56:740-9.

13273 *Stabuc B, Vrhovec L, Stabuc-Silih M, Cizej TE. Improved prediction of decreased creatinine clearance by serum cystatin C: use in cancer patients before and during chemotherapy. *Clin Chem*. 2000;46:193-7.

13274 *Stefanowicz J, Kosiak M, Kosiak W, Korbus-Kosiak A, Sierota D, Owczuk R. Cystatin C and parenchymal thickness/kidney length ratio in Wilms tumor survivors. *Pediatr Blood Cancer*. 2010;54:41-6.

14082 *Stefanowicz J, Kosiak M, Romanowicz G, Owczuk R, Adamkiewicz-Drozynska E, Balcerska A. Glomerular filtration rate and prevalence of chronic kidney disease in Wilms' tumour survivors. *Pediatric Nephrology* 2011;26:759-766.

7874 *Stefanowicz J, Owczuk R, Sierota D, Kaczorowska-Hac B, Balcerska A. Does antineoplasm treatment decrease the glomerular filtration rate in children? *Kidney Blood Press Res* 2009;32:194-9.

7884 *Sterner G, Bjork J, Carlson J, Grubb A, Nyman U. Validation of a new plasma cystatin C-based formula and the Modification of Diet in Renal Disease creatinine-based formula for determination of glomerular filtration rate. *Scand J Urol Nephrol* 2009;43:242-9.

7890 *Stevens LA, Coresh J, Feldman HI, Greene T, Lash JP, Nelson RG, et al. Evaluation of the modification of diet in renal disease study equation in a large diverse population. *J Am Soc Nephrol* 2007;18:2749-57.

13276 *Stevens LA, Coresh J, Greene T, Levey AS. Assessing kidney function--measured and estimated glomerular filtration rate. *N Engl J Med*. 2006;354:2473-83.

7891 *Stevens LA, Coresh J, Schmid CH, Feldman HI, Froissart M, Kusek J, et al. Estimating GFR using serum cystatin C alone and in combination with serum creatinine: a pooled analysis of 3,418 individuals with CKD. *Am J Kidney Dis* 2008;51:395-406.

13668 *Stevens LA, Levey AS. Use of the MDRD study equation to estimate kidney function for drug dosing. *Clin Pharmacol Ther*. 2009;86:465-7.

7896 *Stevens LA, Schmid CH, Greene T, Li L, Beck GJ, Joffe MM, et al. Factors other than glomerular filtration rate affect serum cystatin C levels. *Kidney Int* 2009;75:652-60.

13351 *Stevens LA, Schmid CH, Zhang YL, Coresh J, Manzi J, Landis R, et al. Development and validation of GFR-estimating equations using diabetes, transplant and weight. *Nephrol Dial Transplant* 2010;25:449-57.

1587 *Stickle D, Cole B, Hock K, Hruska KA, Scott MG. Correlation of plasma concentrations of cystatin C and creatinine to inulin clearance in a pediatric population. *Clin Chem* 1998;44:1334-8.

14062 *Stojanoski S, Gjorceva DP, Gruev T, Ristevska-Miceva S, Ristevska N. Impact of thyroid dysfunction on serum Cystatin C, serum Creatinine and Glomerular Filtration Rate. *Macedonian Journal of Medical Sciences* 2011;4:25-30.

13279 *Stevens H, Wide-Svensson D, Grubb A. Serum cystatin C is a better marker for preeclampsia than serum creatinine or serum urate. *Scand J Clin Lab Invest*. 2001;61:575-80.

3977 *Stevens H, Wide-Svensson D, Torffvit O, Grubb A. Serum cystatin C for assessment of glomerular filtration rate in pregnant and non-pregnant women. Indications of altered filtration process in pregnancy. *Scand J Clin Lab Invest* 2002;62:141-7.

7941 *Sun X, Chen Y, Chen X, Wang J, Xi C, Lin S, et al. Change of glomerular filtration rate in healthy adults with aging. *Nephrology (Carlton)* 2009;14:506-13.

13280 *Sun Y, Jiang T, Zeng Z, Chen P. Performance evaluation of a particle-enhanced turbidimetric cystatin C assay using the Abbott Aeroset analyser and assessment of cystatin C-based equations for estimating glomerular filtration rate in chronic kidney disease. *Nephrol Dial Transplant*. 2009.

- 13606 *Suzuki A, Imanishi Y, Nakano S, Niwa T, Ohmori T, Shirai K, et al. Usefulness of serum cystatin C to determine the dose of vancomycin in critically ill patients. *J Pharm Pharmacol.* 2010;62:901-7.
- 13607 *Svenmarker S, Haggmark S, Holmgren A, Naslund U. Serum markers are not reliable measures of renal function in conjunction with cardiopulmonary bypass. *Interact Cardiovasc Thorac Surg.* 2011.
- 13282 *Szewczyk M, Wielkoszynski T, Zakliczynski M, Zembala M. Plasma neutrophil gelatinase-associated lipocalin (NGAL) correlations with cystatin C, serum creatinine, and glomerular filtration rate in patients after heart and lung transplantation. *Transplant Proc.* 2009;41:3242-3.
- 7976 *Taes YE, Wuyts B, Boelaert JR, De Vriese AS, Delanghe JR. Prohepcidin accumulates in renal insufficiency. *Clin Chem Lab Med* 2004;42:387-9.
- 13283 *Taglieri N, Fernandez-Berges DJ, Koenig W, Consuegra-Sanchez L, Fernandez JM, Robles NR, et al. Plasma cystatin C for prediction of 1-year cardiac events in Mediterranean patients with non-ST elevation acute coronary syndrome. *Atherosclerosis.* 2009.
- 1602 *Takuwa S, Ito Y, Ushijima K, Uchida K. Serum cystatin-C values in children by age and their fluctuation during dehydration. *Pediatr Int* 2002;44:28-31.
- 1604 *Tan GD, Lewis AV, James TJ, Altmann P, Taylor RP, Levy JC. Clinical usefulness of cystatin C for the estimation of glomerular filtration rate in type 1 diabetes: reproducibility and accuracy compared with standard measures and iohexol clearance. *Diabetes Care* 2002;25:2004-9.
- 13286 *Tanaka A, Suemaru K, Otsuka T, Ido K, Nishimiya T, Sakai I, et al. Estimation of the initial dose setting of vancomycin therapy with use of cystatin C as a new marker of renal function. *Ther Drug Monit.* 2007;29:261-4.
- 7991 *Tanaka A, Suemaru K, Otsuka T, Ido K, Nishimiya T, Sakai I, et al. Hoek's formula, a cystatin C-based prediction formula for determining the glomerular filtration rate, is the most effective method for original adjusting the dosage of vancomycin. *Int J Clin Pharmacol Ther* 2007;45:592-7.
- 13687 *Tangri N, Stevens LA, Griffith J, Tighiouart H, Djurdjev O, Naimark D, et al. A Predictive Model for Progression of Chronic Kidney Disease to Kidney Failure. *JAMA* 2011.
- 13608 *Tangri N, Stevens LA, Schmid CH, Zhang YL, Beck GJ, Greene T, et al. Changes in dietary protein intake has no effect on serum cystatin C levels independent of the glomerular filtration rate. *Kidney Int.* 2011;79:471-7.
- 13288 *Tarif N, Alwakeel JS, Mitwalli AH, Durdana H, Memon NA, Askar A, et al. Serum cystatin C as a marker of renal function in patients with acute renal failure. *Saudi J Kidney Dis Transpl.* 2008;19:918-23.
- 13666 *Tent H, Rook M, Stevens LA, Van Son WJ, Van Pelt LJ, Hofker HS, et al. Renal function equations before and after living kidney donation: A within-individual comparison of performance at different levels of renal function. *Clinical Journal of the American Society of Nephrology* 2010;5:1960-1968.
- 8036 *Thomas F, Seronie-Vivien S, Gladiéff L, Dalenc F, Durrand V, Malard L, et al. Cystatin C as a new covariate to predict renal elimination of drugs: application to carboplatin. *Clin Pharmacokinet* 2005;44:1305-16.
- 8043 *Thomassen SA, Johannesen IL, Erlandsen EJ, Abrahamsen J, Randers E. Serum cystatin C as a marker of the renal function in patients with spinal cord injury. *Spinal Cord* 2002;40:524-8.
- 3944 *Tian S, Kusano E, Ohara T, Tabei K, Itoh Y, Kawai T, et al. Cystatin C measurement and its practical use in patients with various renal diseases. *Clin Nephrol* 1997;48:104-8.
- 8059 *Tidman M, Sjostrom P, Jones I. A Comparison of GFR estimating formulae based upon s-cystatin C and s-creatinine and a combination of the two. *Nephrol Dial Transplant* 2008;23:154-60.
- 1616 *Toffaletti JG, McDonnell EH. Variation of serum creatinine, cystatin C, and creatinine clearance tests in persons with normal renal function. *Clin Chim Acta* 2008;395:115-9.

- 13293 *Tombach B, Bremer C, Reimer P, Kisters K, Schaefer RM, Geens V, et al. Renal tolerance of a neutral gadolinium chelate (gadobutrol) in patients with chronic renal failure: results of a randomized study. *Radiology*. 2001;218:651-7.
- 13295 *Tondel C, Ramaswami U, Aakre KM, Wijburg F, Bouwman M, Svarstad E. Monitoring renal function in children with Fabry disease: comparisons of measured and creatinine-based estimated glomerular filtration rate. *Nephrol Dial Transplant*. 2009.
- 13693 *Tonelli M, Manns B. Supplementing Creatinine-Based Estimates of Risk in Chronic Kidney Disease: Is It Time? *JAMA* 2011.
- 4197 *Torner A, Odar-Cederlof I, Kallner A, Akner G. Renal function in community-dwelling frail elderly. Comparison between measured and predicted glomerular filtration rate in the elderly and proposal for a new cystatin C-based prediction equation. *Aging Clin Exp Res* 2008;20:216-25.
- 8094 *Toyoda H, Kumada T, Kiriya S, Sone Y, Tanikawa M, Hisanaga Y, et al. Correlation of serum ribavirin concentration with pretreatment renal function estimates in patients with chronic hepatitis C receiving combination antiviral therapy with peginterferon and ribavirin. *J Viral Hepat* 2008;15:651-8.
- 903 *Treiber M, Pecovnik-Balon B, Gorenjak M. Cystatin C versus creatinine as a marker of glomerular filtration rate in the newborn. *Wien Klin Wochenschr* 2006;118 Suppl 2:66-70.
- 13669 *Tsai JP, Wu SW, Hung TW, Kao WT, Hong CL, Lian JD, et al. Diagnostic performance of serum cystatin C and serum creatinine in the prediction of chronic kidney disease in renal transplant recipients. *Transplant Proc*. 2010;42:4530-3.
- 13814 *Uemura O, Ushijima K, Nagai T, Yamada T, Yamakawa S, Hibi Y, et al. Measurements of serum cystatin C concentrations underestimate renal dysfunction in pediatric patients with chronic kidney disease. *Clin Exp Nephrol* 2011;15:535-8.
- 13302 *Ustundag Y, Samsar U, Acikgoz S, Cabuk M, Kiran S, Kulah E, et al. Analysis of glomerular filtration rate, serum cystatin C levels, and renal resistive index values in cirrhosis patients. *Clin Chem Lab Med*. 2007;45:890-4.
- 8163 *Uzun H, Ozmen Keles M, Ataman R, Aydin S, Kalender B, Uslu E, et al. Serum cystatin C level as a potentially good marker for impaired kidney function. *Clin Biochem* 2005;38:792-8.
- 14083 *Wagner D, Kniepeiss D, Stiegler P, Zitta S, Bradatsch A, Meinitzer A, et al. The assessment of GFR after orthotopic liver transplantation using cystatin c and creatinine-based equations. *Transplant International* 2011;24:203.
- 13305 *Walczak K, Rosniak-Bak K, Paradowski M, Moczulski D. Cystatin C as a marker of renal function in diabetic kidney disease. *Diabetologia Doswiadczalna i Kliniczna* 2009;9:69-72.
- 8219 *Van Den Noortgate NJ, Janssens WH, Delanghe JR, Afschrift MB, Lameire NH. Serum cystatin C concentration compared with other markers of glomerular filtration rate in the old old. *J Am Geriatr Soc* 2002;50:1278-82.
- 13609 *van der Velde M, Bakker SJ, de Jong PE, Gansevoort RT. Influence of age and measure of eGFR on the association between renal function and cardiovascular events. *Clin J Am Soc Nephrol*. 2010;5:2053-9.
- 13610 *van Deventer HE, Paiker JE, Katz IJ, George JA. A comparison of cystatin C- and creatinine-based prediction equations for the estimation of glomerular filtration rate in black South Africans. *Nephrol Dial Transplant*. 2011:1553-1558.
- 13433 *van Pottelbergh G, van Heden L, Mathei C, Degryse J. Methods to evaluate renal function in elderly patients: A systematic literature review. *Age and Ageing* 2010;39:542-548.
- 8238 *van Rossum LK, Zietse R, Vulto AG, de Rijke YB. Renal extraction of cystatin C vs 125I-iothalamate in hypertensive patients. *Nephrol Dial Transplant* 2006;21:1253-6.

13306 *Wang J, Sim AS, Wang XL, Salonikas C, Moriatis M, Naidoo D, et al. Relations between markers of renal function, coronary risk factors and the occurrence and severity of coronary artery disease. *Atherosclerosis*. 2008;197:853-9.

8251 *Wang QP, Gu JW, Zhan XH, Li H, Luo XH. Assessment of glomerular filtration rate by serum cystatin C in patients undergoing coronary artery bypass grafting. *Ann Clin Biochem* 2009;46:495-500.

13307 *Wasen E, Isoaho R, Mattila K, Vahlberg T, Kivela SL, Irjala K. Serum cystatin C in the aged: relationships with health status. *Am J Kidney Dis*. 2003;42:36-43.

8269 *Wasen E, Isoaho R, Mattila K, Vahlberg T, Kivela SL, Irjala K. Estimation of glomerular filtration rate in the elderly: a comparison of creatinine-based formulae with serum cystatin C. *J Intern Med* 2004;256:70-8.

8270 *Wasen E, Isoaho R, Vahlberg T, Kivela SL, Irjala K. Association between markers of renal function and C-reactive protein level in the elderly: confounding by functional status. *Scand J Clin Lab Invest* 2008;68:484-91.

8272 *Wasilewska A, Zoch-Zwierz W, Jadeszko I, Porowski T, Biernacka A, Niewiarowska A, et al. Assessment of serum cystatin C in children with congenital solitary kidney. *Pediatr Nephrol* 2006;21:688-93.

13312 *Weinert LS, Prates AB, do Amaral FB, Vaccaro MZ, Camargo JL, Silveiro SP. Gender does not influence cystatin C concentrations in healthy volunteers. *Clin Chem Lab Med*. 2009.

14084 *Weir MR, Mulgaonkar S, Chan L, Shidban H, Waid TH, Preston D, et al. Mycophenolate mofetil-based immunosuppression with sirolimus in renal transplantation: a randomized, controlled Spare-the-Nephron trial. *Kidney International* 2011;79:897-907.

13316 *Westphal S, Dierkes J, Luley C. Effects of fenofibrate and gemfibrozil on plasma homocysteine. *Lancet*. 2001;358:39-40.

13611 *Wetmore JB, Palsson R, Belmont JM, Sigurdsson G, Franzson L, Indridason OS. Discrepancies between creatinine- and cystatin C-based equations: implications for identification of chronic kidney disease in the general population. *Scand J Urol Nephrol*. 2010;44:242-50.

8333 *White C, Akbari A, Hussain N, Dinh L, Filler G, Lepage N, et al. Estimating glomerular filtration rate in kidney transplantation: a comparison between serum creatinine and cystatin C-based methods. *J Am Soc Nephrol* 2005;16:3763-70.

8334 *White C, Akbari A, Hussain N, Dinh L, Filler G, Lepage N, et al. Chronic kidney disease stage in renal transplantation classification using cystatin C and creatinine-based equations. *Nephrol Dial Transplant* 2007;22:3013-20.

13318 *White CA, Akbari A, Doucette S, Fergusson D, Knoll GA. Estimating Glomerular Filtration Rate in Kidney Transplantation: Is the New Chronic Kidney Disease Epidemiology Collaboration Equation Any Better? *Clin Chem*. 2009.

8337 *White CA, Akbari A, Doucette S, Fergusson D, Ramsay T, Hussain N, et al. Effect of Clinical Variables and Immunosuppression on Serum Cystatin C and Beta-Trace Protein in Kidney Transplant Recipients. *Am J Kidney Dis* 2009;54:922-30.

13320 *Viberg A, Lannergard A, Larsson A, Cars O, Karlsson MO, Sandstrom M. A population pharmacokinetic model for cefuroxime using cystatin C as a marker of renal function. *Br J Clin Pharmacol*. 2006;62:297-303.

8362 *Wielosz E, Dryglewska M, Majdan M. Antiphospholipid antibodies and kidney involvement in patients with systemic sclerosis. *Clin Rheumatol* 2009;28:955-9.

13322 *Villa P, Jimenez M, Soriano MC, Manzanares J, Casasnovas P. Serum cystatin C concentration as a marker of acute renal dysfunction in critically ill patients. *Crit Care*. 2005;9:R139-43.

8385 *Willems D, Wolff F, Mekhali F, Gillet C. Cystatin C for early detection of renal impairment in diabetes. *Clin Biochem* 2009;42:108-10.

2409 *Willems HL, Hilbrands LB, van de Calseyde JF, Monnens LA, Swinkels DW. Is serum cystatin C the marker of choice to predict glomerular filtration rate in paediatric patients? *Ann Clin Biochem* 2003;40:60-4.

8399 *Vinge E, Lindergard B, Nilsson-Ehle P, Grubb A. Relationships among serum cystatin C, serum creatinine, lean tissue mass and glomerular filtration rate in healthy adults. *Scand J Clin Lab Invest* 1999;59:587-92.

13323 *Winston JA. Estimating glomerular filtration rate in patients with HIV infection. *Semin Nephrol*. 2008;28:576-80.

13324 *Viollet L, Gailey S, Thornton DJ, Friedman NR, Flanigan KM, Mahan JD, et al. Utility of cystatin C to monitor renal function in Duchenne muscular dystrophy. *Muscle Nerve*. 2009;40:438-42.

8406 *Visvardis G, Griveas I, Zilidou R, Papadopoulou D, Mitsopoulos E, Kyriklidou P, et al. Glomerular filtration rate estimation in renal transplant patients based on serum cystatin-C levels: comparison with other markers of glomerular filtration rate. *Transplant Proc* 2004;36:1757-9.

1671 *Woitas RP, Stoffel-Wagner B, Flommersfeld S, Poege U, Schiedermaier P, Klehr HU, et al. Correlation of serum concentrations of cystatin C and creatinine to inulin clearance in liver cirrhosis. *Clin Chem* 2000;46:712-5.

13327 *Wong H, Vivian L, Weiler G, Filler G. Patients with autosomal dominant polycystic kidney disease hyperfiltrate early in their disease. *Am J Kidney Dis*. 2004;43:624-8.

13870 *Wuerzner G, Bochud M, Giusti V, Burnier M. Measurement of glomerular filtration rate in obese patients: pitfalls and potential consequences on drug therapy. *Obes Facts* 2011;4:238-43.

8435 *Vupputuri S, Fox CS, Coresh J, Woodward M, Muntner P. Differential estimation of CKD using creatinine-versus cystatin C-based estimating equations by category of body mass index. *Am J Kidney Dis* 2009;53:993-1001.

13328 *Vynckier L, Stove V, Delanghe JR. Macromolecular cystatin C can be a caveat for estimating glomerular filtration rate in biliary obstruction. *Clinical Chemistry* 2008;54:1257-1259.

8440 *Xia LH, Bing XG, An XT. Serum cystatin C assay for the detection of early renal impairment in diabetic patients. *J Clin Lab Anal* 2004;18:31-5.

13612 *Xirouchakis E, Marelli L, Cholongitas E, Manousou P, Calvaruso V, Pleguezuelo M, et al. Comparison of cystatin C and creatinine-based glomerular filtration rate formulas with ⁵¹Cr-EDTA clearance in patients with cirrhosis. *Clin J Am Soc Nephrol*. 2011;6:84-92.

8443 *Xu H, Lu Y, Teng D, Wang J, Wang L, Li Y. Assessment of glomerular filtration rate in renal transplant patients using serum cystatin C. *Transplant Proc* 2006;38:2006-8.

1681 *Xu X, Zou J, Ding X, Xin D, Ren Y. Clinical value of serum cystatin C by ELISA for estimation of glomerular filtration rate. *J Clin Lab Anal* 2004;18:61-4.

13882 *Yang Q, Li R, Zhong Z, Mao H, Fan J, Lin J, et al. Is cystatin C a better marker than creatinine for evaluating residual renal function in patients on continuous ambulatory peritoneal dialysis? *Nephrol Dial Transplant* 2011;26:3358-3365.

13329 *Yang YH, He XJ, Chen SR, Wang L, Li EM, Xu LY. Changes of serum and urine neutrophil gelatinase-associated lipocalin in type-2 diabetic patients with nephropathy: one year observational follow-up study. *Endocrine*. 2009;36:45-51.

1685 *Yang YS, Peng CH, Lin CK, Wang CP, Huang CN. Use of serum cystatin C to detect early decline of glomerular filtration rate in type 2 diabetes. *Intern Med* 2007;46:801-6.

13330 *Yashiro M, Kamata T, Segawa H, Kadoya Y, Murakami T, Muso E. Comparisons of cystatin C with creatinine for evaluation of renal function in chronic kidney disease. *Clin Exp Nephrol*. 2009;13:598-604.

13331 *Yeo Y, Han DJ, Moon DH, Park JS, Yang WS, Chang JW, et al. Suitability of the IDMS-Traceable MDRD Equation Method to Estimate GFR in Early Postoperative Renal Transplant Recipients. *Nephron Clin Pract*. 2010;114:108-117.

8467 *Ylinen EA, Ala-Houhala M, Harmoinen AP, Knip M. Cystatin C as a marker for glomerular filtration rate in pediatric patients. *Pediatr Nephrol* 1999;13:506-9.

13332 *Yokoyama H, Inoue T, Node K. Effect of insulin-unstimulated diabetic therapy with miglitol on serum cystatin C level and its clinical significance. *Diabetes Res Clin Pract*. 2009;83:77-82.

8478 *Young JA, Hwang SJ, Sarnak MJ, Hoffmann U, Massaro JM, Levy D, et al. Association of visceral and subcutaneous adiposity with kidney function. *Clin J Am Soc Nephrol* 2008;3:1786-91.

8495 *Zahran A, El-Husseini A, Shoker A. Can cystatin C replace creatinine to estimate glomerular filtration rate? A literature review. *Am J Nephrol* 2007;27:197-205.

2426 *Zahran A, Qureshi M, Shoker A. Comparison between creatinine and cystatin C-based GFR equations in renal transplantation. *Nephrol Dial Transplant* 2007;22:2659-68.

1691 *Zappitelli M, Parvex P, Joseph L, Paradis G, Grey V, Lau S, et al. Derivation and validation of cystatin C-based prediction equations for GFR in children. *Am J Kidney Dis* 2006;48:221-30.

13613 *Zhang PP, Zhan JF, Xie HL, Li LS, Liu ZH. Evaluation of glomerular filtration rate using cystatin C in diabetic patients analysed by multiple factors including tubular function. *J Int Med Res*. 2010;38:473-83.

13614 *Zhang QL, Brenner H, Koenig W, Rothenbacher D. Prognostic value of chronic kidney disease in patients with coronary heart disease: role of estimating equations. *Atherosclerosis*. 2010;211:342-7.

13335 *Zhu J, Yin R, Wu H, Yi J, Luo L, Dong G, et al. Cystatin C as a reliable marker of renal function following heart valve replacement surgery with cardiopulmonary bypass. *Clin Chim Acta*. 2006;374:116-21.