

**Bilaga 3 Exkluderade studier/Excluded studies**

Referens/Reference	Exklusionsorsak/ Reason for exclusion
A pelvic floor muscle exercise programme for urinary incontinence following childbirth. Nurs Stand. 2006 2006-01-01;20(33):46–50.	Duplikat/duplicat
Alexander J, Thomas PW, Sanghera J. Treatments for secondary postpartum haemorrhage. Cochrane Database Syst Rev. 2002 2002-01-01;0:N.PAG-N.PAG 1p.	Fel population/ wrong population
Amorim-Costa C, Mota R, Rebelo C, Silva PT. Uterine compression sutures for postpartum hemorrhage: is routine postoperative cavity evaluation needed? Acta Obstet Gynecol Scand.90(7):701–6.	Fel population/ wrong population
Badawy SZ, Etman A, Singh M, Murphy K, Mayelli T, Philadelphia M. Uterine artery embolization: the role in obstetrics and gynecology. Clinical Imaging;25:288–95.	Fel population/intervention
Bø K, Hilde G. Long term effect of pelvic floor muscle training for female stress and mixed urinary incontinence—a systematic review. Neurourol Urodyn. 2011 2011-01-01;30(6):1008–9.	Ingen systematisk översikt/No systematic review
Boyle R, Hay-Smith J, Mørkved S, Fairbrother KA, Herbison GP. Pelvic floor muscle training for prevention and treatment of urinary and faecal incontinence in antenatal and postnatal women. Cochrane Database Syst Rev 2012;0:N.PAG-N.PAG 1p.	Fel population/ wrong population
Cerruto MA, D'Elia C, Aloisi A, Fabrello M, Artibani W. Prevalence, incidence and obstetric care impact for women with urinary incontinence in Europe: a systematic and qualitative review of the literatur. Neurourol Urodyn. 2011 2011-01-01;30:2-3.	Ingen intervention/ no intervention
Chou D, Abalos E, Gyte GML, Gülmezoglu AM. Drugs for perineal pain in the early postpartum period. Cochrane Database Syst Rev. 2009 2009-01-01;0(3).	Protokoll/protocool
Dodd J, Dare MR, Middleton P. Treatment for women with postpartum iron deficiency anaemia. Cochrane Database Syst Rev (Online). 2004 2004-01-01;0(4).	Fel population/ wrong population.
Doumouchtsis S, Nikolopoulos K, Talaulikar V, Krishna A, Arulkumaran S. Menstrual and fertility outcomes following the surgical management of postpartum haemorrhage: a systematic review. BJOG. 2014 2014-01-01;121(4):382–8 7p.	Fel population/ wrong population.

Referens/Reference	Exklusionsorsak/ Reason for exclusion
Doumouchtsis SK, Papageorghiou AT, Arulkumaran S. Systematic review of conservative management of postpartum hemorrhage: What to do when medical treatment fails. <i>Obstet Gynecol Surv.</i> 2007 2007-01-01;62(8):540–7.	Fel population/ wrong population.
Draper J, Newell R. A discussion of some of the literature relating to history, repair and consequences of perineal trauma. <i>Midwifery.</i> 1996. 1996-01-01;12(3):140–5.	Not a systematic review/Ej en systematisk översikt
Dudding TC, Vaizey CJ, Kamm MA. Obstetric anal sphincter injury: incidence, risk factors, and management. <i>Ann Surg.</i> 247(2):224–37.	Ingen intervention/outcom e. No intervention or outcome.
Ekelund K, Hanke G, Stensballe J, Wikkelsoe A, Albrechtsen CK, Afshari A. Hemostatic resuscitation in postpartum hemorrhage – a supplement to surgery. <i>Acta Obstet Gynecol Scand.</i> 94(7):680–92.	Fel population/ wrong population.
Farrar D, Tuffnell DJ, Ramage C. Interventions for women in subsequent pregnancies following obstetric anal sphincter injury to reduce the risk of recurrent injury and associated harms. <i>The Cochrane Database Syst Rev.</i> 2014;11:CD010374.	Wrong intervention/fel intervention
Farrell SA. Overlapping compared with end-to-end repair of third and fourth degree obstetric anal sphincter tears. <i>Curr Opin Obstet Gynecol.</i> 2011 23(5):386–90.	Not a systematic review/Ej en systematisk översikt
Ferland RJ, Adler LM. Management of postpartum uterine atony. <i>R I Med J (Providence, RI).</i> 1990 1990-01-01;73(12):626–7.	Fel population/ wrong population.
Ferrer P, Roberts I, Sydenham E, Blackhall K, Shakur H. Anti-fibrinolytic agents in post partum haemorrhage: A systematic review. <i>BMC Pregnancy Childbirth.</i> 2009 2009-01-01;9.	Fel population/ wrong population.
Fotopoulou C, Dudenhausen JW. Uterine compression sutures for preserving fertility in severe postpartum haemorrhage: An overview 13 years after the first description. <i>J Obstet Gynaecol.</i> 2010. 2010-01-01;30(4):339–49.	Fel population/wrong intervention.
Franchini M, Franchi M, Bergamini V, Montagnana M, Salvagno GL, Targher G, et al. The use of recombinant activated FVII in postpartum hemorrhage. <i>Clin Obstet Gynecol.</i> 2010 2010-01-01;53(1):219–27.	Fel population/ wrong population.
Franchini M, Franchi M, Bergamini V, Salvagno GL, Montagnana M, Lippi G. A critical review on the use of recombinant factor VIIa	Fel population/ wrong population.

Referens/Reference	Exklusionsorsak/ Reason for exclusion
in life-threatening obstetric postpartum hemorrhage. <i>Semin Thromb Hemost.</i> 2008 34(1):104–12.	
French LM, Smaill FM. Antibiotic regimens for endometritis after delivery. <i>Cochrane Database Syst Rev (Online).</i> 2004 2004-01-01;0(4):CD001067.	Fel population/ wrong population.
Genadry RR, Creanga AA, Roenneburg ML, Wheelless CR. Complex obstetric fistulas. <i>Int J Gynaecol Obstet</i> ;99:S51–6.	Fel population/ wrong population.
Gracia V-D. The use of magnesium sulfate for women with severe preeclampsia or eclampsia diagnosed during the postpartum period. <i>Journal of Maternal-Fetal &amp; Neonatal Medicine</i> 2015;28:2207–09 3p.	Fel population/ wrong population.
Haddow G, Watts R, Robertson J. Effectiveness of a pelvic floor muscle exercise program on urinary incontinence following childbirth. <i>JB Library of Systematic Reviews</i> 2014;3:153–214.	Fel population/ wrong population.
Hartmann K, Viswanathan M, Palmieri R, Gartlehner G, Thorp J, Jr., Lohr KN. Clinical review. Outcomes of routine episiotomy: a systematic review. <i>JAMA</i> ; 2005;293:2141–48 8p.	Fel population/ wrong population.
Hicks TL, Goodall SF, Quattrone EM, Lydon-Rochelle MT. Postpartum sexual functioning and method of delivery: summary of the evidence. <i>J Midwifery Womens Health.</i> 2004 2004-01-01;49(5):430–6.	Intervention och behandling saknas/ No intervention/ treatment
Hofmeyr GJ, Gülmezoglu AM, Novikova N, Linder V, Ferreira S, Piaggio G. Misoprostol to prevent and treat postpartum haemorrhage: a systematic review and meta-analysis of maternal deaths and dose-related effects. <i>Bull World Health Organ.</i> 2009 2009-01-01;87(9):666-F 12p.	Fel population/ wrong population.
Hofmeyr GJ, Walraven G, Gulmezoglu AM, Maholwana B, Alfirovic Z, Villar J. Misoprostol to treat postpartum haemorrhage: a systematic review. <i>BJOG.</i> 2005 2005-01-01;112(5):547–53.	Fel population/ wrong population.
Kamel A, Khaled M. Episiotomy and obstetric perineal wound dehiscence: beyond soreness. <i>J Obstet Gynaecol.</i> 34(3):215–7.	Not a systematic review/Ej systematisk översikt
Kettle C, Hills RK, Ismail KMK. Continuous versus interrupted sutures for repair of episiotomy or second degree tears. <i>Cochrane Database Syst Rev.</i> 2007 2007-01-01;0(4).	Denna SR är sedan uppdaterad 2012 med nya sökningar – se ref 226.

Referens/Reference	Exklusionsorsak/ Reason for exclusion
Kettle C, Johanson RB. Continuous versus interrupted sutures for perineal repair. <i>Cochrane Database Syst Rev</i> (Online). 2000 2000-01-01;0(2):CD000947.	Duplicate/Duplikat
Knight M, Kurinczuk JJ, Spark P, Brocklehurst P. Emergency postpartum hysterectomy for uncontrolled postpartum bleeding: a systematic review. <i>Obstetrics &amp; Gynecology</i> . 2010 2010-01-01;115(6):1306–7 2p.	Not a systematic review/Ej systematisk översikt
Kotto-Kome AC, Calhoun DA, Montenegro R, Sosa R, Maldonado L, Christensen RD. Effect of administering recombinant erythropoietin to women with postpartum anemia: a meta-analysis. <i>Journal of Perinatology</i> . 2004 2004-01-01;24(1):11–5 5p.	Fel population/ wrong population
Likis FE, Sathe NA, Morgans AK, Hartmann KE, Young JL, Carlson-Bremer D, et al. Management of postpartum hemorrhage (Structured abstract). <i>Health Technology Assessment Database</i> . 2015 2015-01-01;0(4).	Fel population/ wrong population
Mackeen AD, Packard Roger E, Ota E, Speer L. Antibiotic regimens for postpartum endometritis. <i>Cochrane Database Syst Rev</i> . 2015 2015-01-01;0(2).	Fel population/ wrong population
Markova V, Norgaard A, Jørgensen Karsten J, Langhoff-Roos J. Treatment for women with postpartum iron deficiency anaemia. <i>Cochrane Database Syst Rev</i> . 2015 2015-01-01;0(8).	Fel population/ wrong population
Martí-Carvajal AJ, Comunián-Carrasco G, Peña-Martí GE. Haematological interventions for treating disseminated intravascular coagulation during pregnancy and postpartum. <i>Cochrane Database Syst Rev</i> . 2011 2011-01-01;0:N.PAG-N.PAG 1p.	Fel population/ wrong population
Matsubara S, Yano H, Ohkuchi A, Kuwata T, Usui R, Suzuki M. Uterine compression sutures for postpartum hemorrhage: An overview. <i>Acta Obstet Gynecol Scand</i> . 2013 2013-01-01;92(4):378–85.	Not a systematic review/Ej systematisk översikt
Miller S, El Ayadi A. Meta-analysis of 3,651 women with severe obstetric hemorrhage/hypovolemic shock treated with non-pneumatic anti-shock garment. <i>International Journal of Gynecology and Obstetrics</i> . 2012 2012-01-01;119:S223.	Ingen systematisk översikt/Not a systematic review

Referens/Reference	Exklusionsorsak/ Reason for exclusion
Moore RA, Derry S, Mason L, McQuay HJ, Edwards J. Single dose oral indometacin for the treatment of acute postoperative pain. <i>Cochrane Database Syst Rev.</i> 2004 2004-01-01;0:N.PAG-N.PAG 1p.	Ej systematisk översikt/Not a systematic review
Mousa HA, Alfirevic Z. Treatment for primary postpartum haemorrhage. <i>Cochrane Database Syst Rev.</i> 2014 2014-01-01;0(2):N.PAG-N.PAG 1p.	Fel population/ Wrong population
Oblasser C, Christie J, McCourt C. Vaginal cones or balls to improve pelvic floor muscle performance and urinary continence in women postpartum: a quantitative systematic review and meta-analysis protocol. <i>Journal of Advanced Nursing.</i> 2015 2015-01-01;71(4):933–41 9p.	Ingen systematisk översikt/ No systematic review
Okafor UV. Emergency postpartum hysterectomy for uncontrolled postpartum bleeding: a systematic review. <i>Obstetrics &amp; Gynecology.</i> 2010 2010-01-01;116(6):1453– 1p.	Ingen systematisk översikt Not a systematic review/
Oladapo OT, Fawole B, Blum J, Abalos E. Advance misoprostol distribution for preventing and treating postpartum haemorrhage. <i>Cochrane Database Syst Rev.</i> 2012 2012-01-01;0(2):N.PAG-N.PAG 1p.	Fel population/ wrong population.
Olefile KM, Khondowe O, M'Rithaa D. Misoprostol for prevention and treatment of postpartum haemorrhage: A systematic review. <i>Curationis.</i> 2013 2013-01-01;36(1):E1-e10.	Fel population/ wrong population.
Poujade O, Ceccaldi PF, Davitian C, Amate P, Chatel P, Khater C, et al. Uterine necrosis following pelvic arterial embolization for post-partum hemorrhage: review of the literature. <i>European Journal of Obstetrics, Gynecology, &amp; Reproductive Biology.</i> 2013;170(2):309–14.	Ingen systematisk översikt Not a systematic review/
Rath W, Hackethal A, Bohlmann MK. Second-line treatment of postpartum haemorrhage (PPH). <i>Archives of Gynecology &amp; Obstetrics.</i> 286(3):549–61.	Fel population/ wrong population.
Roberge S, Bujold E. Single versus double layer closure and risk of uterine rupture: Systematic review and meta-analysis. <i>American Journal of Obstetrics and Gynecology.</i> 2009 2009-01-01;201(6):S198.	Fel population/ wrong population.
Rossi AC, Lee RH, Chmait RH. Emergency postpartum hysterectomy for uncontrolled postpartum bleeding: a systematic review. <i>Obstetrics &amp; Gynecology.</i> 2010 2010-01-01;115(3):637–44 8p.	Fel population/ wrong population.

Referens/Reference	Exklusionsorsak/ Reason for exclusion
Scheufele L. Pelvic floor muscle exercise and the postpartum female: a review of current literature. Journal of the Section on Women's Health. 2003 2003-01-01;27(2):29–36 8p.	Not a systematic review/Ej systematisk översikt
Sheldon WR, Blum J, Durocher J, Winikoff B. Misoprostol for the prevention and treatment of postpartum hemorrhage. Expert Opinion on Investigational Drugs.21(2):235–50.	Wrong population/Fel population
The effectiveness of a pelvic floor muscle exercise program on urinary incontinence following childbirth. Best Practice. 2005 2005-01-01;9(2):1–6 p.	Not a systematic review/Ej systematisk översikt
Tindell K, Garfinkel R, Abu-Haydar E, Ahn R, Burke T, Conn K, et al. Uterine balloon tamponade for the treatment of postpartum haemorrhage in resource-poor settings: a systematic review. BJOG. 2013 2013-01-01;120(1):5–14 0p.	Wrong population/Fel population
Wagg A, Bunn F. Unassisted pelvic floor exercises for postnatal women: a systematic review. Journal of Advanced Nursing. 2007 2007-01-01;58(5):407–17 11p.	Wrong population/Fel population
Wright CE, Chauhan SP, Abuhamad AZ. Bakri Balloon in the Management of Postpartum Hemorrhage: A Review. American Journal of Perinatology. 2014 2014-01-01;31(11):957–63 7p.	Wrong population/Fel population