

Bilaga 3 Exklusionslistor**Exklusionsorsaker**

3 – Fel population

4 – Fel intervention

5 – Fel kontrollgrupp

6 – Fel utfall

7 – Fel design

8 – Protokoll

9 – Fel publikationstyp

10 – Duplikat

11 – Fel språk

Farmaka

Referens	Exklusions- orsak
(1993). "How long should the elderly take antidepressants? A double-blind placebo-controlled study of continuation/prophylaxis therapy with dothiepin. Old Age Depression Interest Group." <i>Br J Psychiatry</i> 162: 175-182.	3
(1994). "A multicentre comparative trial of moclobemide, imipramine and placebo in major depressive disorder. UK Moclobemide Study Group." <i>Int Clin Psychopharmacol</i> 9: 109-113.	4
(2005). "Curbing drinking alleviates depression in dually diagnosed seniors: study examines effectiveness of naltrexone plus SSRI." <i>Brown University Geriatric Psychopharmacology Update</i> 9(8): 1.	4
(2009). "Nortriptyline outperforms paroxetine CR for treating PD with major depression... Parkinson disease." <i>Brown University Geriatric Psychopharmacology Update</i> 13: 1.	3
(2011). "Adjunctive T'ai Chi Chih with escitalopram for geriatric depression." <i>Brown University Psychopharmacology Update</i> 22(5): 3-4.	3
Ackerman, D. L., S. Greenland, A. Bystritsky and G. W. Small (1997). "Characteristics of fluoxetine versus placebo responders in a randomized trial of geriatric depression." <i>Psychopharmacol Bull</i> 33(4): 707-714.	3
Ackerman, D. L., S. Greenland, A. Bystritsky and G. W. Small (2000). "Side effects and time course of response in a placebo-controlled trial of fluoxetine for the treatment of geriatric depression." <i>J Clin Psychopharmacol</i> 20: 658-665.	4
Addington, D., J. Addington, S. Patten, G. Remington, J. Moamai, A. Labelle and L. Beauclair (2002). "Double-blind, placebo-controlled comparison of the efficacy of sertraline as treatment for a major depressive episode in patients with remitted schizophrenia." <i>J Clin Psychopharmacol</i> 22(1): 20-25.	3
Adeoye, O. M., R. A. Sweet, B. G. Pollock, M. D. Miller, B. H. Mulsant, K. B. Kastango and C. F. Reynolds (2000). "Bupropion plasma concentration and antidepressant response in elderly patients: A prospective, randomized, double-blind study." <i>International Journal of Geriatric Psychopharmacology</i> 2(3): 132-136.	4
Aguera-Ortiz, L. and G. Ramos (2006). "[Effectiveness of venlafaxine extended release and conventional antidepressants in elderly patients with depressive disorder]." <i>Actas Esp Psiquiatr</i> 34: 153-161.	11
Ahles, S., H. Gwirtsman and A. Halaris (1984). "Comparative cardiac effects of maprotiline and doxepin in elderly depressed patients." <i>Journal of Clinical Psychiatry</i> 45(11): 460-465.	4
Akerblad, A. C., F. Bengtsson, K. von and L. Ekselius (2006). "Response, remission and relapse in relation to adherence in primary care treatment of depression: a 2-year outcome study." <i>Int Clin Psychopharmacol</i> 21: 117-124.	4
Alam, M. Y., P. L. Jacobsen, Y. Chen, M. Serenko and A. R. Mahableshwarkar (2014). "Safety, tolerability, and efficacy of vortioxetine (Lu AA21004) in major depressive disorder: results of an open-label, flexible-dose, 52-week extension study." <i>Int Clin Psychopharmacol</i> 29(1): 36-44.	7
Alexopoulos, G. S., B. S. Meyers, R. C. Young, B. Kalayam, T. Kakuma, M. Gabrielle, J. A. Sirey and J. Hull (2000). "Executive dysfunction and long-term outcomes of geriatric depression." <i>Arch Gen Psychiatry</i> 57: 285-290.	4

Alexopoulos, G. S., C. F. Murphy, F. M. Gunning-Dixon, V. Latoussakis, D. Kanellopoulos, S. Klimstra, K. O. Lim and M. J. Hoptman (2008). "Microstructural white matter abnormalities and remission of geriatric depression." <i>American Journal of Psychiatry</i> 165: 238-244.	3
Almeida, O. P., N. T. Lautenschlager, S. Vasikaran, P. Leedman, A. Gelavis and L. Flicker (2006). "A 20-week randomized controlled trial of estradiol replacement therapy for women aged 70 years and older: effect on mood, cognition and quality of life." <i>Neurobiol Aging</i> 27(1): 141-149.	3
Alpert, J. E., L. A. Uebelacker, N. E. McLean, M. Abraham, J. F. Rosenbaum and M. Fava (1995). "The Mini-Mental State Examination among adult outpatients with major depressive disorder." <i>Psychother Psychosom</i> 63: 207-211.	6
Altamura, A. C., M. C. Mauri, F. Colacurcio, P. L. Scapicchio, C. Hadjchristos, G. Carucci, M. Minervini, R. Montanini, M. Perini and N. Rudas (1988). "Trazodone in late life depressive states: a double-blind multicenter study versus amitriptyline and mianserin." <i>Psychopharmacology (Berl)</i> : S34-36.	4
Altamura, A. C., M. C. Mauri, N. Rudas, B. Carpiello, R. Montanini, M. Perini, P. L. Scapicchio, C. Hadjchristos, G. Carucci and M. Minervini (1989) "Clinical activity and tolerability of trazodone, mianserin, and amitriptyline in elderly subjects with major depression: a controlled multicenter trial." <i>Clin Neuropharmacol</i> , S25-33; s34-27.	10
Altamura, A. C., N. De, G. Guercetti, G. Invernizzi, M. Percudani and S. A. Montgomery (1989). "Fluoxetine compared with amitriptyline in elderly depression: a controlled clinical trial." <i>Int J Clin Pharmacol Res</i> 9: 391-396.	7
Altintoprak, A. E., N. Zorlu, H. Coskunol, F. Akdeniz and G. Kitapcioglu (2008). "Effectiveness and tolerability of mirtazapine and amitriptyline in alcoholic patients with co-morbid depressive disorder: A randomized, double-blind study." <i>Hum Psychopharmacol</i> 23: 313-319.	3
Amsterdam, J. D. and D. J. Brunswick (2002). "Site variability in treatment outcome in antidepressant trials." <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> 26: 989-993.	4
Andersen, G., K. Vestergaard and L. Lauritzen (1994). "Effective treatment of poststroke depression with the selective serotonin reuptake inhibitor citalopram." <i>Stroke</i> 25: 1099-1104.	3
Anderson, I. M. (2000). "Selective serotonin reuptake inhibitors versus tricyclic antidepressants: A meta-analysis of efficacy and tolerability." <i>J Affect Disord</i> 58: 19-36.	3
Andreeescu, C., E. J. Lenze, B. H. Mulsant, J. L. Wetherell, A. E. Begley, S. Mazumdar, C. F. Reynolds and rd (2009). "High worry severity is associated with poorer acute and maintenance efficacy of antidepressants in late-life depression." <i>Depress Anxiety</i> 26: 266-272.	3
Anstey, K. and H. Brodaty (1995). "Antidepressants and the elderly: Double-blind trials 1987-1992." <i>Int J Geriatr Psychiatry</i> 10(4): 265-279.	3
Antonini, A., S. Tesei, A. Zecchinelli, P. Barone, G. De, M. Canesi, G. Sacilotto, N. Meucci, C. Mariani and G. Pezzoli (2006). "Randomized study of sertraline and low-dose amitriptyline in patients with Parkinson's disease and depression: Effect on quality of life." <i>Movement Disorders</i> 21: 1119-1122.	3
Appelberg, B. (2003). "Long term citalopram prevents recurrent depression in the elderly and is well tolerated." <i>Evidence Based Mental Health</i> 6: 24-24.	3

Areán, P. A. (2006). "One in 10 elderly people with minor or subsyndromal depression develops major depression within a year." <i>Evidence Based Mental Health</i> 9: 94-94.	7
Arminen, S. L., U. Ikonen, M. Pulkkinen, E. Leinonen, A. Mahlalanen, H. Koponen, K. Kourula, J. Ryynä, V. Korpela, M. L. Lehtonen, H. Vartiainen, V. Lehtinen and T. Tamminen (1992). "Paroxetine and imipramine: A 12-week, double-blind multicentre study in hospitalized depressed patients." <i>Nordic Journal of Psychiatry, Supplement</i> 46: 27-31.	3
Asakura, M. and O. Tajima (2005). "A randomized, double-blind study of fluvoxamine maleate in patients with depression or depressive state: A comparison in anticholinergic effects and cardiovascular adverse reactions between fluvoxamine maleate and imipramine hydrochloride." <i>Japanese Pharmacology and Therapeutics</i> 33: 773-787.	5
Ash, G., C. M. Dickens, F. H. Creed, M. I. Jayson and B. Tomenson (1999). "The effects of dothiepin on subjects with rheumatoid arthritis and depression." <i>Rheumatology (Oxford)</i> 38: 959-967.	3
Aslan, A., C. Balaceanu, A. Manoiu and M. Erdos (1986). "A double-blind study on the antidepressive effects of Gerovital H3 and Aslavital in the elderly." <i>Romanian Journal of Gerontology and Geriatrics</i> 7(2): 79-88.	3
Ather, S. A., S. I. Ankier and R. S. W. Middleton (1985). "A double-blind evaluation of trazodone in the treatment of depression in the elderly." <i>British Journal of Clinical Practice</i> 39(5): 192-199.	3
Baca, E., C. Gonzalez de, M. Garcia-Toro, F. Perez-Arnau and A. Porras-Chavarino (2003). "Sertraline is more effective than imipramine in the treatment of non-melancholic depression: results from a multicentre, randomized study." <i>Prog Neuropsychopharmacol Biol Psychiatry</i> 27: 493-500.	4
Bakhshi, V., M. Elliott, A. Gentili, M. Godschalk and T. Mulligan (2000). "Testosterone improves rehabilitation outcomes in ill older men." <i>J Am Geriatr Soc</i> 48: 550-553.	3
Bakish, D., A. Ravindran, C. Hooper and Y. Lapierre (1994). "Psychopharmacological treatment response of patients with a DSM-III diagnosis of dysthymic disorder." <i>Psychopharmacol Bull</i> 30: 53-59.	4
Bakker, T. J. E. M., H. J. Duivenvoorden, J. van der Lee, M. G. M. Olde Rikkert, A. T. F. Beekman and M. W. Ribbe (2013). "Benefit of an integrative psychotherapeutic nursing home program to reduce multiple psychiatric symptoms of psychogeriatric patients and caregiver burden after six months of follow-up: A re-analysis of a randomized controlled trial." <i>International Psychogeriatrics</i> 25(1): 34-46.	4
Balaceanu-Stolnici, C., M. Covic, A. Manoiu and M. Vrabiescu (1996). "Double blind study concerning the antidepressive effects and the clinical tolerance of Gerovital H3 without potassium metabisulphite." <i>Romanian Journal of Gerontology and Geriatrics</i> 17(1-2): 46-61.	3
Baldomero, E. B., J. G. Ubago, C. L. Cercos, J. V. Ruiloba, C. G. Calvo and R. P. Lopez (2005). "Venlafaxine extended release versus conventional antidepressants in the remission of depressive disorders after previous antidepressant failure: ARGOS study." <i>Depress Anxiety</i> 22: 68-76.	3
Baldwin, D. S., H. Loft and M. Dragheim (2012). "A randomised, double-blind, placebo controlled, duloxetine-referenced, fixed-dose study of three dosages of Lu AA21004 in acute treatment of major depressive disorder (MDD)." <i>Eur Neuropsychopharmacol</i> 22: 482-491.	3

Baldwin, D. S., T. Hansen and I. Florea (2012). "Vortioxetine (Lu AA21004) in the long-term open-label treatment of major depressive disorder." <i>Curr Med Res Opin</i> 28: 1717-1724.	3
Banerjee, S., J. Hellier, R. Romeo, M. Dewey, M. Knapp, C. Ballard, R. Baldwin, P. Bentham, C. Fox, C. Holmes, C. Katona, C. Lawton, J. Lindesay, G. Livingston, N. McCrae, E. Moniz-Cook, J. Murray, S. Nurock, M. Orrell, J. O'Brien, M. Poppe, A. Thomas, R. Walwyn, K. Wilson and A. Burns (2013). "Study of the use of antidepressants for depression in dementia: The HTA-SADD trial- A multicentre, randomised, double-blind, placebo-controlled trial of the clinical effectiveness and cost-effectiveness of sertraline and mirtazapine." <i>Health Technology Assessment</i> 17: 1-43.	10
Bao, Y., E. P. Post, T. R. Ten Have, B. R. Schackman and M. L. Bruce (2009). "Achieving effective antidepressant pharmacotherapy in primary care: the role of depression care management in treating late-life depression." <i>J Am Geriatr Soc</i> 57(5): 895-900.	3
Barone, P., L. Scarzella, R. Marconi, A. Antonini, L. Morgante, F. Bracco, M. Zappia, B. Musch, Depression and G. Parkinson Italian Study (2006). "Pramipexole versus sertraline in the treatment of depression in Parkinson's disease: a national multicenter parallel-group randomized study." <i>J Neurol</i> 253: 601-607.	3
Barone, P., W. Poewe, S. Albrecht, C. Debieuvre, D. Massey, O. Rascol, E. Tolosa and D. Weintraub (2010). "Pramipexole for the treatment of depressive symptoms in patients with Parkinson's disease: a randomised, double-blind, placebo-controlled trial." <i>Lancet Neurol</i> 9: 573-580.	3
Battaglia, A. and M. Bejor (2001). "Influence of poststroke depression on functional outcome." <i>Europa Medicophysica</i> 37: 25-37.	3
Bauer, M., K. Demyttenaere, N. El-Khalili, M. E. Thase, G. I. Papakostas, J. Szamosi, W. R. Earley and H. Eriksson (2014). "Pooled analysis of adjunct extended-release quetiapine fumarate in patients with major depressive disorder according to ongoing SSRI or SNRI treatment." <i>Int Clin Psychopharmacol</i> 29(1): 16-25.	3
Bauer, M., L. Dell'osso, S. Kasper, W. Pitchot, E. Dencker Vansvik, J. Kohler, L. Jorgensen and S. A. Montgomery (2013). "Extended-release quetiapine fumarate (quetiapine XR) monotherapy and quetiapine XR or lithium as add-on to antidepressants in patients with treatment-resistant major depressive disorder." <i>J Affect Disord</i> 151(1): 209-219.	3
Bauer, M., N. El-Khalili, C. Datto, J. Szamosi and H. Eriksson (2010). "A pooled analysis of two randomised, placebo-controlled studies of extended release quetiapine fumarate adjunctive to antidepressant therapy in patients with major depressive disorder." <i>J Affect Disord</i> 127(1-3): 19-30.	3
Bauer, M., R. Zaninelli, B. Muller-Oerlinghausen and W. Meister (1999). "Paroxetine and amitriptyline augmentation of lithium in the treatment of major depression: a double-blind study." <i>J Clin Psychopharmacol</i> 19(2): 164-171.	7
Baumhackl, U., K. Biziere, R. Fischbach, C. Geretsegger, G. Hebenstreit, E. Radmayr and M. Stabl (1989). "Efficacy and tolerability of moclobemide compared with imipramine in depressive disorder (DSM-III): an Austrian double-blind, multicentre study." <i>Br J Psychiatry Suppl</i> : 78-83.	4
Bavazzano, A., R. Guarducci, G. Gestri and P. Sala (1980). "Clinical trial with amantadine and hydergine in elderly patients." <i>Journal of Clinical and Experimental Gerontology</i> 2(4): 289-299.	3

Bayer, A. J., M. S. Pathy, A. Cameron, T. Venkateswalu, S. A. Ather, G. D. Mankikar and A. K. Ghosh (1989). "A comparative study of conventional and controlled-release formulations of trazodone in elderly depressed patients." <i>Clin Neuropharmacol</i> 12 Suppl 1: S50-55; Discussion S56-57.	3
Beasley, Jr., M. C, M. E. Sayler, J. C. Bosomworth and J. F. Wernicke (1991). "High-dose fluoxetine: Efficacy and activating-sedating effects in agitated and retarded depression." <i>J Clin Psychopharmacol</i> 11: 166-174.	6
Beaumont, G., M. Gringras, F. D. Hobbs, V. W. Drury, P. Freeling, A. Tylee, D. F. Shanley, H. Comber, S. Little and S. Hill (1993). "A randomized, double-blind, multi-centre, parallel-group study comparing the tolerability and efficacy of moclobemide and dothiepin hydrochloride in depressed patients in general practice." <i>Int Clin Psychopharmacol</i> 7: 159-165.	4
Bech, P., S. L. Lonn and K. F. Overo (2010). "Relapse prevention and residual symptoms: A closer analysis of placebo-controlled continuation studies with escitalopram in major depressive disorder, generalized anxiety disorder, social anxiety disorder, and obsessive-compulsive disorder." <i>Journal of Clinical Psychiatry</i> 71: 121-129.	7
Behnke, K., B. Mejer-Nielsen, A. Korner, P. Arup, A. Geisler, M. Sastre-y-Hernandez, R. Bischoff, M. Schratzer and B. Voet (1992). "Roliplram versus nortriptyline in gerontopsychiatric inpatients with major depression." <i>Nord J Psychiatry</i> 46: 407-411.	4
Behnke, K., J. Sogaard, S. Martin, J. Bauml, A. V. Ravindran, H. Agren and E. D. Vester-Blokland (2003). "Mirtazapine orally disintegrating tablet versus sertraline: a prospective onset of action study." <i>J Clin Psychopharmacol</i> 23: 358-364.	8
Bell, I. R., J. S. Edman, F. D. Morrow, D. W. Marby, G. Perrone, H. L. Kayne, M. Greenwald and J. O. Cole (1992). "Brief communication. Vitamin B1, B2, and B6 augmentation of tricyclic antidepressant treatment in geriatric depression with cognitive dysfunction." <i>J Am Coll Nutr</i> 11(2): 159-163.	4
Bella, R., R. Biondi, R. Raffaele and G. Pennisi (1990). "Effect of acetyl-L-carnitine on geriatric patients suffering from dysthymic disorders." <i>Int J Clin Pharmacol Res</i> 10(6): 355-360.	3
Bellino, S., G. Barzega, F. Bogetto, G. Maina, S. Venturello and L. Ravizza (1997). "An open-label, randomized, prospective comparison of sertraline and amisulpride in the treatment of dysthymia in the elderly." <i>Current Therapeutic Research - Clinical and Experimental</i> 58: 798-808.	3
Benes, H., W. Mattern, I. Peglau, T. Dreykluft, L. Bergmann, C. Hansen, R. Kohnen, N. Banik, S. W. Schoen and M. Hornyak (2011). "Ropinirole improves depressive symptoms and restless legs syndrome severity in RLS patients: a multicentre, randomized, placebo-controlled study." <i>J Neurol</i> 258: 1046-1054.	3
Benkert, O., A. Szegedi and R. Kohnen (2000). "Mirtazapine compared with paroxetine in major depression." <i>J Clin Psychiatry</i> 61: 656-663.	6
Benkert, O., A. Szegedi, M. Philipp, R. Kohnen, C. Heinrich, A. Heukels, V.-S. van der, R. A. Baker, J. H. Simmons and A. J. Schutte (2006). "Mirtazapine orally disintegrating tablets versus venlafaxine extended release: a double-blind, randomized multicenter trial comparing the onset of antidepressant response in patients with major depressive disorder." <i>J Clin Psychopharmacol</i> 26: 75-78.	9

Bergh, S., G. Selbaek and K. Engedal (2012). "Discontinuation of antidepressants in people with dementia and neuropsychiatric symptoms (DESEP study): double blind, randomised, parallel group, placebo controlled trial." <i>BMJ</i> 344: e1566.	3
Bergman, J., C. Miodownik, A. Palatnik and V. Lerner (2011). "Efficacy of bupropion XR in treatment-resistant elderly patients: a case series study." <i>Clin Neuropharmacol</i> 34: 17-20.	6
Berman, R. M., A. M. Darnell, H. L. Miller, A. Anand and D. S. Charney (1997). "Effect of pindolol in hastening response to fluoxetine in the treatment of major depression: a double-blind, placebo-controlled trial." <i>Am J Psychiatry</i> 154: 37-43.	3
Berman, R. M., M. Fava, M. E. Thase, M. H. Trivedi, R. Swanink, R. D. McQuade, W. H. Carson, D. Adson, L. Taylor, J. Hazel and R. N. Marcus (2009). "Aripiprazole augmentation in major depressive disorder: a double-blind, placebo-controlled study in patients with inadequate response to antidepressants." <i>CNS Spectr</i> 14(4): 197-206.	3
Berman, R. M., R. N. Marcus, R. Swanink, R. D. McQuade, W. H. Carson, P. K. Corey-Lisle and A. Khan (2007). "The efficacy and safety of aripiprazole as adjunctive therapy in major depressive disorder: A multicenter, randomized, double-blind, placebo-controlled study." <i>Journal of Clinical Psychiatry</i> 68: 843-853.	7
Bersani, G., G. Meco, A. Denaro, D. Liberati, C. Colletti, R. Nicolai, F. S. Bersani and A. Kovarech (2013). "L-Acetylcarnitine in dysthymic disorder in elderly patients: A double-blind, multicenter, controlled randomized study vs. fluoxetine." <i>European Neuropsychopharmacology</i> .	4
Bettini, R., M. Ferrari, G. Gobbi, F. Volanti and O. Piccinelli (1994). "A clinical and instrumental evaluation of the effect of tiapride on senile involution syndrome." <i>Clinica Terapeutica</i> 144(3): 231-236.	11
Bignamini, A. and V. Rapisarda (1992). "A double-blind multicentre study of paroxetine and amitriptyline in depressed outpatients. Italian Paroxetine Study Group." <i>Int Clin Psychopharmacol</i> 6 Suppl 4: 37-41.	3
Bird, H. and M. Broggini (2000). "Paroxetine versus amitriptyline for treatment of depression associated with rheumatoid arthritis: a randomized, double blind, parallel group study." <i>J Rheumatol</i> 27: 2791-2797.	3
Birkenhager, T. K., B. Van Den, W. W, P. G. Mulder, P. Moleman and J. A. Bruijn (2008). "Efficacy of imipramine in psychotic versus nonpsychotic depression." <i>J Clin Psychopharmacol</i> 28: 166-170.	3
Birkenhager, T. K., W. W. van den Broek, P. G. Mulder, J. A. Bruijn and P. Moleman (2004). "Comparison of two-phase treatment with imipramine or fluvoxamine, both followed by lithium addition, in inpatients with major depressive disorder." <i>Am J Psychiatry</i> 161(11): 2060-2065.	3
Birkenhager, T. K., W. W. van den Broek, P. G. Mulder, J. A. Bruijn and P. Moleman (2006). "[Comparison of the effectiveness of two treatment strategies in inpatients with a depressive disorder. A double-blind study of imipramine followed by lithium addition versus fluvoxamine followed by lithium addition]." <i>Tijdschr Psychiatr</i> 48(4): 271-281.	11
Blasey Christine, M., S. Block Thaddeus, K. Belanoff Joseph and L. Roe Robert (2011). "Efficacy and safety of mifepristone for the treatment of psychotic depression." <i>J Clin Psychopharmacol</i> 31: 436-440.	4

Blumenfield, M., N. B. Levy, B. Spinowitz, C. Charytan, C. M. Beasley, Jr, A. K. Dubey, R. J. Solomon, R. Todd, A. Goodman and R. F. Bergstrom (1997). "Fluoxetine in depressed patients on dialysis." <i>Int J Psychiatry Med</i> 27: 71-80.	3
Bobo, W. V., H. Chen, M. H. Trivedi, J. W. Stewart, A. A. Nierenberg, M. Fava, B. T. Kurian, D. Warden, D. W. Morris, J. F. Luther, M. M. Husain, I. A. Cook, I. M. Lesser, S. G. Kornstein, S. R. Wisniewski, A. J. Rush and R. C. Shelton (2011). "Randomized comparison of selective serotonin reuptake inhibitor (escitalopram) monotherapy and antidepressant combination pharmacotherapy for major depressive disorder with melancholic features: a CO-MED report." <i>J Affect Disord</i> 133: 467-476.	9
Bogner, H. R., K. H. Morales, C. F. Reynolds, 3rd, M. S. Cary and M. L. Bruce (2012). "Course of depression and mortality among older primary care patients." <i>Am J Geriatr Psychiatry</i> 20(10): 895-903.	3
Bogner, H. R., K. H. Morales, E. P. Post and M. L. Bruce (2007). "Diabetes, depression, and death: a randomized controlled trial of a depression treatment program for older adults based in primary care (PROSPECT)." <i>Diabetes Care</i> 30(12): 3005-3010.	4
Bogolepova, A. N., E. I. Chukanova, M. Y. Smirnova, A. S. Chukanova, I. Y. Gracheva and E. G. Semushkina (2012). "Experience in the use of Valdoxan in the treatment of post-stroke depression." <i>Neuroscience and Behavioral Physiology</i> 42: 556-560.	4
Bohm, C. and D. S. G. R. E. Robinson (1990) "Buspirone therapy for elderly patients with anxiety or depressive neurosis letter." <i>J-Clin-Psychiatry</i> , 309.	3
Bohm, C., D. S. Robinson, R. E. Gammans, R. C. Shrotriya, D. R. Alms, A. Leroy and M. Placchi (1990). "Buspirone therapy in anxious elderly patients: a controlled clinical trial." <i>J Clin Psychopharmacol</i> 10(3 Suppl): 47S-51S.	3
Borson, S., G. J. McDonald, T. Gayle, M. Deffebach, S. Lakshminarayan and C. VanTuinen (1992). "Improvement in mood, physical symptoms, and function with nortriptyline for depression in patients with chronic obstructive pulmonary disease." <i>Psychosomatics</i> 33: 190-201.	3
Bortnick, B., N. El-Khalili, M. Banov, D. Adson, C. Datto, S. Raines, W. Earley and H. Eriksson (2011). "Efficacy and tolerability of extended release quetiapine fumarate (quetiapine XR) monotherapy in major depressive disorder: a placebo-controlled, randomized study." <i>J Affect Disord</i> 128: 83-94.	6
Bose, A., D. Li and C. Gandhi (2008). "Escitalopram in the acute treatment of depressed patients aged 60 years or older." <i>Am J Geriatr Psychiatry</i> 16: 14-20.	7
Bot, M., R. M. Carney, K. E. Freedland, E. H. Rubin, M. W. Rich, B. C. Steinmeyer and D. L. Mann (2011). "Inflammation and treatment response to sertraline in patients with coronary heart disease and comorbid major depression." <i>J Psychosom Res</i> 71: 13-17.	7
Boulenger, J. P., H. Loft and I. Florea (2012). "A randomized clinical study of Lu AA21004 in the prevention of relapse in patients with major depressive disorder." <i>J Psychopharmacol</i> 26: 1408-1416.	3
Boyer, P., J. M. Danion, J. C. Bisserbe, J. M. Hotton and S. Troy (1998). "Clinical and economic comparison of sertraline and fluoxetine in the treatment of depression. A 6-month double-blind study in a primary-care setting in France." <i>Pharmacoeconomics</i> 13(1 Pt 2): 157-169.	3

Branconnier, R. J., J. O. Cole and G. F. Oxenkrug (1983). "Cardiovascular effects of imipramine and bupropion in aged depressed patients." <i>Psychopharmacol Bull</i> 19(4): 658-662.	3
Branconnier, R. J., J. O. Cole and S. Ghazvinian (1981). "The therapeutic profile of mianserin in mild elderly depressives." <i>Psychopharmacol Bull</i> 17: 129-131.	3
Branconnier, R. J., J. O. Cole and S. Ghazvinian (1983). "Clinical pharmacology of bupropion and imipramine in elderly depressives." <i>Journal of Clinical Psychiatry</i> 44(5 II): 130-133.	10
Brannan, S. K., C. H. Mallinckrodt, E. B. Brown, M. M. Wohlreich, J. G. Watkin and A. F. Schatzberg (2005). "Duloxetine 60 mg once-daily in the treatment of painful physical symptoms in patients with major depressive disorder." <i>J Psychiatr Res</i> 39(1): 43-53.	3
Brannan, S. K., C. H. Mallinckrodt, M. J. Detke, J. G. Watkin and G. D. Tollefson (2005). "Onset of action for duloxetine 60 mg once daily: Double-blind, placebo-controlled studies." <i>J Psychiatr Res</i> 39: 161-172.	3
Brecht, S., C. Courtecuisse, C. Debieuvre, J. Croenlein, D. Desaiah, J. Raskin, C. Petit and K. Demyttenaere (2007). "Efficacy and safety of duloxetine 60 mg once daily in the treatment of pain in patients with major depressive disorder and at least moderate pain of unknown etiology: a randomized controlled trial." <i>J Clin Psychiatry</i> 68: 1707-1716.	3
Bricker, J. B., J. Russo, M. B. Stein, C. Sherbourne, M. Craske, T. J. Schraufnagel and P. Roy-Byrne (2007). "Does occasional cannabis use impact anxiety and depression treatment outcomes?: Results from a randomized effectiveness trial." <i>Depress Anxiety</i> 24(6): 392-398.	3
Brion, S., S. Audrain and B. de (1996). "[Major depressive episodes in patients over 70 years of age. Evaluation of the efficiency and acceptability of tianeptine and mianserin]." <i>Presse Med</i> 25: 461-468.	11
Brion, S., S. Audrain and B. De (1996). "Episodes of severe depression in subjects over 70 years of age. Efficacy and acceptability of tianeptine and mianserine." <i>Presse Medicale</i> 25: 461-468.	11
Brodie, N. H., R. L. McGhie and H. O'Hara (1975). "Anxiety/depression in elderly patients. A double blind comparative study of fluphenazine/nortriptyline and promazine." <i>Practitioner</i> 215(1289): 660-664.	9
Bruijn, J. A., P. Moleman, P. G. Mulder and W. W. van den Broek (1998). "Comparison of 2 treatment strategies for depressed inpatients: imipramine and lithium addition or mirtazapine and lithium addition." <i>J Clin Psychiatry</i> 59(12): 657-663.	3
Bump, G. M., B. H. Mulsant, B. G. Pollock, S. Mazumdar, A. E. Begley, M. A. Dew, C. F. Reynolds and rd (2001). "Paroxetine versus nortriptyline in the continuation and maintenance treatment of depression in the elderly." <i>Depress Anxiety</i> 13: 38-44.	3
Burke, W. J., I. Gergel and A. Bose (2002). "Fixed-dose trial of the single isomer SSRI escitalopram in depressed outpatients." <i>J Clin Psychiatry</i> 63: 331-336.	3
Butters, M. A., J. T. Becker, R. D. Nebes, M. D. Zmuda, B. H. Mulsant, B. G. Pollock and C. F. Reynolds, 3rd (2000). "Changes in cognitive functioning following treatment of late-life depression." <i>Am J Psychiatry</i> 157(12): 1949-1954.	4
Cai, X. J., X. P. Bi, Z. Zhao and L. Wang (2006). "[The effects of antidepressant treatment on efficacy of antihypertensive therapy in elderly hypertension]." <i>Zhonghua Nei Ke Za Zhi</i> 45: 639-641.	11

Carney, R. M., K. E. Freedland, P. K. Stein, B. C. Steinmeyer, W. S. Harris, E. H. Rubin, R. J. Krone and M. W. Rich (2010). "Effect of omega-3 fatty acids on heart rate variability in depressed patients With Coronary Heart Disease." <i>Psychosom Med</i> 72: 748-754.	3
Carta, M. G., F. Zairo, G. Mellino and M. C. Hardoy (2007). "Add-on quetiapine in the treatment of major depressive disorder in elderly patients with cerebrovascular damage." <i>Clin Pract Epidemiol Ment Health</i> 3: 28.	4
Casacchia, M., F. Bolino, W. Marola, R. Pirro, G. Nivoli, V. Rapisarda and P. Pancheri (1994). "Controlled multicentre study of teniloxazine in mild depression of the elderly." <i>New Trends in Experimental and Clinical Psychiatry</i> 10(4): 187-192.	4
Casper, R. C., G. D. Tollefson and M. E. Nilsson (2001). "No gender differences in placebo responses of patients with major depressive disorder." <i>Biol Psychiatry</i> 49: 158-160.	3
Cavallini, G., S. Caracciolo, G. Vitali, F. Modenini and G. Biagiotti (2004). "Carnitine versus androgen administration in the treatment of sexual dysfunction, depressed mood, and fatigue associated with male aging." <i>Urology</i> 63(4): 641-646.	3
Chan, A. S., M. C. Cheung, W. J. Tsui, S. L. Sze and D. Shi (2011). "Dejian mind-body intervention on depressive mood of community-dwelling adults: a randomized controlled trial." <i>Evid Based Complement Alternat Med</i> 2011: 473961.	3
Chan, H. N., A. John Rush, A. A. Nierenberg, M. Trivedi, S. R. Wisniewski, G. K. Balasubramani, E. S. Friedman, B. N. Gaynes, L. Davis, D. Morris and M. Fava (2012). "Correlates and outcomes of depressed out-patients with greater and fewer anxious symptoms: A CO-MED report." <i>International Journal of Neuropsychopharmacology</i> 15(10): 1387-1399.	4
Chauvet-Gelinier, J. C. (2010). "Efficacy of escitalopram vs paroxetine on severe depression with associated anxiety: Data from the "boulenger" study." <i>Encephale</i> 36: 425-432.	11
Chemerinski, E., R. G. Robinson, S. Arndt and J. T. Kosier (2001). "The effect of remission of poststroke depression on activities of daily living in a double-blind randomized treatment study." <i>J Nerv Ment Dis</i> 189: 421-425.	4
Chen, S., Y. Fang, L. Li and H. Fan (2012). "Collaborative late-life depression care management in urban China primary care." <i>Asia-Pacific Psychiatry</i> 4: 97.	3
Chen, Y. M., X. M. Huang, R. Thompson and Y. B. Zhao (2011). "Clinical features and efficacy of escitalopram treatment for geriatric depression." <i>J Int Med Res</i> 39(5): 1946-1953.	3
Chesrow, E. J., S. E. Kaplitz, R. Sabatini, H. Vetra and G. H. Marquardt (1965). "A new psychotherapeutic agent effective in the management of geriatric anxiety, depression and behavioral reactions." <i>J Amer. Geriat. Soc.</i> .. 449-454.	3
Choi-Kwon, S., J. Choi, S. U. Kwon, D. W. Kang and J. S. Kim (2008). "Fluoxetine improves the quality of life in patients with poststroke emotional disturbances." <i>Cerebrovasc Dis</i> 26: 266-271.	3
Choi-Kwon, S., S. W. Han, S. U. Kwon, D. W. Kang, J. M. Choi and J. S. Kim (2006). "Fluoxetine treatment in poststroke depression, emotional incontinence, and anger proneness: a double-blind, placebo-controlled study." <i>Stroke</i> 37: 156-161.	3
Christensen, H., A. Aiken, P. J. Batterham, J. Walker, A. J. Mackinnon, M. Fenech and I. B. Hickie (2011). "No clear potentiation of antidepressant medication effects by folic acid+vitamin B12 in a large community sample." <i>J Affect Disord</i> 130: 37-45.	4

Christensen, P., H. Y. Thomsen and O. L. Pedersen (1985). "Cardiovascular effects of amitriptyline in the treatment of elderly depressed patients." <i>Psychopharmacology (Berl)</i> 87(2): 212-215.	9
Ciechanowski, P. S., J. E. Russo, W. J. Katon, M. V. Korff, G. E. Simon, E. H. Lin, E. J. Ludman and B. A. Young (2006). "The association of patient relationship style and outcomes in collaborative care treatment for depression in patients with diabetes." <i>Med Care</i> 44(3): 283-291.	3
Claghorn James, L., Q. Earl Craig, D. Walczak Donna and A. Stoner Kim (1996). "Fluvoxamine maleate in the treatment of depression: A single-center, double-blind, placebo-controlled comparison with imipramine in outpatients." <i>J Clin Psychopharmacol</i> 16: 113-120.	4
Claghorn, J. L. and J. P. Feighner (1993). "A double-blind comparison of paroxetine with imipramine in the long-term treatment of depression." <i>J Clin Psychopharmacol</i> 13(6 Suppl 2): 23S-27S.	7
Clarke, R. (2012). "Folic acid and vitamin B12 for prevention of cognitive decline in older people with depression." <i>Aging Health</i> 8: 115-117.	3
Clayton, A. H., S. G. Kornstein, G. Rosas, C. Guico-Pabia and K. A. Tourian (2009). "An integrated analysis of the safety and tolerability of desvenlafaxine compared with placebo in the treatment of major depressive disorder." <i>CNS Spectr</i> 14: 183-195.	3
Clayton, A., S. Kornstein, A. Prakash, C. Mallinckrodt and M. Wohlreich (2007). "Changes in sexual functioning associated with duloxetine, escitalopram, and placebo in the treatment of patients with major depressive disorder." <i>Journal of Sexual Medicine</i> 4(4 I): 917-929.	3
Cohn, C. K., R. Shrivastava, J. Mendels, J. B. Cohn, L. F. Fabre, J. L. Claghorn, E. C. Dessain, T. M. Itil and A. Lautin (1990). "Double-blind, multicenter comparison of sertraline and amitriptyline in elderly depressed patients." <i>J Clin Psychiatry</i> 51 Suppl B: 28-33.	3
Cohn, J. B., L. Varga and A. Lyford (1984). "A two-center double-blind study of nomifensine, imipramine and placebo in depressed geriatric outpatients." <i>Journal of Clinical Psychiatry</i> 45: 68-72.	3
Cohn, J., J. Claghorn and D. Ostrow (1991). "A comparative trial of trimipramine and doxepin in depressed elderly patients." <i>Adv Ther</i> 8(2): 92-102.	9
Coleman, C. C., B. R. King, C. Bolden-Watson, M. J. Book, S. Taylor, N. Richard, J. Ascher, S. Batey, B. Jamerson and A. Metz (2001). "A placebo-controlled comparison of the effects on sexual functioning of bupropion sustained release and fluoxetine." <i>Clin Ther</i> 23: 1040-1058.	4
Coleman, C. C., L. A. Cunningham, V. J. Foster, S. R. Batey, R. M. Donahue, T. L. Houser and J. A. Ascher (1999). "Sexual dysfunction associated with the treatment of depression: a placebo-controlled comparison of bupropion sustained release and sertraline treatment." <i>Ann Clin Psychiatry</i> 11: 205-215.	3
Colonna, L., H. F. Andersen and E. H. Reines (2005). "A randomized, double-blind, 24-week study of escitalopram (10 mg/day) versus citalopram (20 mg/day) in primary care patients with major depressive disorder." <i>Curr Med Res Opin</i> 21: 1659-1668.	3
Cook, B. L., P. M. Helms, R. E. Smith and M. Tsai (1986). "Unipolar depression in the elderly: Reoccurrence on discontinuation of tricyclic antidepressants." <i>J Affect Disord</i> 10: 91-94.	3

Cookson, J., I. Gilaberte, D. Desaiah and D. K. Kajdasz (2006). "Treatment benefits of duloxetine in major depressive disorder as assessed by number needed to treat." <i>Int Clin Psychopharmacol</i> 21: 267-273.	3
Corruble, E., C. Goldberger and M. Spann (2010). "Relationship between TSH levels in the normal range and short-term duloxetine efficacy." <i>J Affect Disord</i> 123: 312-316.	7
Cravello, L., C. Caltagirone and G. Spalletta (2009). "The SNRI venlafaxine improves emotional unawareness in patients with post-stroke depression." <i>Hum Psychopharmacol</i> 24: 331-336.	3
Croft, H., E. Settle, Jr, T. Houser, S. R. Batey, R. M. Donahue and J. A. Ascher (1999). "A placebo-controlled comparison of the antidepressant efficacy and effects on sexual functioning of sustained-release bupropion and sertraline." <i>Clin Ther</i> 21: 643-658.	4
Croft, H., T. L. Houser, B. D. Jamerson, R. Leadbetter, C. Bolden-Watson, R. Donahue and A. Metz (2002). "Effect on body weight of bupropion sustained-release in patients with major depression treated for 52 weeks." <i>Clin Ther</i> 24: 662-672.	6
Culang, M., J. R. Snead, J. G. Keilp and S. P. Roose (2010). "Change in cognitive functioning following antidepressant treatment in late-life depression." <i>American Journal of Geriatric Psychiatry</i> 18(3): S59.	3
Culang-Reinlieb, M. E., J. R. Snead, J. G. Keilp and S. P. Roose (2012). "Change in cognitive functioning in depressed older adults following treatment with sertraline or nortriptyline." <i>Int J Geriatr Psychiatry</i> 27(8): 777-784.	4
Culang-Reinlieb, M., J. R. Snead, J. Keilp and S. Roose (2011). "The impact of antidepressant treatment on cognitive functioning in depressed older adults." <i>American Journal of Geriatric Psychiatry</i> 19: S59-S60.	3
Cunningham, L. A. (1997). "Once-daily venlafaxine extended release (XR) and venlafaxine immediate release (IR) in outpatients with major depression. Venlafaxine XR 208 Study Group." <i>Ann Clin Psychiatry</i> 9: 157-164.	3
Cutler, A. J., S. A. Montgomery, D. Feifel, A. Lazarus, M. Astrom and M. Brecher (2009). "Extended release quetiapine fumarate monotherapy in major depressive disorder: A placebo- and duloxetine-controlled study." <i>Journal of Clinical Psychiatry</i> 70: 526-539.	3
Dalery, J. and A. Honig (2003). "Fluvoxamine versus fluoxetine in major depressive episode: a double-blind randomised comparison." <i>Hum Psychopharmacol</i> 18: 379-384.	6
Dalery, J. and V. Aubin (2001). "[Comparative study of paroxetine and mianserin in depression in elderly patients: efficacy, tolerance, serotonin dependence]." <i>Encephale</i> 27(1): 71-81.	4 och 11
Dalery, J., C. Rochat, E. Peyron and G. Bernard (1997). "The efficacy and acceptability of amineptine versus fluoxetine in major depression." <i>Int Clin Psychopharmacol</i> 12 Suppl 3: S35-38.	3
Danjou, P. and D. Hackett (1995). "Safety and tolerance profile of venlafaxine." <i>Int Clin Psychopharmacol</i> 10: 15-20.	3
Datto, C. J., R. Thompson, K. Knott and I. R. Katz (2006). "Older adult report of change in depressive symptoms as a treatment decision tool." <i>J Am Geriatr Soc</i> 54: 627-631.	3

Dawes, S. E., B. W. Palmer, T. Meeks, S. Golshan, J. Kasckow, S. Mohamed and S. Zisook (2012). "Does antidepressant treatment improve cognition in older people with schizophrenia or schizoaffective disorder and comorbid subsyndromal depression?" <i>Neuropsychobiology</i> 65(3): 168-172.	3
Davies, G., S. Hamilton and E. Hendrickson (1977). "The effect of cyclandelate in depressed and demented patients: A controlled study in psychogeriatric patients." <i>Age Ageing</i> 6(3): 156-162.	3
Davis, L. L., P. Pilkinton, S. R. Wisniewski, M. H. Trivedi, B. N. Gaynes, R. H. Howland, S. Zisook, G. K. Balasubramani, M. Fava and A. J. Rush (2012). "Effect of concurrent substance use disorder on the effectiveness of single and combination antidepressant medications for the treatment of major depression: An exploratory analysis of a single-blind randomized trial." <i>Depress Anxiety</i> 29(2): 111-122.	3
Davis, L. L., S. R. Wisniewski, R. H. Howland, M. H. Trivedi, M. M. Husain, M. Fava, P. J. McGrath, G. K. Balasubramani, D. Warden and A. J. Rush (2010). "Does comorbid substance use disorder impair recovery from major depression with SSRI treatment? An analysis of the STAR*D level one treatment outcomes." <i>Drug Alcohol Depend</i> 107(2-3): 161-170.	3
Davis, M. R., D. Cohen and C. Eisdorfer (1982). "Anxiety and depression in the aged with sleep disturbances." <i>Journal of Clinical and Experimental Gerontology</i> 4(3): 239-247.	3
De Fruyt, F., K. Van Leeuwen, R. M. Bagby, J. P. Rolland and F. Rouillon (2006). "Assessing and interpreting personality change and continuity in patients treated for major depression." <i>Psychol Assess</i> 18(1): 71-80.	3
de Jonghe, F., M. Hendricksen, G. van Aalst, S. Kool, V. Peen, R. Van, E. van den Eijnden and J. Dekker (2004). "Psychotherapy alone and combined with pharmacotherapy in the treatment of depression." <i>Br J Psychiatry</i> 185: 37-45.	3
de Vasconcelos, C., G. U, R. Lopes, M. Avila de, V. Alves, N. de Souza, J. J, B. Mendes, S. Magalhaes, S. Araujo, O. Martins de, J. da Costa, L. A, A. Faria, X. V and E. Sakurai (2007). "A placebo-controlled double-blind randomized study of venlafaxine in the treatment of depression in dementia." <i>Dement Geriatr Cogn Disord</i> 24: 36-41.	3
De, L., A. Ceola and G. Magni (1984). "Viloxazine against placebo in a double-blind study in depressed elderly patients." <i>Current Therapeutic Research - Clinical and Experimental</i> 36: 239-244.	3
De, L., A. Ceola and G. Magni (1984). "Viloxazine in the treatment of depressed elderly people. A double-blind trial against placebo." <i>Minerva Psichiatrica</i> 25: 127-130.	4
De, R., P. Rucci, M. Lodi, G. Ravaglia, P. Forti and V. Volterra (1998). "Fluoxetine and amitriptyline in elderly depressed patients. A 10-week, double-blind study on course of neurocognitive adverse events and depressive symptoms." <i>Arch Gerontol Geriatr</i> 27: 125-140.	3
De, V., J. Kummer, A. Agnoli, P. Gentili, A. Lorizio and R. Anand (1990). "Moclobemide compared with second-generation antidepressants in elderly people." <i>Acta Psychiatr Scand Suppl</i> 360: 64-66.	5
DeBattista, C., J. Belanoff, S. Glass, A. Khan, R. L. Horne, C. Blasey, L. L. Carpenter and G. Alva (2006). "Mifepristone versus placebo in the treatment of psychosis in patients with psychotic major depression." <i>Biol Psychiatry</i> 60: 1343-1349.	3

Dechant, K. L. and S. P. Clissold (1991). "Paroxetine. A review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in depressive illness." Drugs 41: 225-253.	3
Dehlin, O., G. Bjornsson and J. Lundstrom (1981). "Zimelidine to geriatric patients: A pharmacokinetic and clinical study." Acta Psychiatrica Scandinavica 63: 410-424.	3
Delle, C., P. Panzeri, S. Seripa, L. Trillo, M. Donnini and E. Vicario (1992). "Efficacy of moclobemide in a geriatric population with symptoms of depression and cognitive decline." Clin Neuropharmacol: 461.	3
Demyttenaere, K., H. F. Andersen and E. H. Reines (2008). "Impact of escitalopram treatment on Quality of Life Enjoyment and Satisfaction Questionnaire scores in major depressive disorder and generalized anxiety disorder." Int Clin Psychopharmacol 23: 276-286.	4
Dew, M. A., C. F. Reynolds, rd, B. Mulsant, E. Frank, P. R. Houck, S. Mazumdar, A. Begley and D. J. Kupfer (2001). "Initial recovery patterns may predict which maintenance therapies for depression will keep older adults well." J Affect Disord 65: 155-166.	3
Devanand, D. P. and S. P. Roose (2014). "Antidepressants in older adults: complexities, confounds and clinical effects." Am J Geriatr Psychiatry 22(1): 1-3.	7
Devanand, D. P., M. S. Nobler, J. Cheng, N. Turret, G. H. Pelton, S. P. Roose and H. A. Sackeim (2005). "Randomized, double-blind, placebo-controlled trial of fluoxetine treatment for elderly patients with dysthymic disorder." Am J Geriatr Psychiatry 13: 59-68.	4
Devanand, D. P., N. Juszczak, S. Nobler Mitchell, N. Turret, L. Fitzsimons, A. Sackeim Harold and P. Roose Steven (2004). "An Open Treatment Trial of Venlafaxine for Elderly Patients With Dysthymic Disorder." J Geriatr Psychiatry Neurol 17: 219-224.	9
Devos, D., K. Dujardin, I. Poirot, C. Moreau, O. Cottencin, P. Thomas, A. Destee, R. Bordet and L. Defebvre (2008). "Comparison of desipramine and citalopram treatments for depression in Parkinson's disease: a double-blind, randomized, placebo-controlled study." Mov Disord 23(6): 850-857.	4
Dierick, M., L. Ravizza, R. Realini and A. Martin (1996). "A double-blind comparison of venlafaxine and fluoxetine for treatment of major depression in outpatients." Prog Neuropsychopharmacol Biol Psychiatry 20(1): 57-71.	9
Dinan, T. G. (2001). "Efficacy and safety of weekly treatment with enteric-coated fluoxetine in patients with major depressive disorder." J Clin Psychiatry 62 Suppl 22: 48-52.	3
Dinan, T. G. and S. Barry (1989). "A comparison of electroconvulsive therapy with a combined lithium and tricyclic combination among depressed tricyclic nonresponders." Acta Psychiatr Scand 80(1): 97-100.	8
Distefano, A., L. Mariani, G. Tiralosi and G. Pavano (1984). "A new antidepressant for treatment of depressions in aged patients." Giornale di Gerontologia 32: 517-532.	11
Divyashree, M., Jayanthi, C.R. and Chandrashekhar, H., A comparative study of efficacy and safety of conventional versus newer antidepressants in patients with depressive episode in a tertiary care hospital, Journal of Chemical and Pharmaceutical Research, 6 (2014) 516-524.	7

Dobkin, R. D., M. Menza, K. L. Bienfait, M. Gara, H. Marin, M. H. Mark, A. Dicke and A. Troster (2010). "The impact of antidepressant treatment on cognitive functioning in depressed patients with Parkinson's disease." <i>J Neuropsychiatry Clin Neurosci</i> 22: 188-195.	4
Dobkin, R. D., M. Menza, K. L. Bienfait, M. Gara, H. Marin, M. H. Mark, A. Dicke and J. Friedman (2011). "Depression in Parkinson's disease: Symptom improvement and residual symptoms after acute pharmacologic management." <i>American Journal of Geriatric Psychiatry</i> 19(3): 222-229.	3
Dobscha, S. K., K. Corson, D. H. Hickam, N. A. Perrin, D. F. Kraemer and M. S. Gerrity (2006). "Depression decision support in primary care: a cluster randomized trial." <i>Ann Intern Med</i> 145(7): 477-487.	3
Dombrovski, A. Y. and B. H. Mulsant (2007). "The evidence for electroconvulsive therapy (ECT) in the treatment of severe late-life depression. ECT: the preferred treatment for severe depression in late life." <i>Int Psychogeriatr</i> 19(1): 10-14, 27-35; discussion 24-16.	9
Doogan, D. P. (1991). "Toleration and safety of sertraline: Experience worldwide." <i>Int Clin Psychopharmacol</i> 6: 47-56.	4
Doogan, D. P. and V. Caillard (1992). "Sertraline in the prevention of depression." <i>Br J Psychiatry</i> 160: 217-222.	7
Doriswamy, P. M., K. R. Krishnan, T. Oxman, L. R. Jenkyn, D. J. Coffey, T. Burt and C. M. Clary (2003). "Does antidepressant therapy improve cognition in elderly depressed patients?" <i>J Gerontol A Biol Sci Med Sci</i> 58(12): M1137-1144.	9
Doriswamy, P. M., Z. M. Khan, R. M. Donahue and N. E. Richard (2001). "Quality of life in geriatric depression: a comparison of remitters, partial responders, and nonresponders." <i>Am J Geriatr Psychiatry</i> 9: 423-428.	3
Dording, C. M., D. Mischoulon, I. Shyu, J. E. Alpert and G. I. Papakostas (2012). "SAME and sexual functioning." <i>Eur Psychiatry</i> 27: 451-454.	4
Doree, J. P., J. Des Rosiers, V. Lew, A. Gendron, R. Elie, E. Stip and S. V. Tourjman (2007). "Quetiapine augmentation of treatment-resistant depression: a comparison with lithium." <i>Curr Med Res Opin</i> 23(2): 333-341.	3
Drye, L. T., B. K. Martin, C. E. Frangakis, C. L. Meinert, J. E. Mintzer, C. A. Munro, A. P. Porsteinsson, P. V. Rabins, P. B. Rosenberg, L. S. Schneider, D. Weintraub, C. G. Lyketsos and D.-R. Group (2011). "Do treatment effects vary among differing baseline depression criteria in depression in Alzheimer's disease study +/- 2 (DIADS-2)?" <i>Int J Geriatr Psychiatry</i> 26(6): 573-583.	4
Dube, S., M. A. Dellva, M. Jones, W. Kielbasa, R. Padich, A. Saha and P. Rao (2010). "A study of the effects of LY2216684, a selective norepinephrine reuptake inhibitor, in the treatment of major depression." <i>J Psychiatr Res</i> 44: 356-363.	3
Dubini, A., M. Bosc and V. Polin (1997). "Noradrenaline-selective versus serotonin-selective antidepressant therapy: differential effects on social functioning." <i>J Psychopharmacol</i> 11: S17-23.	10
Duboff, E. A. (1993). "Long-term treatment of major depressive disorder with paroxetine." <i>J Clin Psychopharmacol</i> 13: 28S-33S.	3

Dumville, J. C., J. N. Miles, J. Porthouse, S. Cockayne, L. Saxon and C. King (2006). "Can vitamin D supplementation prevent winter-time blues? A randomised trial among older women." <i>J Nutr Health Aging</i> 10(2): 151-153.	7
Dunbar, G. C., J. L. Claghorn, A. Kiev, K. Rickels and W. T. Smith (1993). "A comparison of paroxetine and placebo in depressed outpatients." <i>Acta Psychiatr Scand</i> 87(5): 302-305.	4
Dunlop, S. R., B. E. Dornseif, J. F. Wernicke and J. H. Potvin (1990). "Pattern analysis shows beneficial effect of fluoxetine treatment in mild depression." <i>Psychopharmacol Bull</i> 26: 173-180.	4
Dunner, D. L., J. B. Cohn, T. Walshe, rd, C. K. Cohn, J. P. Feighner, R. R. Fieve, J. P. Halikas, J. T. Hartford, E. D. Hearst, E. C. Settle, Jr and et al. (1992). "Two combined, multicenter double-blind studies of paroxetine and doxepin in geriatric patients with major depression." <i>J Clin Psychiatry</i> 53 Suppl: 57-60.	3
Dunner, D. L., J. D. Amsterdam, R. C. Shelton, A. Loebel and S. J. Romano (2007). "Efficacy and tolerability of adjunctive ziprasidone in treatment-resistant depression: a randomized, open-label, pilot study." <i>J Clin Psychiatry</i> 68(7): 1071-1077.	3
Durham, L. K., S. M. Webb, P. M. Milos, C. M. Clary and A. B. Seymour (2004). "The serotonin transporter polymorphism, 5HTTLPR, is associated with a faster response time to sertraline in an elderly population with major depressive disorder." <i>Psychopharmacology (Berl)</i> 174: 525-529.	4
Edwards, D. R. (1998) "A double- blind, placebo-controlled trial of lithium augmentation of antidepressants in treatment-resistant depression in elderly patients." National Research Register.	9
Eiser, N., R. Harte, K. Spiros, C. Phillips and M. T. Isaac (2005). "Effect of treating depression on quality-of-life and exercise tolerance in severe COPD." <i>COPD</i> 2: 233-241.	3
Eklund, K., G. C. Dunbar, R. M. Pinder and K. Steffensen (1985). "Mianserin and imipramine in the treatment of elderly depressed patients." <i>Acta Psychiatrica Scandinavica</i> 72: 54-59.	9
Ekselius, L., F. Bengtsson and K. Von (2000). "Non-compliance with pharmacotherapy of depression is associated with a sensation seeking personality." <i>Int Clin Psychopharmacol</i> 15: 273-278.	3
Ekselius, L., L. von Knorring and G. Eberhard (1997). "A double-blind multicenter trial comparing sertraline and citalopram in patients with major depression treated in general practice." <i>Int Clin Psychopharmacol</i> 12(6): 323-331.	9
El-Khalili, N., M. Joyce, S. Atkinson, R. J. Buynak, C. Datto, P. Lindgren and H. Eriksson (2010). "Extended-release quetiapine fumarate (quetiapine XR) as adjunctive therapy in major depressive disorder (MDD) in patients with an inadequate response to ongoing antidepressant treatment: a multicentre, randomized, double-blind, placebo-controlled study." <i>Int J Neuropsychopharmacol</i> 13: 917-932.	10
Elsborg, L. (1991). "Safety aspects of treatment of the elderly with dementia and/or depression: Side effect profile of a specific 5-HT uptake inhibitor." <i>Nordisk Psykiatrisk Tidsskrift</i> : 29-35.	4
Entsuah, A. R., H. Huang and M. E. Thase (2001). "Response and remission rates in different subpopulations with major depressive disorder administered venlafaxine, selective serotonin reuptake inhibitors, or placebo." <i>Journal of Clinical Psychiatry</i> 62: 869-877.	3

Evans, M., M. Hammond, K. Wilson, M. Lye and J. Copeland (1997). "Placebo-controlled treatment trial of depression in elderly physically ill patients." <i>Int J Geriatr Psychiatry</i> 12: 817-824.	10
Fabre, L. F. (1990). "Buspirone in the management of major depression: a placebo-controlled comparison." <i>J Clin Psychiatry</i> 51 Suppl: 55-61.	3
Fabre, L. F., F. S. Abuzzahab, M. Amin, J. L. Claghorn, J. Mendels, W. M. Petrie, S. Dube and J. G. Small (1995). "Sertraline safety and efficacy in major depression: a double-blind fixed-dose comparison with placebo." <i>Biol Psychiatry</i> 38: 592-602.	3
Fabre, L. F., M. B. Scharf and T. M. Itil (1991). "Comparative efficacy and safety of nortriptyline and fluoxetine in the treatment of major depression: a clinical study." <i>J Clin Psychiatry</i> 52 Suppl: 62-67.	3
Fairbairn, A. F., K. George and T. Dorman (1989). "Lofepramine versus dothiepin in the treatment of depression in elderly patients." <i>Br J Clin Pract</i> 43: 55-60.	3
Fairweather, D. B., J. S. Kerr, D. A. Harrison, C. A. Moon and I. Hindmarch (1993). "A double blind comparison of the effects of fluoxetine and amitriptyline on cognitive function in elderly depressed patients." <i>Hum Psychopharmacol</i> 8(1): 41-47.	9
Falk, W. E., J. F. Rosenbaum, M. W. Otto, P. M. Zusky, J. B. Weilburg and R. A. Nixon (1989). "Fluoxetine versus trazodone in depressed geriatric patients." <i>J Geriatr Psychiatry Neurol</i> 2: 208-214.	4
Fang, Y., C. Yuan, Y. Xu, J. Chen, Z. Wu, L. Cao, Z. Yi, W. Hong, Y. Wang, K. Jiang, X. Cui, J. R. Calabrese and K. Gao (2011). "A pilot study of the efficacy and safety of paroxetine augmented with risperidone, valproate, buspirone, trazodone, or thyroid hormone in adult chinese patients with treatment-resistant major depression." <i>J Clin Psychopharmacol</i> 31: 638-642.	4
Fava, M., A. J. Rush, J. E. Alpert, G. K. Balasubramani, S. R. Wisniewski, C. N. Carmin, M. M. Biggs, S. Zisook, A. Leuchter, R. Howland, D. Warden and M. H. Trivedi (2008). "Difference in treatment outcome in outpatients with anxious versus nonanxious depression: a STAR*D report." <i>Am J Psychiatry</i> 165(3): 342-351.	4
Fava, M., C. H. Mallinckrodt, M. J. Detke, J. G. Watkin and M. M. Wohlrreich (2004). "The effect of duloxetine on painful physical symptoms in depressed patients: Do improvements in these symptoms result in higher remission rates?" <i>Journal of Clinical Psychiatry</i> 65(4): 521-530.	3
Fava, M., D. Mischoulon, D. Iosifescu, J. Witte, M. Pencina, M. Flynn, L. Harper, M. Levy, K. Rickels and M. Pollack (2012). "A double-blind, placebo-controlled study of aripiprazole adjunctive to antidepressant therapy among depressed outpatients with inadequate response to prior antidepressant therapy (ADAPT-A Study)." <i>Psychother Psychosom</i> 81(2): 87-97.	4
Fava, M., M. E. Schmidt, S. Zhang, J. Gonzales, N. J. Raute and R. Judge (2002). "Treatment approaches to major depressive disorder relapse. Part 2: reinitiation of antidepressant treatment." <i>Psychother Psychosom</i> 71: 195-199.	4
Fawcett, J. and R. L. Barkin (1998). "Review of the results from clinical studies on the efficacy, safety and tolerability of mirtazapine for the treatment of patients with major depression." <i>J Affect Disord</i> 51: 267-285.	4

Fawcett, J., R. N. Marcus, S. F. Anton, K. O'Brien and U. Schwiderski (1995). "Response of anxiety and agitation symptoms during nefazodone treatment of major depression." Journal of Clinical Psychiatry 56: 37-42.	4
Feiger, A. D., K. A. Tourian, G. R. Rosas and S. K. Padmanabhan (2009). "A placebo-controlled study evaluating the efficacy and safety of flexible-dose desvenlafaxine treatment in outpatients with major depressive disorder." CNS Spectr 14: 41-50.	4
Feighner, J. P. and J. B. Cohn (1985). "Double-blind comparative trials of fluoxetine and doxepin in geriatric patients with major depressive disorder." Journal of Clinical Psychiatry 46(3 II): 20-25.	3
Feighner, J. P., W. F. Boyer, C. H. Meredith and G. Hendrickson (1988). "An overview of fluoxetine in geriatric depression." Br-J-Psychiatry-Suppl: 105-108.	3
Feighner, J. P., W. F. Boyer, G. G. Hendrickson, R. A. Pambakian and V. S. Doroski (1990). "A controlled trial of adinazolam versus desipramine in geriatric depression." Int Clin Psychopharmacol 5(3): 227-232.	4
Felmet, K., S. Zisook and J. W. Kasckow (2011). "Elderly patients with schizophrenia and depression: Diagnosis and treatment." Clinical Schizophrenia and Related Psychoses 4: 239-250E.	4
Feng, Y., B. G. Pollock, R. E. Ferrell, M. A. Kimak, C. F. Reynolds, rd and R. R. Bies (2006). "Paroxetine: population pharmacokinetic analysis in late-life depression using sparse concentration sampling." Br J Clin Pharmacol 61: 558-569.	4
Ferguson, J. M., J. Mendels and G. E. Schwart (2002). "Effects of reboxetine on Hamilton Depression Rating Scale factors from randomized, placebo-controlled trials in major depression." Int Clin Psychopharmacol 17: 45-51.	9
Fields, C., L. Drye, V. Vaidya, C. Lyketsos and A. R. Group (2012). "Celecoxib or naproxen treatment does not benefit depressive symptoms in persons age 70 and older: findings from a randomized controlled trial." Am J Geriatr Psychiatry 20(6): 505-513.	4
Fisch, C. and S. B. Knoebel (1992). "Electrocardiographic findings in sertraline depression trials." Drug Investigation 4: 305-312.	9
Fischer-Cornelissen, K. A. and P. Berchier (1980). "Geriatric patients with depressive disturbances. Double-blind multicenter study and long-term trial of pizotifen, placebo, doxepine and cinnarizine." Arzneimittelforschung: 1216-1217.	11
Fitch, K., F. J. Molnar, B. Power, D. Wilkins and M. Man-Son-Hing (2005). "Antidepressant use in older people: family physicians' knowledge, attitudes, and practices." Can Fam Physician 51: 80-81.	5
Flint, A. J., A. Iaboni, B. H. Mulsant, A. J. Rothschild, E. M. Whyte, B. S. Meyers and S. Group Stop-Pd (2013). "Effect of Sertraline on Risk of Falling in Older Adults with Psychotic Depression on Olanzapine: Results of a Randomized Placebo-Controlled Trial." Am J Geriatr Psychiatry.	4
Florea, I., N. Danchenko, H. Loft, B. Rive, S. Pendlebury and L. Abetz (2013). "The effect of vortioxetine (LU AA21004) on health-related quality of life (HRQOL) in patients with major depressive disorder (MDD)." Value in Health 16: A65.	6

Ford, A. H., L. Flicker, J. Thomas, P. Norman, K. Jamrozik and O. P. Almeida (2008). "Vitamins B12, B6, and folic acid for onset of depressive symptoms in older men: results from a 2-year placebo-controlled randomized trial." <i>J Clin Psychiatry</i> 69: 1203-1209.	4
Ford, A. H., L. Flicker, K. McCaul, F. van Bockxmeer, S. Hegarty, V. Hirani, S. Fenner and O. P. Almeida (2010). "The B-VITAGE trial: A randomized trial of homocysteine lowering treatment of depression in later life." <i>Trials</i> 11.	3
Forlenza, O. V., A. Stoppe Junior, E. S. Hirata and R. C. Ferreira (2000). "Antidepressant efficacy of sertraline and imipramine for the treatment of major depression in elderly outpatients." <i>Sao Paulo Med J</i> 118(4): 99-104.	10
Forlenza, O. V., O. P. Almeida, A. Stoppe, Jr., E. S. Hirata and R. C. R. Ferreira (2001). "Antidepressant efficacy and safety of low-dose sertraline and standard-dose imipramine for the treatment of depression in older adults: results from a double-blind, randomized, controlled clinical trial." <i>Int Psychogeriatr</i> 13(1): 75-84.	7 och 8
Fortner, M. R., K. Brown, I. M. Varia, K. R. Gersing, C. O'Connor and P. M. Doraiswamy (1999). "Effect of Bupropion SR on the Quality of Life of Elderly Depressed Patients With Comorbid Medical Disorders." <i>Prim Care Companion J Clin Psychiatry</i> 1(6): 174-179.	3
Frederiksen, S. O., G. d'Elia and B. O. Bengtsson (1985). "ACTH 4-9 analogue (org 2766) in depressed elderly patients. I Effect on depressed mood." <i>Acta Psychiatrica Scandinavica</i> 72(4): 341-348.	4
Frederiksen, S. O., G. d'Elia and B. O. Bengtsson (1985). "ACTH 4-9 analogue (ORG 2766) in depressed elderly patients. II. Effect on memory and vigilance." <i>Acta Psychiatrica Scandinavica</i> 72(4): 349-357.	4
Freed, E., R. Goldney, T. Lambert, J. Tiller and R. Johnston (1999). "A double-blind, multicentre study to assess the tolerability and efficacy of paroxetine compared with amitriptyline in the treatment of depressed patients in Australian general practice." <i>Aust N Z J Psychiatry</i> 33(3): 416-421.	3
Fregni, F., C. R. Ono, C. M. Santos, F. Bermpohl, C. Buchpiguel, E. R. Barbosa, M. A. Marcolin, A. Pascual-Leone and K. D. Valente (2006). "Effects of antidepressant treatment with rTMS and fluoxetine on brain perfusion in PD." <i>Neurology</i> 66: 1629-1637.	9
Friedman, E. S., L. L. Davis, S. Zisook, S. R. Wisniewski, M. H. Trivedi, M. Fava, A. J. Rush and C.-M. S. Team (2012). "Baseline depression severity as a predictor of single and combination antidepressant treatment outcome: results from the CO-MED trial." <i>Eur Neuropsychopharmacol</i> 22(3): 183-199.	3
Fu, Y. X., H. Q. Meng and Q. H. Luo (2006). "Interventional effect of fluoxetine on patients with acute myocardial infarction accompanied by anxiety and depression." <i>Chinese Journal of Clinical Rehabilitation</i> 10: 16-17.	11
Fuchs, A., U. Hehnke, C. Erhart, C. Schell, B. Pramshohler, B. Danninger and F. Schautzer (1993). "Video rating analysis of effect of maprotiline in patients with dementia and depression." <i>Pharmacopsychiatry</i> 26: 37-41.	3
Galecki, P., A. Florkowski, T. Pietras, I. Kolodziejska, T. Nowakowski, A. Pawelczyk and K. Zboralski (2004). "Efficiency and safety of citalopram and venlafaxine in treatment of depressive disorders in elderly patients." <i>Polski Merkuriusz Lekarski</i> : 621-624.	11

Gao, A. M., X. B. Xue and S. T. Qiao (2004) "A clinical control study of paroxetine and venlafaxine in the treatment of depression in elderly." Chinese Clinical New Medicine, 506-507.	3
Garakani, A., J. M. Martinez, S. Marcus, J. Weaver, K. Rickels, M. Fava and J. Hirschowitz (2008). "A randomized, double-blind, and placebo-controlled trial of quetiapine augmentation of fluoxetine in major depressive disorder." <i>Int Clin Psychopharmacol</i> 23(5): 269-275.	3
Gariballa, S. (2011). "Testing homocysteine-induced neurotransmitter deficiency, and depression of mood hypothesis in clinical practice." <i>Age Ageing</i> 40: 702-705.	9
Garlow, S. J., B. Kinkead, M. E. Thase, L. L. Judd, A. J. Rush, K. A. Yonkers, D. J. Kupfer, E. Frank, P. J. Schettler and M. H. Rapaport (2013). "Fluoxetine increases suicide ideation less than placebo during treatment of adults with minor depressive disorder." <i>J Psychiatr Res</i> 47(9): 1199-1203.	3
Gastpar, M., A. Singer and K. Zeller (2005). "Efficacy and tolerability of hypericum extract STW3 in long-term treatment with a once-daily dosage in comparison with sertraline." <i>Pharmacopsychiatry</i> 38: 78-86.	3
Gaynes, B. N., S. B. Dusetzina, A. R. Ellis, R. A. Hansen, J. F. Farley, W. C. Miller and T. Sturmer (2012). "Treating depression after initial treatment failure: directly comparing switch and augmenting strategies in STAR*D." <i>J Clin Psychopharmacol</i> 32: 114-119.	4
Gaynor, P. J., M. Gopal, W. Zheng, J. M. Martinez, M. J. Robinson and L. B. Marangell (2011). "A randomized placebo-controlled trial of duloxetine in patients with major depressive disorder and associated painful physical symptoms." <i>Curr Med Res Opin</i> 27: 1849-1858.	4
Gaynor, P. J., M. Gopal, W. Zheng, J. M. Martinez, M. J. Robinson, D. Hann and L. B. Marangell (2011). "Duloxetine versus placebo in the treatment of major depressive disorder and associated painful physical symptoms: a replication study." <i>Curr Med Res Opin</i> 27(10): 1859-1867.	9
Gentili, P. (1974). "Bromazepam in treatment of dysthymic symptoms in the elderly." <i>Giornale di Gerontologia</i> 22(7): 457-466.	11
Gentili, P., F. De Maria and M. De Vanna (1984). "Diclofensine in the treatment of depressed geriatric hospitalized patients: A placebo-controlled double-blind trial." <i>Current Therapeutic Research - Clinical and Experimental</i> 35(3): 386-395.	3
Georgotas, A. and R. E. McCue (1989) "The additional benefit of extending an antidepressant trial past seven weeks in the depressed elderly." <i>Int J Geriatr Psychiatry</i> , 191-195.	9
Georgotas, A. and R. E. McCue (1989). "Relapse of depressed patients after effective continuation therapy." <i>J Affect Disord</i> 17: 159-164.	3
Georgotas, A., B. Reisberg and S. Ferris (1983). "First results on the effects of MAO inhibition on cognitive functioning in elderly depressed patients." <i>Arch Gerontol Geriatr</i> 2: 249-254.	9
Georgotas, A., P. Stokes and R. E. McCue (1986). "The usefulness of DST in predicting response to antidepressants: A placebo-controlled study." <i>J Affect Disord</i> 11: 21-28.	3
Georgotas, A., R. E. McCue and T. B. Cooper (1989). "A placebo-controlled comparison of nortriptyline and phenelzine in maintenance therapy of elderly depressed patients." <i>Arch Gen Psychiatry</i> 46: 783-786.	7

Georgotas, A., R. E. McCue and W. Hapworth (1986). "Comparative efficacy and safety of MAOIs versus TCAs in treating depression in the elderly." <i>Biol Psychiatry</i> 21(12): 1155-1166.	9
Georgotas, A., R. E. McCue, B. Reisberg, S. H. Ferris, N. Nagachandran, I. Chang and P. Mir (1989). "The effects of mood changes and antidepressants on the cognitive capacity of elderly depressed patients." <i>Int Psychogeriatr</i> 1: 135-143.	4
Georgotas, A., R. E. McCue, E. Friedman and T. B. Cooper (1987). "A placebo-controlled comparison of the effect of nortriptyline and phenelzine on orthostatic hypotension in elderly depressed patients." <i>J Clin Psychopharmacol</i> : 413-416.	3
Georgotas, A., R. E. McCue, T. B. Cooper, N. Nagachandran and I. Chang (1988). "How effective and safe is continuation therapy in elderly depressed patients? Factors affecting relapse rate." <i>Arch Gen Psychiatry</i> 45: 929-932.	3
Geretsegger, C., F. Bohmer and M. Ludwig (1994). "Paroxetine in the elderly depressed patient: randomized comparison with fluoxetine of efficacy, cognitive and behavioural effects." <i>Int Clin Psychopharmacol</i> 9: 25-29.	9
Gerner, R. H. (1984). "Antidepressant selection in the elderly." <i>Psychosomatics</i> 25(7): 528-535.	3
Gerner, R., W. Estabrook, J. Steuer and L. Jarvik (1980). "Treatment of geriatric depression with trazodone, imipramine, and placebo: A double-blind study." <i>Journal of Clinical Psychiatry</i> 41: 216-220.	6
Gerner, R., W. Estabrook, J. Steuer, L. Waltuch, P. Kakkar and L. Jarvik (1980). "A placebo-controlled double-blind study of imipramine and trazodone in geriatric depression." <i>Psychopathology in the Aged</i> : 167-182.	3
Gershon, S., J. Mann, R. Newton and B. J. Gunther (1981). "Evaluation of trazodone in the treatment of endogenous depression: Results of a multicenter double-blind study." <i>J Clin Psychopharmacol</i> 1: 39S-44S.	3
Gex-Fabry, M., F. Jermann, M. Kosel, M. F. Rossier, L. Van der, G. Bertschy, G. Bondolfi and J. M. Aubry (2012). "Salivary cortisol profiles in patients remitted from recurrent depression: one-year follow-up of a mindfulness-based cognitive therapy trial." <i>J Psychiatr Res</i> 46: 80-86.	4
Ghose, K. (1997). "Tolerance and side-effects of paroxetine in elderly depressed patients." <i>Arch Gerontol Geriatr</i> 24: 35-45.	4
Gibiino, S., E. Mori, D. De Ronchi and A. Serretti (2013). "Potential benefits of slow titration of paroxetine treatment in an elderly population: eight-week results from a naturalistic setting." <i>J Clin Psychopharmacol</i> 33(4): 565-569.	9
Goldstein, D. J., S. H. Hamilton, D. N. Masica, C. M. Beasley and Jr (1997). "Fluoxetine in medically stable, depressed geriatric patients: effects on weight." <i>J Clin Psychopharmacol</i> 17: 365-369.	3
Goldstein, S. E., F. Birnbom and R. Laliberte (1982). "Nomifensine in the treatment of depressed geriatric patients." <i>Journal of Clinical Psychiatry</i> 43: 287-289.	3
Goodwin, G. M., R. Emsley, S. Rembry, F. Rouillon and G. Agomelatine Study (2009). "Agomelatine prevents relapse in patients with major depressive disorder without evidence of a discontinuation syndrome: a 24-week randomized, double-blind, placebo-controlled trial." <i>J Clin Psychiatry</i> 70(8): 1128-1137.	4

Gopalakrishnan, R., J. Ross, C. O'Brien and D. Oslin (2009). "Course of late-life depression with alcoholism following combination therapy." <i>J Stud Alcohol Drugs</i> 70(2): 237-241.	3
Gottfries, C. G. (1996). "Scandinavian experience with citalopram in the elderly." <i>Int Clin Psychopharmacol</i> 11: 41-44.	3
Goveas, J. S., P. E. Hogan, J. M. Kotchen, J. W. Smoller, N. L. Denburg, J. E. Manson, A. Tummala, W. J. Mysiw, J. K. Ockene, N. F. Woods, M. A. Espeland and S. Wassertheil-Smoller (2012). "Depressive symptoms, antidepressant use, and future cognitive health in postmenopausal women: the Women's Health Initiative Memory Study." <i>Int Psychogeriatr</i> 24(8): 1252-1264.	3 och 9
Gram, L. F. and G. Danish University Antidepressant (2008). "Acute and continuation therapy in unipolar depression: observations from the run-in phase of a maintenance trial." <i>Acta Psychiatr Scand</i> 118(2): 123-129.	4
Green, T. D., C. F. Reynolds, 3rd, B. H. Mulsant, B. G. Pollock, M. D. Miller, P. R. Houck, S. Mazumdar, M. A. Dew and D. J. Kupfer (1999). "Accelerating antidepressant response in geriatric depression: a post hoc comparison of combined sleep deprivation and paroxetine versus monotherapy with paroxetine, nortriptyline, or placebo." <i>J Geriatr Psychiatry Neurol</i> 12(2): 67-71.	3
Greenlee, A., J. F. Karp, M. A. Dew, P. Houck, C. Andreeescu and C. F. Reynolds, 3rd (2010). "Anxiety impairs depression remission in partial responders during extended treatment in late-life." <i>Depress Anxiety</i> 27(5): 451-456.	3
Greil, W., W. Ludwig-Mayerhofer, N. Erazo, R. R. Engel, A. Czernik, H. Giedke, B. Muller-Oerlinghausen, M. Osterheider, G. A. Rudolf, H. Sauer, J. Tegeler and T. Wetterling (1996). "Comparative efficacy of lithium and amitriptyline in the maintenance treatment of recurrent unipolar depression: a randomised study." <i>J Affect Disord</i> 40(3): 179-190.	3
Greist, J., R. K. McNamara, C. H. Mallinckrodt, J. N. Rayamajhi and J. Raskin (2004). "Incidence and duration of antidepressant-induced nausea: duloxetine compared with paroxetine and fluoxetine." <i>Clin Ther</i> 26: 1446-1455.	3
Grunebaum, M. F., J. G. Keilp, S. P. Ellis, K. Sudol, N. Bauer, A. K. Burke, M. A. Oquendo and J. J. Mann (2013). "SSRI versus bupropion effects on symptom clusters in suicidal depression: post hoc analysis of a randomized clinical trial." <i>J Clin Psychiatry</i> 74(9): 872-879.	3
Gu, Y., T. Jiang, J.-B. Guo, G. Wang, Z.-J. Yu, B.-S. Jiang, B. Zhang and W.-Y. Ma (2010). "Efficacy and safety of escitalopram in elderly patients with major depression: A randomized and control study." <i>Chinese Mental Health Journal</i> 24: 445-449.	11
Gual, A., M. Balcells, M. Torres, M. Madrigal, T. Diez and L. Serrano (2003). "Sertraline for the prevention of relapse in detoxicated alcohol dependent patients with a comorbid depressive disorder: a randomized controlled trial." <i>Alcohol Alcohol</i> 38: 619-625.	9
Guelfi, J. D., C. White, D. Hackett, J. Y. Guichoux and G. Magni (1995). "Effectiveness of venlafaxine in patients hospitalized for major depression and melancholia." <i>J Clin Psychiatry</i> 56: 450-458.	9
Guelfi, J. D., M. Ansseau, L. Timmerman, S. Korsgaard and G. Mirtazapine-Venlafaxine Study (2001). "Mirtazapine versus venlafaxine in hospitalized severely depressed patients with melancholic features." <i>J Clin Psychopharmacol</i> 21(4): 425-431.	3

Guelfi, J. D., N. Strub and H. Loft (2000). "Efficacy of intravenous citalopram compared with oral citalopram for severe depression. Safety and efficacy data from a double-blind, double-dummy trial." <i>J Affect Disord</i> 58: 201-209.	3
Guerin, R. and P. Erny (1968). "The effect of Noveril 80 in elderly subjects. Clinical trials with Noveril 80 in the treatment of depressive conditions in the aged. [French]." <i>Gazette medicale de France</i> : 1286-1287.	11
Guillibert, E., Y. Pelicier, J. C. Archambault, J. P. Chabannes, G. Clerc, M. Desvilles, M. Guibert, R. Pagot, J. L. Poisat and Y. Thobie (1989). "A double-blind, multicentre study of paroxetine versus clomipramine in depressed elderly patients." <i>Acta Psychiatrica Scandinavica, Supplement</i> 80(350): 132-134.	3
Gunning, F. M., J. Cheng, C. F. Murphy, D. Kanellopoulos, J. Acuna, M. J. Hoptman, S. Klimstra, S. Morimoto, J. Weinberg and G. S. Alexopoulos (2009). "Anterior cingulate cortical volumes and treatment remission of geriatric depression." <i>Int J Geriatr Psychiatry</i> 24(8): 829-836.	9
Gunning-Dixon, F. M., M. Walton, J. Cheng, J. Acuna, S. Klimstra, M. E. Zimmerman, A. M. Brickman, M. J. Hoptman, R. C. Young and G. S. Alexopoulos (2010). "MRI signal hyperintensities and treatment remission of geriatric depression." <i>J Affect Disord</i> 126: 395-401.	3
Gunning-Dixon, F., J. Cheng, J. Acuna, D. Kanellopoulos, C. Murphy, M. Hoptman, J. Weinberg and G. S. Alexopoulos (2009). "Anterior cingulate regional volumes and treatment remission of geriatric depression." <i>American Journal of Geriatric Psychiatry</i> 17: A63.	9
Gwirtsman, H. E., S. Ahles and A. Halaris (1983). "Therapeutic superiority of maprotiline versus doxepin in geriatric depression." <i>Journal of Clinical Psychiatry</i> 44: 449-453.	3
Habra, M. E., B. Baker, N. Frasure-Smith, J. R. Swenson, D. Koszycki, G. Butler, Z. van, T. L and F. Lesperance (2010). "First episode of major depressive disorder and vascular factors in coronary artery disease patients: Baseline characteristics and response to antidepressant treatment in the CREATE trial." <i>J Psychosom Res</i> 69: 133-141.	6
Hahm, W. (2010). "An open label study of mirtazapine in the treatment of depression with Alzheimer's disease." <i>European Neuropsychopharmacology</i> 20: S552.	3
Haight, T. and H. Cheng (2013). "Antidepressant treatment for older adults with dementia and depression: Re-visit the evidence from randomized controlled trials (RCT)." <i>J Am Geriatr Soc</i> 61: S224.	3
Hale, A., R. M. Corral, C. Mencacci, J. S. Ruiz, C. A. Severo and V. Gentil (2010). "Superior antidepressant efficacy results of agomelatine versus fluoxetine in severe MDD patients: a randomized, double-blind study." <i>Int Clin Psychopharmacol</i> 25: 305-314.	6
Halikas, J. A. (1995). "Org 3770 (mirtazapine) versus trazodone: A placebo controlled trial in depressed elderly patients." <i>Hum Psychopharmacol</i> 10: S125-S133.	3
Han, L., J. McCusker, M. Cole, R. Capek and M. Abrahamowicz (2011). "Antidepressant use and cognitive functioning in older medical patients with major or minor depression: a prospective cohort study with database linkage." <i>J Clin Psychopharmacol</i> 31(4): 429-435.	3

Hansen, R. A., S. B. Dusetzina, A. R. Ellis, T. Sturmer, J. F. Farley and B. N. Gaynes (2012). "Risk of adverse events in treatment-resistant depression: propensity-score-matched comparison of antidepressant augment and switch strategies." <i>Gen Hosp Psychiatry</i> 34: 192-200.	7
Hardy, B. G., K. I. Shulman and C. Zuccheri (1997). "Gradual discontinuation of lithium augmentation in elderly patients with unipolar depression." <i>J Clin Psychopharmacol</i> 17(1): 22-26.	6
Havle, N., Y. Altunkaynak, C. Dayan, M. C. Ilinem and B. Arpacı (2010). "Escitalopram in treatment of post-stroke depression." <i>Klinik Psikofarmakoloji Bulteni</i> 20: 74-78.	11
Hayes, R. L., R. H. Gerner and L. Fairbanks (1983). "ECG findings in geriatric depressives given trazodone, placebo, or imipramine." <i>Journal of Clinical Psychiatry</i> 44(5 I): 180-183.	4
He, P., Y. Kong and L. Z. Xu (2004). "Randomized controlled observation on the effect of early application of fluoxetine in preventing depression after stroke." <i>Chinese Journal of Clinical Rehabilitation</i> 8: 6016-6017.	11
Hebenstreit, G. F., K. Fellerer, R. Zochling, A. Zentz and G. C. Dunbar (1989). "A pharmacokinetic dose titration study in adult and elderly depressed patients." <i>Acta Psychiatrica Scandinavica, Supplement</i> 80: 81-84.	6
Hebenstreit, G. F., M. Loidl, U. Baumhackl, B. Gallhofer, C. Geretsegger, E. Radmayr, M. Saletu, W. Schony and M. Stabl (1990). "Efficacy and safety of moclobemide compared with imipramine in the treatment of major depressive disorder. Double-blind multicenter study, Austria." <i>J Neural Transm Suppl</i> 32: 177-184.	4
Heiligenstein, J. H., J. E. Ware, Jr., K. M. Beusterien, P. J. Roback, C. Andrejasich and G. D. Tollefson (1995). "Acute effects of fluoxetine versus placebo on functional health and well-being in late-life depression." <i>Int Psychogeriatr</i> 7 Suppl: 125-137.	4
Hellerstein, D. J., S. A. Little, L. W. Samstag, S. Batchelder, J. C. Muran, M. Fedak, D. Kreditor, R. N. Rosenthal and A. Winston (2001). "Adding group psychotherapy to medication treatment in dysthymia: a randomized prospective pilot study." <i>J Psychother Pract Res</i> 10(2): 93-103.	3
Heras, P., K. Kritikos, A. Hatzopoulos, N. Kritikos, V. Heras and D. Mitsibounas (2013). "The role of paroxetine in fatigue and depression of patients under chemotherapeutic treatment." <i>Am J Ther</i> 20(3): 254-256.	3
Hernandez, C. R., G. S. Smith, P. R. Houck, B. G. Pollock, B. Mulsant and M. A. Dew (2000) "The clinical response to total sleep deprivation and recovery sleep in geriatric depression: potential indicators of antidepressant treatment outcome." <i>Psychiatry Res</i> , 41-49.	4
Heun, R., Corral, R.M., Ahokas, A., Nicolini, H., Teixeira, J.M. and Dehelean, P., Efficacy of agomelatine in more anxious elderly depressed patients. A randomized, double-blind study vs placebo, European Psychiatry, 28 (2013). Accession:	9
Holtzheimer, P. E., 3rd, T. W. Meeks, M. E. Kelley, M. Mufti, R. Young, K. McWhorter, N. Vito, R. Chismar, S. Quinn, S. Dey, E. H. Byrd and W. M. McDonald (2008). "A double blind, placebo-controlled pilot study of galantamine augmentation of antidepressant treatment in older adults with major depression." <i>Int J Geriatr Psychiatry</i> 23(6): 625-631.	3

Honig, A., A. M. Kuyper, A. H. Schene, M. van, P. J, J. de, D. M. Tulner, A. Schins, H. J. Crijns, P. M. Kuijpers, H. Vossen, R. Lousberg, J. Ormel and M.-I. investigators (2007). "Treatment of post-myocardial infarction depressive disorder: a randomized, placebo-controlled trial with mirtazapine." <i>Psychosom Med</i> 69: 606-613.	3
Hoog, S., R. Tepner, M. E. Nilsson, S. Romano and J. S. Kennedy (1999). "Changes in anxiety, agitation, and insomnia during treatment of depression for patients age 55 years and older: Analysis from fluoxetine double-blind, placebo-controlled trials." <i>International Journal of Geriatric Psychopharmacology</i> 2(1): 33-39.	3
Hostmaelingen, H. J., O. Asskilt, S. G. Austad, J. Fjellheim, E. A. Hostmaelingen, P. H. Kristiansen, T. I. Olsen, T. Skotte and E. Ofsti (1989). "Primary care treatment of depression in the elderly: a double-blind, multi-centre study of flupenthixol ('Fluanxol') and sustained-release amitriptyline." <i>Curr Med Res Opin</i> 11: 593-599.	3
Hoyberg, O. J., B. Maragakis, J. Mullin, D. Norum, E. Stordall, P. Ekdahl, E. Ose, K. M. Moksnes and C. Sennef (1996). "A double-blind multicentre comparison of mirtazapine and amitriptyline in elderly depressed patients." <i>Acta Psychiatr Scand</i> 93(3): 184-190.	6
Huang, L. L. (2005). "Comparison of therapeutic effects and side effects between fluoxetine hydrochloride and clomipramine in patients with vascular depression." <i>Chinese Journal of Clinical Rehabilitation</i> 9(24): 226-228.	3
Huezo-Diaz, P., R. Uher, R. Smith, M. Rietschel, N. Henigsberg, A. Marusic, O. Mors, W. Maier, J. Hauser, D. Souery, A. Placentino, A. Zobel, E. R. Larsen, P. M. Czerski, B. Gupta, F. Hoda, N. Perroud, A. Farmer, I. Craig, K. J. Aitchison and P. McGuffin (2009). "Moderation of antidepressant response by the serotonin transporter gene." <i>Br J Psychiatry</i> 195: 30-38.	3
Hutchinson, D. R., S. Tong, C. A. Moon, M. Vince and A. Clarke (1992). "Paroxetine in the treatment of elderly depressed patients in general practice: a double-blind comparison with amitriptyline." <i>Int Clin Psychopharmacol</i> 6 Suppl 4: 43-51.	10
Hwang, J. P., C. H. Yang and S. J. Tsai (2004). "Comparison study of venlafaxine and paroxetine for the treatment of depression in elderly Chinese inpatients." <i>Int J Geriatr Psychiatry</i> 19(2): 189-190.	3
Imamura, K., N. Okayasu and T. Nagatsu (2011). "The relationship between depression and regional cerebral blood flow in Parkinson's disease and the effect of selegiline treatment." <i>Acta Neurol Scand</i> 124: 28-39.	3
Iovieno, N., N. van, A. Clain, L. Baer and A. A. Nierenberg (2011). "Residual symptoms after remission of major depressive disorder with fluoxetine and risk of relapse." <i>Depress Anxiety</i> 28: 137-144.	3
Iwata, N., K. A. Tourian, E. Hwang, L. Mele and C. Vialet (2013). "Efficacy and safety of desvenlafaxine 25 and 50 Mg/Day in a randomized, placebo-controlled study of depressed outpatients." <i>Journal of Psychiatric Practice</i> 19: 5-14.	3
Jain, R., R. Mahableshwarkar Atul, L. Jacobsen Paula, Y. Chen and E. Thase Michael (2013). "A randomized, double-blind, placebo-controlled 6-wk trial of the efficacy and tolerability of 5 mg vortioxetine in adults with major depressive disorder." <i>International Journal of Neuropsychopharmacology</i> 16: 313-321.	3
Jainer, A. K., N. Soni and D. Onalaja (2003). "Re: Prophylactic therapy with lithium in elderly patients with Unipolar Major Depression." <i>Int J Geriatr Psychiatry</i> 18(4): 353-354; author reply 354.	9

Jansen, W., G. W. Brückner, S. Henauer and L. M. Omer (1982) "[Clinical double blind comparison of diclofensine and placebo in geriatric patients with depressive syndromes]." Pharmacopsychiatria, 205-209 DOI: 10.1055/s-2007-1019539.	11
Januel, D., M. F. Poirier, F. D'Alche-Biree, M. Dib and J. P. Olie (2003). "Multicenter double-blind randomized parallel-group clinical trial of efficacy of the combination clomipramine (150 mg/day) plus lithium carbonate (750 mg/day) versus clomipramine (150 mg/day) plus placebo in the treatment of unipolar major depression." J Affect Disord 76(1-3): 191-200.	3
Jefferson, J. W., A. J. Rush, J. C. Nelson, S. A. VanMeter, A. Krishen, K. D. Hampton, D. S. Wightman and J. G. Modell (2006). "Extended-release bupropion for patients with major depressive disorder presenting with symptoms of reduced energy, pleasure, and interest: findings from a randomized, double-blind, placebo-controlled study." J Clin Psychiatry 67: 865-873.	4
Jensen, H. V., K. Olafsson, L. Lykke-Olesen, A. Mork, A. Geisler and P. Arup (1992). "Combining nortriptyline and lithium in elderly depressed patients: A double-blind, placebo-controlled study." Lithium 3(4): 259-262.	10
Jessef, H. J., I. Jessef and G. Wegner (1981). "Treatment of elderly depressive patients. Lofepramin and amitriptyline under double-blind conditions." Zeitschrift für Allgemeinmedizin: 784-788.	11
Jessel, H. J., I. Jessel and G. Wegener (1981). "Therapy of elderly depressive patients. Lofepramine and amitriptyline under double-blind conditions." Zeitschrift für Allgemeinmedizin: 784-788.	3
Ji, J. L., W. J. Liu, N. Zhang, Z. Q. Chen, A. L. Zheng, Q. Y. Mei, J. Y. Pan, Z. X. Zhao, M. Tao, Y. P. Wang and J. Wei (2007). "Effects of paroxetine with or without zolpidem on depression with insomnia: A multi-center randomized comparative study." National Medical Journal of China 87: 1585-1589.	11
Jia, W., X. L. Zhang, D. B. Zhang and M. Y. Liu (2005). "Effect of early intervention on recovery of motor function and recurrent stroke in patients with post-stroke depression." Chinese Journal of Clinical Rehabilitation 9(12): 4-5.	11
Jiang, W., H. Oken, M. Fiuzat, L. K. Shaw, C. Martsberger, M. Kuchibhatla, R. Kaddurah-Daouk, D. C. Steffens, R. Baillie, M. Cuffe, R. Krishnan, C. O'Connor and S.-C. Investigators (2012). "Plasma omega-3 polyunsaturated fatty acids and survival in patients with chronic heart failure and major depressive disorder." J Cardiovasc Transl Res 5: 92-99.	3
Jindal, R. D., E. S. Friedman, S. R. Berman, A. L. Fasiczka, R. H. Howland and M. E. Thase (2003). "Effects of Sertraline on Sleep Architecture in Patients with Depression." J Clin Psychopharmacol 23: 540-548.	3
Joel, I., A. E. Begley, B. H. Mulsant, E. J. Lenze, S. Mazumdar, M. A. Dew, D. Blumberger, M. Butters, C. F. Reynolds, 3rd and I. G. I. Team (2013). "Dynamic Prediction of Treatment Response in Late-Life Depression." Am J Geriatr Psychiatry.	3
Jorge, R. E., R. G. Robinson, S. Arndt and S. Starkstein (2003). "Mortality and poststroke depression: a placebo-controlled trial of antidepressants." Am J Psychiatry 160: 1823-1829.	9

Kamijima, K., T. Higuchi, J. Ishigooka, T. Ohmori, N. Ozaki, S. Kanba, T. Kinoshita, T. Koyama and A. S. Group (2013). "Aripiprazole augmentation to antidepressant therapy in Japanese patients with major depressive disorder: a randomized, double-blind, placebo-controlled study (ADMIRE study)." <i>J Affect Disord</i> 151(3): 899-905.	3
Kanemoto, K., M. Matsubara, K. Yamashita, Y. Tarao, E. Inada and T. Sekine (2004). "Controlled comparison of two different doses of milnacipran in major depressive outpatients." <i>Int Clin Psychopharmacol</i> 19(6): 343-346.	3
Karp, J. F., D. Weiner, K. Seligman, M. Butters, M. Miller, E. Frank, J. Stack, B. H. Mulsant, B. Pollock, M. A. Dew, D. J. Kupfer and C. F. Reynolds lii (2005). "Body pain and treatment response in late-life depression." <i>American Journal of Geriatric Psychiatry</i> 13(3): 188-194.	3
Kasckow, J., I. Fellows, S. Golshan, E. Solorzano, T. Meeks and S. Zisook (2010). "Treatment of subsyndromal depressive symptoms in middle-age and older patients with schizophrenia: Effect of age on response." <i>American Journal of Geriatric Psychiatry</i> 18: 853-857.	9
Kasckow, J., N. Lanouette, T. Patterson, I. Fellows, S. Golshan, E. Solorzano and S. Zisook (2010). "Treatment of subsyndromal depressive symptoms in middle-aged and older adults with schizophrenia: effect on functioning." <i>Int J Geriatr Psychiatry</i> 25: 183-190.	7
Kasper, S., I. G. Anhelescu, A. Szegedi, A. Dienel and M. Kieser (2006). "Superior efficacy of St John's wort extract WS 5570 compared to placebo in patients with major depression: a randomized, double-blind, placebo-controlled, multi-center trial [ISRCTN77277298]." <i>BMC Med</i> 4: 14.	3
Kasper, S., O. M. Lemming and S. de (2006). "Escitalopram in the long-term treatment of major depressive disorder in elderly patients." <i>Neuropsychobiology</i> 54: 152-159.	3
Kasper, S., S. Baldwin David, L. Lönn Sara and J.-P. Boulenger (2009). "Superiority of escitalopram to paroxetine in the treatment of depression." <i>European Neuropsychopharmacology</i> 19: 229-237.	6
Katila, H., I. Mezhebovsky, A. Mulroy, L. Berggren, H. Eriksson, W. Earley and C. Datto (2013). "Randomized, Double-Blind Study of the Efficacy and Tolerability of Extended Release Quetiapine Fumarate (Quetiapine XR) Monotherapy in Elderly Patients With Major Depressive Disorder." <i>Am J Geriatr Psychiatry</i> .	4
Katon, W., M. Von Korff, E. Lin, E. Walker, G. E. Simon, T. Bush, P. Robinson and J. Russo (1995). "Collaborative management to achieve treatment guidelines. Impact on depression in primary care." <i>JAMA</i> 273(13): 1026-1031.	7
Katona, C. L. E., B. N. Hunter and J. Bray (1998). "A double-blind comparison of the efficacy and safety of paroxetine and imipramine in the treatment of depression with dementia." <i>Int J Geriatr Psychiatry</i> 13(2): 100-108.	3
Katona, C., Does being elderly alter response in MDD?, <i>International Journal of Psychiatry in Clinical Practice</i> , 17 (2013) 10. Accession:	9
Katona, C., T. Hansen and C. K. Olsen (2012). "A randomised, double-blind, placebocontrolled, active-referenced study of the multimodal antidepressant Lu AA21004 in the treatment of elderly depressed patients." <i>European Neuropsychopharmacology</i> 22: S258-S259.	3

Katz, A. J., S. B. Dusetzina, J. F. Farley, A. R. Ellis, B. N. Gaynes, W. C. Castillo, T. Sturmer and R. A. Hansen (2012). "Distressing adverse events after antidepressant switch in the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) trial: influence of adverse events during initial treatment with citalopram on development of subsequent adverse events with an alternative antidepressant." <i>Pharmacotherapy</i> 32(3): 234-243.	6
Katz, I. R., G. M. Simpson, S. M. Curlik, P. A. Parmelee and C. Muhly (1990). "Pharmacologic treatment of major depression for elderly patients in residential care settings." <i>Journal of Clinical Psychiatry</i> 51: 41-47.	9
Kavoussi, R. J., R. T. Segraves, A. R. Hughes, J. A. Ascher and J. A. Johnston (1997). "Double-blind comparison of bupropion sustained release and sertraline in depressed outpatients." <i>J Clin Psychiatry</i> 58(12): 532-537.	3
Keegan, D., R. C. Bowen, S. Blackshaw, S. Saleh, N. Dayal, F. Remillard, S. Shrikhande, S. Cebrian Perez and A. Boulton (1991). "A comparison of fluoxetine and amitriptyline in the treatment of major depression." <i>Int Clin Psychopharmacol</i> 6(2): 117-124.	3
Keers, R., C. Bonvicini, C. Scassellati, R. Uher, A. Placentino, C. Giovannini, M. Rietschel, N. Henigsberg, D. Kozel, O. Mors, W. Maier, J. Hauser, D. Souery, J. Mendlewicz, C. Schmal, A. Zobel, E. R. Larsen, A. Szczepankiewicz, Z. Kovacic, A. Elkin, I. Craig, P. McGuffin, A. E. Farmer, K. J. Aitchison and M. Gennarelli (2011). "Variation in GNB3 predicts response and adverse reactions to antidepressants." <i>Journal of Psychopharmacology</i> 25: 867-874.	3
Keers, R., R. Uher, B. Gupta, M. Rietschel, G. Schulze Thomas, J. Hauser, M. Skibinska, N. Henigsberg, P. Kalember, W. Maier, A. Zobel, O. Mors, S. Kristensen Ann, D. Kozel, C. Giovannini, J. Mendlewicz, S. Kumar, P. McGuffin, E. Farmer Anne and J. Aitchison Katherine (2010). "Stressful life events, cognitive symptoms of depression and response to antidepressants in GENDEP." <i>J Affect Disord</i> 127: 337-342.	3
Keitner, G. I., S. J. Garlow, C. E. Ryan, P. T. Ninan, D. A. Solomon, C. B. Nemeroff and M. B. Keller (2009). "A randomized, placebo-controlled trial of risperidone augmentation for patients with difficult-to-treat unipolar, non-psychotic major depression." <i>J Psychiatr Res</i> 43(3): 205-214.	3
Keller Martin, B., J. L. Ruwe Frank, J. J. G. Janssens, J. M. A. Sitsen, R. Jokinen and J. Janczewski (2005). "Relapse Prevention With Gepirone ER in Outpatients With Major Depression." <i>J Clin Psychopharmacol</i> 25: 79-84.	3
Keller, M. B., A. J. Gelenberg, R. M. Hirschfeld, A. J. Rush, M. E. Thase, J. H. Kocsis, J. C. Markowitz, J. A. Fawcett, L. M. Koran, D. N. Klein, J. M. Russell, S. G. Kornstein, J. P. McCullough, S. M. Davis and W. M. Harrison (1998). "The treatment of chronic depression, part 2: a double-blind, randomized trial of sertraline and imipramine." <i>J Clin Psychiatry</i> 59: 598-607.	3
Kellner, C. H., R. G. Knapp, G. Petrides, T. A. Rummans, M. M. Husain, K. Rasmussen, M. Mueller, H. J. Bernstein, K. O'Connor, G. Smith, M. Biggs, S. H. Bailine, C. Malur, E. Yim, S. McClintock, S. Sampson and M. Fink (2006). "Continuation electroconvulsive therapy vs pharmacotherapy for relapse prevention in major depression: a multisite study from the Consortium for Research in Electroconvulsive Therapy (CORE)." <i>Arch Gen Psychiatry</i> 63(12): 1337-1344.	10

Kennedy, G. J. (2006). "The sequenced treatment alternatives to relieve depression studies: How applicable are the results for older adults?" Primary Psychiatry: 33-36.	3
Kennedy, S. H. and R. Emsley (2006). "Placebo-controlled trial of agomelatine in the treatment of major depressive disorder." Eur Neuropsychopharmacol 16: 93-100.	3
Kennedy, S. H., A. Avedisova, N. Gimenez-Montesinos, C. Belaidi, C. de Bodinat and G. Agomelatine Study (2014). "A placebo-controlled study of three agomelatine dose regimens (10mg, 25mg, 25-50mg) in patients with major depressive disorder." Eur Neuropsychopharmacol 24(4): 553-563.	3
Kerber, K. B., S. R. Wisniewski, J. F. Luther, A. F. Leuchter, I. D'Empaire, M. H. Trivedi and A. J. Rush (2012). "Effects of heart disease on depression treatment: results from the COMED study." Gen Hosp Psychiatry 34: 24-34.	7
Khan, A. U. (1981). "A comparison of the therapeutic and cardiovascular effects of a single nightly dose of Prothiaden (dothiepin, dosulepin) and Lentizol (sustained-release amitriptyline) in depressed elderly patients." Journal of International Medical Research 9(2): 108-112.	4
Khan, A., A. Bose, G. S. Alexopoulos, C. Gommoll, D. Li and C. Gandhi (2007). "Double-blind comparison of escitalopram and duloxetine in the acute treatment of major depressive disorder." Clin Drug Investig 27(7): 481-492.	9
Khan, A., A. J. Cutler, D. K. Kajdasz, S. Gallipoli, M. Athanasiou, D. S. Robinson, H. Whalen and C. R. Reed (2011). "A randomized, double-blind, placebo-controlled, 8-week study of vilazodone, a serotonergic agent for the treatment of major depressive disorder." J Clin Psychiatry 72: 441-447.	3
Khan, A., A. Sambunaris, J. Edwards, A. Ruth and D. S. Robinson (2014). "Vilazodone in the treatment of major depressive disorder: efficacy across symptoms and severity of depression." Int Clin Psychopharmacol 29(2): 86-92.	3
Khazaie, H., M. Rahimi, F. Tatari, M. Rezaei, F. Najafi and M. Tahmasian (2011). "Treatment of depression in type 2 diabetes with Fluoxetine or Citalopram?" Neurosciences (Riyadh) 16: 42-45.	4
Kiev, A. and A. Feiger (1997). "A double-blind comparison of fluvoxamine and paroxetine in the treatment of depressed outpatients." J Clin Psychiatry 58(4): 146-152.	9
Kilminster, S. and I. Hindmarch (1992) "Cognition and mood in elderly depressed patients treated with either fluoxetine or amitriptyline." Br J Clin Pharmacol, 245p-246p.	3
Kimura, M., A. Tateno and R. G. Robinson (2003). "Treatment of poststroke generalized anxiety disorder comorbid with poststroke depression: merged analysis of nortriptyline trials." Am J Geriatr Psychiatry 11: 320-327.	3
Kimura, M., R. G. Robinson and J. T. Kosier (2000). "Treatment of cognitive impairment after poststroke depression: A double- blind treatment trial." Stroke 31: 1482-1486.	3
Klysner, R., E. Pleidrup, H. L. Hansen, J. Bent-Hansen, P. D. Loldrup, M. Lunde and H. H. E. Petersen (2000). "The effectiveness of citalopram in the prevention of depression recurrence in elderly patients." International Journal of Neuropsychopharmacology: 211.	7
Knight, M. M. and E. A. Houseman (2008) "A collaborative model for the treatment of depression in homebound elders." Issues Ment Health Nurs, 974-991 DOI: 10.1080/01612840802279049.	7

Koesters, M., G. Guiana, A. Cipriani, T. Becker and C. Barbui (2013). "Agomelatine efficacy and acceptability revisited: systematic review and meta-analysis of published and unpublished randomised trials." <i>Br J Psychiatry</i> 203(3): 179-187.	9
Kohler, S., S. Hoffmann, T. Unger, B. Steinacher and T. Fydrich (2012). "Adherence to guidelines and effectiveness of inpatient treatment for unipolar depression." <i>Int J Psychiatry Clin Pract</i> 16(2): 103-112.	4
Kok, R. M. (2013). "What is the role of medications in late life depression?" <i>Psychiatr Clin North Am</i> 36(4): 597-605.	9
Kok, R. M., C. van Baarsen, W. A. Nolen and T. J. Heeren (2009). "Early response as predictor of final remission in elderly depressed patients." <i>Int J Geriatr Psychiatry</i> 24(11): 1299-1303.	3
Kok, R. M., D. Vink, P. Westers, W. A. Nolen and T. J. Heeren (2008). "Early behavioural changes during antidepressant treatment in elderly patients." <i>Int J Geriatr Psychiatry</i> 23(7): 769-771.	4
Kok, R. M., D. Vink, T. J. Heeren and W. A. Nolen (2007). "Lithium augmentation compared with phenelzine in treatment-resistant depression in the elderly: an open, randomized, controlled trial." <i>J Clin Psychiatry</i> 68: 1177-1185.	3
Kok, R. M., W. A. Nolen and T. J. Heeren (2007). "Venlafaxine versus nortriptyline in the treatment of elderly depressed inpatients: a randomised, double-blind, controlled trial." <i>Int J Geriatr Psychiatry</i> 22: 1247-1254.	3
Kok, R. M., W. A. Nolen and T. J. Heeren (2009). "Outcome of late-life depression after 3 years of sequential treatment." <i>Acta Psychiatr Scand</i> 119: 274-281.	3
Kok, R., T. Heeren and W. Nolen (2010). "Treatment of psychotic depression in the elderly compared with nonpsychotic depression." <i>J Clin Psychopharmacol</i> 30: 465-467.	10
Koran, L. M., A. J. Gelenberg, S. G. Kornstein, R. H. Howland, R. A. Friedman, C. DeBattista, D. Klein, J. H. Kocsis, A. F. Schatzberg, M. E. Thase, A. J. Rush, R. M. Hirschfeld, L. M. LaVange and M. B. Keller (2001). "Sertraline versus imipramine to prevent relapse in chronic depression." <i>J Affect Disord</i> 65: 27-36.	4
Koran, L. M., S. H. Hamilton, M. Hertzman, B. S. Meyers, A. E. Halaris, G. D. Tollefson, J. M. Downs, D. G. Folks, D. V. Jeste, L. W. Lazarus and et al. (1995). "Predicting response to fluoxetine in geriatric patients with major depression." <i>J Clin Psychopharmacol</i> 15: 421-427.	4
Koran, L., M. Hertzman, B. Meyers, S. Holman and G. Tollefson (1994). "Predicting response to fluoxetine in geriatric patients with DSM-III-R major depression." <i>Psychopharmacol Bull</i> : 76.	4
Kornstein, S. G., A. Bose, D. Li, K. G. Saikali and C. Gandhi (2006). "Escitalopram maintenance treatment for prevention of recurrent depression: a randomized, placebo-controlled trial." <i>J Clin Psychiatry</i> 67: 1767-1775.	9
Kornstein, S. G., A. F. Schatzberg, M. E. Thase, K. A. Yonkers, J. P. McCullough, G. I. Keitner, A. J. Gelenberg, S. M. Davis, W. M. Harrison and M. B. Keller (2000). "Gender differences in treatment response to sertraline versus imipramine in chronic depression." <i>Am J Psychiatry</i> 157: 1445-1452.	3

Kornstein, S. G., Q. Jiang, S. Reddy, J. J. Musgnung and C. J. Guico-Pabia (2010). "Short-term efficacy and safety of desvenlafaxine in a randomized, placebo-controlled study of perimenopausal and postmenopausal women with major depressive disorder." <i>Journal of Clinical Psychiatry</i> 71: 1088-1096.	6
Kornstein, S., A. Clayton, W. Bao and C. J. Guico-Pabia (2012). "Treatment of major depressive disorder with desvenlafaxine 50 mg/d in perimenopausal and postmenopausal women." <i>Menopause</i> 19: 1390-1391.	3
Koshino, Y., W. M. Bahk, H. Sakai and T. Kobayashi (2013). "The efficacy and safety of bupropion sustained-release formulation for the treatment of major depressive disorder: a multi-center, randomized, double-blind, placebo-controlled study in Asian patients." <i>Neuropsychiatr Dis Treat</i> 9: 1273-1280.	3
Kragh-Sorensen, P., B. Muller, J. V. Andersen, D. Buch and K. B. Stage (1995). "Moclobemide versus clomipramine in depressed patients in general practice. A randomized, double-blind, parallel, multicenter study." <i>J Clin Psychopharmacol</i> 15: 24S-30S.	3
Kranzler Henry, R., T. Mueller, J. Cornelius, M. Pettinati Helen, D. Moak, R. Martin Peter, R. Anthenelli, J. Brower Kirk, S. O'Malley, J. Mason Barbara, D. Hasin and M. Keller (2006). "Sertraline treatment of co-occurring alcohol dependence and major depression." <i>J Clin Psychopharmacol</i> 26: 13-20.	3
Kretschmar, J. H. (1979). "Results of a double-blind trial comparing mianserin and amitriptyline in elderly patients with depressive illness." <i>Excerpta-Medica-Ics</i> : 19-25.	4
Kretschmar, J. H. (1980). "Mianserin and amitriptyline in elderly hospitalized patients with depressive illness: A double-blind trial." <i>Curr Med Res Opin</i> 6: 144-151.	3
Krishnan, K. R., P. M. Doraiswamy and C. M. Clary (2001). "Clinical and treatment response characteristics of late-life depression associated with vascular disease: a pooled analysis of two multicenter trials with sertraline." <i>Prog Neuropsychopharmacol Biol Psychiatry</i> 25(2): 347-361.	10
Kuchibhatla, M. N. and G. G. Fillenbaum (2011). "Trajectory classes of depression in a randomized depression trial of heart failure patients: a reanalysis of the SADHART-CHF trial." <i>Am J Geriatr Pharmacother</i> 9: 483-494.	7
Kumar, V. (1999). "Post-stroke depression and treatment strategies including Aniracetam." <i>International Journal of Geriatric Psychopharmacology</i> 2: 40-46.	4
Laakman, G., M. Faltermaier-Temizel, S. Bossert-Zaudig, T. Baghai and G. Lorkowski (1995). "Treatment of depressive outpatients with lorazepam, alprazolam, amitriptyline and placebo." <i>Psychopharmacology (Berl)</i> 120: 109-115.	3
Laakmann, G., M. Faltermaier-Temizel, S. Bossert-Zaudig, T. Baghai and G. Lorkowski (1996). "Treatment of mild to moderate depressive syndrome. A placebo-controlled double-blind trial on therapy with benzodiazepines or antidepressants." <i>Munchener Medizinische Wochenschrift</i> 138: 39-44.	11
Lacasse, Y., L. Beaudoin, L. Rousseau and F. Maltais (2004). "Randomized trial of paroxetine in end-stage COPD." <i>Monaldi Arch Chest Dis</i> 61: 140-147.	3
Lader, M., H. F. Andersen and T. Baekdal (2005). "The effect of escitalopram on sleep problems in depressed patients." <i>Hum Psychopharmacol</i> 20(5): 349-354.	9

Laghrissi-Thode, F., B. G. Pollock, M. C. Miller, B. H. Mulsant, L. Altieri and M. S. Finkel (1995). "Double-blind comparison of paroxetine and nortriptyline on the postural stability of late-life depressed patients." <i>Psychopharmacol Bull</i> 31(4): 659-663.	9
Lai, H. and R. Tolat (2005). "Paroxetine-related tremor: While this SSRI can significantly improve mood, its adverse effect warrants second look." <i>Geriatrics</i> 60: 18-20.	4
Lakey, S. L., A. Z. LaCroix, S. L. Gray, S. Borson, C. D. Williams, D. Calhoun, J. S. Goveas, J. W. Smoller, J. K. Ockene, K. H. Masaki, M. Coday, M. C. Rosal and N. F. Woods (2012). "Antidepressant Use, Depressive Symptoms, and Incident Frailty in Women Aged 65 and Older from the Women's Health Initiative Observational Study." <i>J Am Geriatr Soc</i> 60(5): 854-861.	9
Lakshmanan, M., L. C. Mion and J. D. Frengley (1986). "Effective low dose tricyclic antidepressant treatment for depressed geriatric rehabilitation patients. A double-blind study." <i>J Am Geriatr Soc</i> 34: 421-426.	9
Lam, R. W., H. F. Andersen and A. G. Wade (2008). "Escitalopram and duloxetine in the treatment of major depressive disorder: A pooled analysis of two trials." <i>Int Clin Psychopharmacol</i> 23: 181-187.	4
Langworth, S., O. Bodlund and H. Agren (2006). "Efficacy and tolerability of reboxetine compared with citalopram: a double-blind study in patients with major depressive disorder." <i>J Clin Psychopharmacol</i> 26: 121-127.	9
Lattanzio, F., A. Cherubini, G. Furneri, B. Di and N. Marchionni (2008). "Sertraline treatment for depression associated with acute coronary syndromes: a cost analysis from the viewpoint of the Italian Healthcare System." <i>Aging Clin Exp Res</i> 20: 76-80.	3
Lauritzen, L., K. Odgaard, L. Clemmesen, M. Lunde, J. Ohrstrom, C. Black and P. Bech (1996). "Relapse prevention by means of paroxetine in ECT-treated patients with major depression: a comparison with imipramine and placebo in medium-term continuation therapy." <i>Acta Psychiatr Scand</i> 94: 241-251.	3
Lauritzen, L., L. Clemmesen, R. Klysner, D. Loldrup, M. Lunde, E. Schaumburg, S. Waarst and P. Bech (1992). "Combined treatment with imipramine and mianserin. A controlled pilot study." <i>Pharmacopsychiatry</i> 25(4): 182-186.	3
Laux, G., M. Friede and W. E. Muller (2013). "Treatment of comorbid anxiety and depression with escitalopram: Results of a post-marketing surveillance study." <i>Pharmacopsychiatry</i> 46: 16-22.	3
Lavretsky, H., Combination of methylphenidate with citalopram is superior to either drug alone in improving clinical and cognitive outcomes in geriatric depression, <i>Neuropsychopharmacology</i> , 38 (2013) S90. Accession:	4
Lavretsky, H., P. Siddarth, A. Kumar, C. F. Reynolds and rd (2008). "The effects of the dopamine and serotonin transporter polymorphisms on clinical features and treatment response in geriatric depression: a pilot study." <i>Int J Geriatr Psychiatry</i> 23: 55-59.	4
Lavretsky, H., S. Park, P. Siddarth, A. Kumar, C. F. Reynolds and rd (2006). "Methylphenidate-enhanced antidepressant response to citalopram in the elderly: a double-blind, placebo-controlled pilot trial." <i>Am J Geriatr Psychiatry</i> 14: 181-185.	3

Lecrubier, Y., G. Clerc, R. Didi and M. Kieser (2002). "Efficacy of St. John's wort extract WS 5570 in major depression: a double-blind, placebo-controlled trial." <i>Am J Psychiatry</i> 159: 1361-1366.	9
Lee, M. S., Y. M. Ahn, S. Chung, R. Walton, J. Raskin and M. S. Kim (2012). "The effect of initial duloxetine dosing strategy on nausea in Korean patients with major depressive disorder." <i>Psychiatry Investigation</i> 9: 391-399.	7
Lee, M. S., Y. M. Ahn, S. Chung, R. Walton, J. Raskin and M. S. Kim (2013). "The effect of initial duloxetine dosing strategy on nausea in Korean patients with major depressive disorder." <i>Psychiatry Investigation</i> 10: 391-399.	4
Leentjens, A. F., F. W. Vreeling, G. J. Luijckx and F. R. Verhey (2003). "SSRIs in the treatment of depression in Parkinson's disease." <i>Int J Geriatr Psychiatry</i> 18: 552-554.	3
Lekakis, J., I. Ikonomidis, Z. Papoutsi, P. Moutsatsou, M. Nikolaou, J. Parassis and D. T. Kremastinos (2010). "Selective serotonin re-uptake inhibitors decrease the cytokine-induced endothelial adhesion molecule expression, the endothelial adhesiveness to monocytes and the circulating levels of vascular adhesion molecules." <i>Int J Cardiol</i> 139: 150-158.	3
Lemoine, P., C. Guilleminault and E. Alvarez (2007). "Improvement in subjective sleep in major depressive disorder with a novel antidepressant, agomelatine: Randomized, double-blind comparison with venlafaxine." <i>Journal of Clinical Psychiatry</i> 68: 1723-1732.	9
Lenderking, W. R., H. Tennen, J. F. Nackley, M. S. Hale, R. R. Turner and M. A. Testa (1999). "The effects of venlafaxine on social activity level in depressed outpatients." <i>J Clin Psychiatry</i> 60: 157-163.	7
Lenoir, S., F. H. Degenring and R. Saller (1999). "A double-blind randomised trial to investigate three different concentrations of a standardised fresh plant extract obtained from the shoot tips of Hypericum perforatum L." <i>Phytomedicine</i> 6(3): 141-146.	4
Lenox-Smith Alan, J. and Q. Jiang (2008). "Venlafaxine extended release versus citalopram in patients with depression unresponsive to a selective serotonin reuptake inhibitor." <i>Int Clin Psychopharmacol</i> 23: 113-119.	3
Lenox-Smith, A. J. and A. Reynolds (2003). "A double-blind, randomised, placebo controlled study of venlafaxine XL in patients with generalised anxiety disorder in primary care." <i>Br J Gen Pract</i> 53(495): 772-777.	3
Lenze, E. J., B. H. Mulsant, M. A. Dew, M. K. Shear, P. Houck, B. G. Pollock, C. F. Reynolds and rd (2003). "Good treatment outcomes in late-life depression with comorbid anxiety." <i>J Affect Disord</i> 77: 247-254.	3
Lenze, E. J., M. Sheffrin, H. C. Driscoll, B. H. Mulsant, B. G. Pollock, M. A. Dew, F. Lotrich, B. Devlin, R. Bies, I. Reynolds and F. C (2008). "Incomplete response in late-life depression: Getting to remission." <i>Dialogues in Clinical Neuroscience</i> 10: 419-430.	10
Leuchter, A. F., I. A. Cook, W. S. Gilmer, L. B. Marangell, K. S. Burgoyne, R. H. Howland, M. H. Trivedi, S. Zisook, R. Jain, M. Fava, D. Iosifescu and S. Greenwald (2009). "Effectiveness of a quantitative electroencephalographic biomarker for predicting differential response or remission with escitalopram and bupropion in major depressive disorder." <i>Psychiatry Res</i> 169: 132-138.	3

Lewis, G., J. Mulligan, N. Wiles, P. Cowen, N. Craddock, M. Ikeda, D. Grozeva, V. Mason, D. Nutt, D. Sharp, D. Tallon, L. Thomas, M. C. O'Donovan and T. J. Peters (2011). "Polymorphism of the 5-HT transporter and response to antidepressants: randomised controlled trial." <i>Br J Psychiatry</i> 198: 464-471.	9
Lewis-Fernandez, R., C. Blanco, C. H. Mallinckrodt, M. M. Wohlrreich, J. G. Watkin and J. M. Plewes (2006). "Duloxetine in the treatment of major depressive disorder: Comparisons of safety and efficacy in U.S. Hispanic and majority Caucasian patients." <i>Journal of Clinical Psychiatry</i> 67(9): 1379-1390.	9
Li, C. M., X. D. Jiang, G. Liao, J. M. Lei, S. Lan and F. W. Ni (2004). "Effect of antidepressant drugs in early period on the recovery of post-stroke depression." <i>Chinese Journal of Clinical Rehabilitation</i> 8(19): 3713-3715.	11
Li, L. T., S. H. Wang, H. Y. Ge, J. Chen, S. W. Yue and M. Yu (2008). "The beneficial effects of the herbal medicine Free and Easy Wanderer Plus (FEWP) and fluoxetine on post-stroke depression." <i>J Altern Complement Med</i> 14(7): 841-846.	3
Li, N., W. D. Ji, D. H. Zhang and Y. Li (2006). "Efficacy and safety of reboxetine versus fluoxetine for the elders with depression." <i>Chinese Journal of New Drugs</i> : 1682-1684.	3 och 11
Li, X. H., H. W. Zhang and Y. D. Pang (2004). "A comparative evaluation on the effect of citalopram and amitriptyline in treating senile depression." <i>Chinese Journal of Clinical Rehabilitation</i> 8(27): 5762-5763.	11
Li, X., S. Zheng and X. Lu (2007) "A comparative study of reboxetine and doxepine in the treatment of patients with depression in old age." <i>Journal of Psychiatry</i> , 35-36.	11
Li, Y. M., G. Y. Li, X. L. Wang and L. F. Yu (2005). "Rehabilitation effect of San John's Wort extract on depression and myocardial function after coronary artery bypass grafting: A randomized grouping, placebo-control and blind evaluation." <i>Chinese Journal of Clinical Rehabilitation</i> 9: 38-39.	4
Liang, K. (2006) "Comparative study of mirtazapine and sertraline in treatment of elderly depressive patient." <i>Journal of Heze Medical College</i> , 5-7.	11
Licht, R. W. and G. Danish University Antidepressant (2013). "Is it possible to evaluate true prophylactic efficacy of antidepressants in severely ill patients with recurrent depression? Lessons from a placebo-controlled trial. The fifth trial of the Danish University Antidepressant Group (DUAG-5)." <i>J Affect Disord</i> 148(2-3): 286-290.	3
Lieberman Daniel, Z., A. Montgomery Stuart, A. Tourian Karen, C. Brisard, G. Rosas, K. Padmanabhan, J.-M. Germain and B. Pitrosky (2008). "A pooled analysis of two placebo-controlled trials of desvenlafaxine in major depressive disorder." <i>Int Clin Psychopharmacol</i> 23: 188-197.	4
Liebowitz, M. R., P. P. Yeung and R. Entsuah (2007). "A randomized, double-blind, placebo-controlled trial of desvenlafaxine succinate in adult outpatients with major depressive disorder." <i>Journal of Clinical Psychiatry</i> 68: 1663-1672.	3
Liebowitz, M., H. A. Croft, D. K. Kajdasz, H. Whalen, S. Gallipoli, M. Athanasiou and C. R. Reed (2011). "The safety and tolerability profile of vilazodone, a novel antidepressant for the treatment of major depressive disorder." <i>Psychopharmacol Bull</i> 44: 2.	3

Liebowitz, M., R. W. Lam, U. Lepola, C. Datto, D. Sweitzer and H. Eriksson (2010). "Efficacy and tolerability of extended release quetiapine fumarate monotherapy as maintenance treatment of major depressive disorder: a randomized, placebo-controlled trial." <i>Depress Anxiety</i> 27: 964-976.	10
Limosin, F., J. C. Samuelian and F. Rouillon (2007). "Multicenter double-blind study of the efficacy of paroxetine versus clomipramine in elderly patients with major depression." <i>Journal of Aging and Pharmacotherapy</i> 13: 7-19.	3
Lin, E. H., L. Tang, W. Katon, M. T. Hegel, M. D. Sullivan and J. Unutzer (2006). "Arthritis pain and disability: response to collaborative depression care." <i>Gen Hosp Psychiatry</i> 28(6): 482-486.	3
Link, C. and G. Dunbar (1992). "An overview of studies comparing the efficacy, safety, and tolerability of paroxetine and clomipramine." <i>Nordic Journal of Psychiatry, Supplement</i> 46: 17-22.	4
Llorca, P. M., J. M. Azorin, N. Despiegel and P. Verpillat (2005). "Efficacy of escitalopram in patients with severe depression: A pooled analysis." <i>Int J Clin Pract</i> 59: 268-275.	6
Locklear, J. C., H. Svedsater, C. Datto and J. Endicott (2013). "Effects of once-daily extended release quetiapine fumarate (quetiapine XR) on quality of life and sleep in elderly patients with major depressive disorder." <i>J Affect Disord</i> .	6
Lustman, P. J., K. E. Freedland, L. S. Griffith and R. E. Clouse (2000). "Fluoxetine for depression in diabetes: a randomized double-blind placebo-controlled trial." <i>Diabetes Care</i> 23: 618-623.	3
Lustman, P. J., L. S. Griffith, R. E. Clouse, K. E. Freedland, S. A. Eisen, E. H. Rubin, R. M. Carney and J. B. McGill (1997). "Effects of nortriptyline on depression and glycemic control in diabetes: results of a double-blind, placebo-controlled trial." <i>Psychosom Med</i> 59: 241-250.	3
Lyketsos, C. G., E. Weiller, C. Katona and P. Gorwood (2011). "Are old-old patients with major depression more likely to relapse than young-old patients during continuation treatment with escitalopram?" <i>BMC Geriatr</i> 11: 2.	3
Lyketsos, C. G., J. M. Sheppard, C. D. Steele, S. Kopunek, M. Steinberg, A. S. Baker, J. Brandt and P. V. Rabins (2000). "Randomized, placebo-controlled, double-blind clinical trial of sertraline in the treatment of depression complicating Alzheimer's disease: initial results from the Depression in Alzheimer's Disease study." <i>Am J Psychiatry</i> 157: 1686-1689.	3
Maas, D. W., G. J. Westendorp Rudi, M. Willems Jorien, C. de, J. M. Anton, M. van der and C. Rose (2010). "TNF- α antagonist infliximab in the treatment of depression in older adults: Results of a prematurely ended, randomized, placebo-controlled trial." <i>J Clin Psychopharmacol</i> 30: 343-345.	9
Mahableshwarkar, A., Jacobsen, P., Chen, Y., Serenko, M. and Palo, W., The Safety and tolerability profile of vortioxetine in the treatment of patients with major depressive disorder aged 65 years and older, <i>American Journal of Geriatric Psychiatry</i> , 22 (2014) S143-S144. Accession:	9
Mahapatra, S. N. and D. Hackett (1997). "A randomised, double-blind, parallel-group comparison of venlafaxine and dothiepin in geriatric patients with major depression." <i>Int J Clin Pract</i> 51: 209-213.	9

Mahmoud, R. A., G. J. Pandina, I. Turkoz, C. Kosik-Gonzalez, C. M. Canuso, M. J. Kujawa and G. M. Gharabawi-Garibaldi (2007). "Risperidone for treatment-refractory major depressive disorder: a randomized trial." <i>Ann Intern Med</i> 147: 593-602.	10
Maksinczyk (1997) "Double-blind, multicentre, phase III study to compare the efficacy, safety and tolerability of citalopram and amitriptyline in elderly depressed patients in general practice." National Research Register.	3
Malaguarnera, M., R. Bella, M. Vacante, M. Giordano, G. Malaguarnera, M. P. Gargante, M. Motta, A. Mistretta, L. Rampello and G. Pennisi (2011). "Acetyl-L-carnitine reduces depression and improves quality of life in patients with minimal hepatic encephalopathy." <i>Scand J Gastroenterol</i> 46(6): 750-759.	3
Mallinckrodt, C. H., J. G. Watkin, C. Liu, M. M. Wohlreich and J. Raskin (2005). "Duloxetine in the treatment of major depressive disorder: A comparison of efficacy in patients with and without melancholic features." <i>BMC Psychiatry</i> 5.	6
Malsch, U. (1984). "Treatment of elderly depressive patients: Not simply a question of mood change." <i>Therapiewoche</i> 34: 2883-2885.	11
Malsch, U., K. Jager, B. Schwalb and W. Fischer (1996). "Antidepressant treatment in geriatric patients. Efficacy and cardiac tolerability of trimipramine vs. mianserin." <i>Munchener Medizinische Wochenschrift</i> 138(34): 29-33.	11
Mamo, D. C., B. G. Pollock, B. Mulsant, P. R. Houck, S. Bensasi, M. C. Miller, M. S. Redfern and C. Reynolds lii (2002). "Effects of nortriptyline and paroxetine on postural sway in depressed elderly patients." <i>Am J Geriatr Psychiatry</i> 10: 199-205.	4
Mamo, D. C., R. A. Sweet, B. H. Mulsant, B. G. Pollock, M. D. Miller, J. A. Stack, A. E. Begley, C. F. Reynolds and rd (2000). "Effect of nortriptyline and paroxetine on extrapyramidal signs and symptoms: A prospective double-blind study in depressed elderly patients." <i>Am J Geriatr Psychiatry</i> 8: 226-231.	3
Mannel, M., U. Kuhn, U. Schmidt, M. Ploch and H. Murck (2010). "St. John's wort extract LI160 for the treatment of depression with atypical features—A double-blind, randomized, and placebo-controlled trial." <i>J Psychiatr Res</i> 44: 760-767.	10
Mao, P. X., Y. L. Tang, F. Jiang, L. Shu, X. Gu, M. Li, M. Qian, C. Ma, P. B. Mitchell and Z. J. Cai (2008). "Escitalopram in major depressive disorder: a multicenter, randomized, double-blind, fixed-dose, parallel trial in a Chinese population." <i>Depress Anxiety</i> 25: 46-54.	7 och 9
Marcus, R. N., R. D. McQuade, W. H. Carson, D. Hennicken, M. Fava, J. S. Simon, M. H. Trivedi, M. E. Thase and R. M. Berman (2008). "The efficacy and safety of aripiprazole as adjunctive therapy in major depressive disorder: A second multicenter, randomized, double-blind, placebo-controlled study." <i>J Clin Psychopharmacol</i> 28: 156-165.	3
Marraccini, R. L., C. F. Reynolds, 3rd, P. R. Houck, M. D. Miller, E. Frank, J. M. Perel, C. Cornes, S. Mazumdar and D. J. Kupfer (1999). "A double-blind, placebo-controlled assessment of nortriptyline's side-effects during 3-year maintenance treatment in elderly patients with recurrent major depression." <i>Int J Geriatr Psychiatry</i> 14(12): 1014-1018.	5

Martinez, J. M., W. Katon, J. H. Greist, K. Kroenke, M. E. Thase, A. L. Meyers, S. E. Edwards, L. B. Marangell, S. Shoemaker and R. Swindle (2012). "A pragmatic 12-week, randomized trial of duloxetine versus generic selective serotonin-reuptake inhibitors in the treatment of adult outpatients in a moderate-to-severe depressive episode." <i>Int Clin Psychopharmacol</i> 27: 17-26.	3
Maust, D. T., D. W. Oslin and M. E. Thase (2013). "Going beyond antidepressant monotherapy for incomplete response in nonpsychotic late-life depression: a critical review." <i>Am J Geriatr Psychiatry</i> 21(10): 973-986.	9
Max, M. B., R. Kishore-Kumar, S. C. Schafer, B. Meister, R. H. Gracely, B. Smoller and R. Dubner (1991). "Efficacy of desipramine in painful diabetic neuropathy: a placebo-controlled trial." <i>Pain</i> 45: 3-9; discussion 1.	3
Mayer, L. S., R. C. Bay, A. Politis, M. Steinberg, C. Steele, A. S. Baker, P. V. Rabins and C. G. Lyketsos (2006). "Comparison of three rating scales as outcome measures for treatment trials of depression in Alzheimer disease: findings from DIADS." <i>Int J Geriatr Psychiatry</i> 21(10): 930-936.	6
McCue, R. E., A. Georgotas, N. Nagachandran, B. Abdul, E. A. Go, R. F. Suckow and T. B. Cooper (1989). "Plasma levels of nortriptyline and 10-hydroxynortriptyline and treatment-related electrocardiographic changes in the elderly depressed." <i>J Psychiatr Res</i> 23: 73-79.	3
McFarlane, A., M. V. Kamath, E. L. Fallen, V. Malcolm, F. Cherian and G. Norman (2001). "Effect of sertraline on the recovery rate of cardiac autonomic function in depressed patients after acute myocardial infarction." <i>Am Heart J</i> 142: 617-623.	9
McGrath, P. J., A. Y. Khan, M. H. Trivedi, J. W. Stewart, D. W. Morris, S. R. Wisniewski, S. Miyahara, A. A. Nierenberg, M. Fava and A. J. Rush (2008). "Response to a selective serotonin reuptake inhibitor (citalopram) in major depressive disorder with melancholic features: a STAR*D report." <i>J Clin Psychiatry</i> 69: 1847-1855.	4
McGrath, P. J., J. W. Stewart, F. M. Quitkin, Y. Chen, J. E. Alpert, A. A. Nierenberg, M. Fava, J. Cheng and E. Petkova (2006). "Predictors of relapse in a prospective study of fluoxetine treatment of major depression." <i>American Journal of Psychiatry</i> 163: 1542-1548.	3
McPartlin, G. M., A. Reynolds, C. Andersen and J. Casoy (1998). "A comparison of once-daily venlafaxine XR and paroxetine in depressed outpatients treated in general practice." <i>Primary Care Psychiatry</i> 4(3): 127-132.	3
Medori, R., R. Marcus and M. Zibellini (1997). "A multicenter, double-blind comparison of nefazodone and maprotiline in the treatment of non-psychotic elderly depressed patients." <i>Pharmacopsychiatry</i> : 197.	3
Meignan-Debray, S., B. Forette and M. Roger (1990). "Double-blind trial of medfoxamine (Cledial(registered trademark)) versus placebo in elderly patients with depressive disorders." <i>Psychologie Medicale</i> 22(9): 883-892.	11
Menchetti, M., B. Bortolotti, P. Rucci, P. Scocco, A. Bombi and D. Berardi (2010). "Depression in Primary care: Interpersonal Counseling vs Selective serotonin reuptake inhibitors. The DEPICS Study. A multicenter randomized controlled trial. Rationale and design." <i>BMC Psychiatry</i> 10.	4
Mendels, J., R. Johnston, J. Mattes and R. Riesenbergs (1993). "Efficacy and safety of b.i.d. doses of venlafaxine in a dose-response study." <i>Psychopharmacol Bull</i> 29: 169-174.	3

Mendonça Lima, C. A., P. Baumann, M. Brawand-Amey, C. Brogli, S. Jacquet, N. Cochard, K. Powell-Golay and C. B. Eap (2005) "Effect of age and gender on citalopram and desmethylcitalopram steady-state plasma concentrations in adults and elderly depressed patients." <i>Prog Neuropsychopharmacol Biol Psychiatry</i> , 952-956 DOI: 10.1016/j.pnpbp.2005.06.001.	3
Menza, M., R. D. Dobkin, H. Marin, M. H. Mark, M. Gara, S. Buyske, K. Bienfait and A. Dicke (2009). "The impact of treatment of depression on quality of life, disability and relapse in patients with Parkinson's disease." <i>Mov Disord</i> 24: 1325-1332.	3
Merideth, C. H., J. P. Feighner and G. Hendrickson (1984). "A double-blind comparative evaluation of the efficacy and safety of nomifensine, imipramine, and placebo in depressed geriatric outpatients." <i>Journal of Clinical Psychiatry</i> 45(4 II): 73-77.	6
Meyers, B. S., A. J. Flint, A. J. Rothschild, B. H. Mulsant, E. M. Whyte, C. Peasley-Miklus, E. Papademetriou, A. C. Leon, M. Heo and S.-P. Group (2009). "A double-blind randomized controlled trial of olanzapine plus sertraline vs olanzapine plus placebo for psychotic depression: the study of pharmacotherapy of psychotic depression (STOP-PD)." <i>Arch Gen Psychiatry</i> 66(8): 838-847.	3
Meyers, B. S., S. A. Klimstra, M. Gabriele, M. Hamilton, T. Kakuma, F. Tirumalasetti and G. S. Alexopoulos (2001). "Continuation treatment of delusional depression in older adults." <i>Am J Geriatr Psychiatry</i> 9(4): 415-422.	3
Meyers, B. S., S. Mattis, M. Gabriele and T. Kakuma (1991). "Effects of nortriptyline on memory self-assessment and performance in recovered elderly depressives." <i>Psychopharmacol Bull</i> : 295-299.	3
Miao, S. Y. and Y. J. Shi (2004). "Related factors of post-stroke depression and therapeutical effect of citalopram." <i>Chinese Journal of Clinical Rehabilitation</i> 8: 3718-3719.	11
Michelson, D., M. Schmidt, J. Lee and R. Tepner (2001). "Changes in sexual function during acute and six-month fluoxetine therapy: a prospective assessment." <i>J Sex Marital Ther</i> 27(3): 289-302.	3
Middleton, R. S. (1976). "Treating elderly depressed patients with amitriptyline." <i>Modern Geriatrics</i> .	3
Middleton, R. S. W. (1975). "A comparison between maprotiline (Ludiomil) and imipramine in the treatment of depressive illness in the elderly." <i>Journal of International Medical Research</i> 3(sup.2): 79-83.	4
Mikami, K., R. E. Jorge, H. P. Adams, Jr., P. H. Davis, E. C. Leira, M. Jang and R. G. Robinson (2011). "Effect of antidepressants on the course of disability following stroke." <i>Am J Geriatr Psychiatry</i> 19(12): 1007-1015.	9
Miller, L. G. and K. Winther (2006). "A placebo-controlled double-blind study of the efficacy of Hypericum perforatum (St John's wort) in minor depressed vs. dysthymic patients." <i>Focus on Alternative & Complementary Therapies</i> 11: 197-198.	3
Miller, M. D., E. I. Curtiss, L. Marino, P. R. Houck, C. F. Paradis, S. Mazumdar, B. G. Pollock, J. Foglia, C. F. Reynolds and rd (1998). "Long-term ECG changes in depressed elderly patients treated with nortriptyline. A double-blind, randomized, placebo-controlled evaluation." <i>Am J Geriatr Psychiatry</i> 6: 59-66.	9

Miotto, P., L. Allegro and G. Ziliotto (1983). "A controlled study into the effects of Nomifensine in the treatment of depression in the elderly." <i>Archivio per le Scienze Mediche</i> 140(3): 245-258.	11
Mitchell, P. H., R. C. Veith, K. J. Becker, A. Buzaitis, K. C. Cain, M. Fruin, D. Tirschwell and L. Teri (2009). "Brief psychosocial-behavioral intervention with antidepressant reduces poststroke depression significantly more than usual care with antidepressant: living well with stroke: randomized, controlled trial." <i>Stroke</i> 40(9): 3073-3078.	4
Miyai, I. and M. J. Reding (1998). "Effects of antidepressants on functional recovery following stroke: A double-blind study." <i>Journal of Neurologic Rehabilitation</i> 12: 5-13.	7
Mocking, R. J. T., J. Assies, M. Bot, E. H. J. M. Jansen, A. H. Schene and F. Pouwer (2012). "Biological Effects of Add-On Eicosapentaenoic Acid Supplementation in Diabetes Mellitus and Co-Morbid Depression: A Randomized Controlled Trial." <i>PLoS One</i> 7(11).	3
Mohapatra, P. K., N. Kar, G. C. Kar and M. Behera (2005). "Effectiveness of sertraline in treatment of depression in a consecutive sample of patients with acute myocardial infarction: Six month prospective study on outcome." <i>Clinical Practice and Epidemiology in Mental Health</i> 1.	9
Moizeszowicz, J. and S. Subira (1977). "Controlled trial of nomifensin (HOE 984) and viloxazine in the treatment of depression in the elderly." <i>J Clin Pharmacol</i> 17(1): 81-83.	4
Mokhber, N. and M. M. Gharavi (2007) "Comparison of nortriptyline and bupropion in major depressive disorder among elderly patients." <i>Journal of Pakistan Psychiatric Society</i> , 83- 87.	3
Moller, H. J. and H. P. Volz (1993). "Brofaromine in elderly major depressed patients--a comparative trial versus imipramine." <i>Eur Neuropsychopharmacol</i> 3(4): 501-510.	3
Moller, H. J., K. Glaser, F. Leverkus and C. Gobel (2000). "Double-blind, multicenter comparative study of sertraline versus amitriptyline in outpatients with major depression." <i>Pharmacopsychiatry</i> 33(6): 206-212.	3
Monteleone, P. and M. Fabrazzo (1994). "Blood levels of mianserin and amitriptyline and clinical response in aged depressed patients." <i>Pharmacopsychiatry</i> 27(6): 238-241.	6
Montgomery, S. A. and G. Dunbar (1993). "Paroxetine is better than placebo in relapse prevention and the prophylaxis of recurrent depression." <i>Int Clin Psychopharmacol</i> 8: 189-195.	6
Montgomery, S. A. and H. F. Andersen (2006). "Escitalopram versus venlafaxine XR in the treatment of depression." <i>Int Clin Psychopharmacol</i> 21: 297-309.	3
Montgomery, S. A., A. K. T. Huusom and J. Bothmer (2004). "A randomised study comparing escitalopram with venlafaxine XR in primary care patients with major depressive disorder." <i>Neuropsychobiology</i> 50: 57-64.	4
Montgomery, S. A., J. G. Rasmussen, K. Lyby, P. Connor and P. Tanghoj (1992). "Dose response relationship of citalopram 20 mg, citalopram 40 mg and placebo in the treatment of moderate and severe depression." <i>Int Clin Psychopharmacol</i> 6 Suppl 5: 65-70.	6
Montgomery, S. A., L. Mansuy, A. Ruth, A. Bose, H. Li and D. Li (2013). "Efficacy and safety of levomilnacipran sustained release in moderate to severe major depressive disorder: A randomized, double-blind, placebo-controlled, proof-of-concept study." <i>Journal of Clinical Psychiatry</i> 74: 363-369.	3

Montgomery, S. A., P. E. Reimetz and M. Zivkov (1998). "Mirtazapine versus amitriptyline in the long-term treatment of depression: a double-blind placebo-controlled study." <i>Int Clin Psychopharmacol</i> 13: 63-73.	3
Montgomery, S., A. C. Altamura, H. Katila, C. Datto, J. Szamosi and H. Eriksson (2011). "Efficacy of once-daily extended release quetiapine fumarate (quetiapine XR) in elderly patients with major depressive disorder (MDD) according to baseline anxiety symptoms, sleep disturbance and pain levels." <i>Int J Psychiatry Clin Pract</i> 15: 32-33.	5
Montgomery, S., A. C. Altamura, H. Katila, C. Datto, J. Szamosi and H. Eriksson (2011). "Extended release quetiapine efficacy in elderly patients with major depressive disorder by anxiety, sleep disturbance and pain levels." <i>European Neuropsychopharmacology</i> 21: S362-S363.	3
Moon, K. T. (2012). "Is donepezil an effective treatment for depression in older persons?" <i>Am Fam Physician</i> 85: 5.	7
Morse, J. Q., P. A. Pilkonis, P. R. Houck, E. Frank, C. F. Reynolds and rd (2005). "Impact of cluster C personality disorders on outcomes of acute and maintenance treatment in late-life depression." <i>Am J Geriatr Psychiatry</i> 13: 808-814.	3
Moscovitch, A., C. A. Blashko, J. M. Eagles, G. Darcourt, C. Thompson, S. Kasper, R. M. Lane, O. International Collaborative Group on Sertraline in the Treatment of and Disorders (2004). "A placebo-controlled study of sertraline in the treatment of outpatients with seasonal affective disorder." <i>Psychopharmacology (Berl)</i> 171: 390-397.	3
Muijsers, R. B. R., G. L. Plosker and S. Noble (2002). "Sertraline: A review of its use in the management of major depressive disorder in elderly patients." <i>Drugs and Aging</i> 19: 377-392.	3
Mukherjee, P. K. and A. Davey (1986). "Differential dosing of trazodone in elderly depressed patients: A study to investigate optimal dosing." <i>Journal of International Medical Research</i> 14(5): 279-284.	3
Muller-Siecheneder, F., M. J. Muller, A. Hillert, A. Szegedi, H. Wetzel and O. Benkert (1998). "Risperidone versus haloperidol and amitriptyline in the treatment of patients with a combined psychotic and depressive syndrome." <i>J Clin Psychopharmacol</i> 18(2): 111-120.	3
Mulsant, B. H., B. G. Pollock, R. D. Nebes, M. D. Miller, J. T. Little, J. Stack, P. R. Houck, S. Bensasi, S. Mazumdar and C. F. Reynolds, 3rd (1999). "A double-blind randomized comparison of nortriptyline and paroxetine in the treatment of late-life depression: 6-week outcome." <i>J Clin Psychiatry</i> 60 Suppl 20: 16-20.	3
Mulsant, B. H., B. G. Pollock, R. Nebes, M. D. Miller, R. A. Sweet, J. Stack, P. R. Houck, S. Bensasi, S. Mazumdar, C. F. Reynolds and rd (2001). "A twelve-week, double-blind, randomized comparison of nortriptyline and paroxetine in older depressed inpatients and outpatients." <i>Am J Geriatr Psychiatry</i> 9: 406-414.	6
Mulsant, B. H., R. A. Sweet, J. Rosen, B. G. Pollock, G. S. Zubenko, T. Flynn, A. E. Begley, S. Mazumdar and C. F. Reynolds, 3rd (2001). "A double-blind randomized comparison of nortriptyline plus perphenazine versus nortriptyline plus placebo in the treatment of psychotic depression in late life." <i>J Clin Psychiatry</i> 62(8): 597-604.	3

Munro, C. A., C. F. Longmire, L. T. Drye, B. K. Martin, C. E. Frangakis, C. L. Meinert, J. E. Mintzer, A. P. Porsteinsson, P. V. Rabins, P. B. Rosenberg, L. S. Schneider, D. Weintraub, C. G. Lyketsos and G. Depression in Alzheimer's Disease Study-2 Research (2012). "Cognitive outcomes after sertraline treatment in patients with depression of Alzheimer disease." Am J Geriatr Psychiatry 20(12): 1036-1044.	3
Murphy, G. M., C. Kremer, H. Rodrigues, A. F. Schatzberg and G. Mitrazapine versus paroxetine Study (2003). "The apolipoprotein E epsilon4 allele and antidepressant efficacy in cognitively intact elderly depressed patients." Biol Psychiatry 54(7): 665-673.	4
Murphy, G. M., Jr, S. B. Hollander, H. E. Rodriguez, C. Kremer and A. F. Schatzberg (2004). "Effects of the serotonin transporter gene promoter polymorphism on mirtazapine and paroxetine efficacy and adverse events in geriatric major depression." Arch Gen Psychiatry 61: 1163-1169.	3
Murray, V., M. von Arbin, A. Bartfai, A. L. Berggren, A. M. Landtblom, J. Lundmark, P. Nasman, J. E. Olsson, M. Samuelsson, A. Terent, R. Varelius, M. Asberg and B. Martensson (2005). "Double-blind comparison of sertraline and placebo in stroke patients with minor depression and less severe major depression." J Clin Psychiatry 66(6): 708-716.	3
Nair, N. P., M. Amin, G. Schwartz, D. Dastoor, J. X. Thavundayil, J. Mirmiran, C. MacDonald and R. Phillips (1993). "A comparison of the cardiac safety and therapeutic efficacy of trimipramine versus doxepin in geriatric depressed patients." J Am Geriatr Soc 41(8): 863-867.	3
Nair, N. P., M. Amin, P. Holm, C. Katona, N. Klitgaard, K. Ng Ying, M. N, P. Kragh-Sorensen, H. Kuhn, C. A. Leek and K. B. Stage (1995). "Moclobemide and nortriptyline in elderly depressed patients. A randomized, multicentre trial against placebo." J Affect Disord 33: 1-9.	3
Narushima, K. and R. G. Robinson (2003). "The Effect of Early versus Late Antidepressant Treatment on Physical Impairment Associated with Poststroke Depression: Is There a Time-Related Therapeutic Window?" Journal of Nervous and Mental Disease 191(10): 645-652.	3
Navari, R. M., M. C. Brenner and M. N. Wilson (2008). "Treatment of depressive symptoms in patients with early stage breast cancer undergoing adjuvant therapy." Breast Cancer Res Treat 112: 197-201.	9
Navarro, V., C. Gasto, X. Torres, T. Marcos and L. Pintor (2001). "Citalopram versus nortriptyline in late-life depression: a 12-week randomized single-blind study." Acta Psychiatr Scand 103(6): 435-440.	7
Nebes, R. D., B. G. Pollock, B. H. Mulsant, M. A. Butters, M. D. Zmuda and C. F. Reynolds, 3rd (1999). "Cognitive effects of paroxetine in older depressed patients." J Clin Psychiatry 60 Suppl 20: 26-29.	4
Nebes, R. D., B. G. Pollock, P. R. Houck, M. A. Butters, B. H. Mulsant, M. D. Zmuda, C. F. Reynolds and rd (2003). "Persistence of cognitive impairment in geriatric patients following antidepressant treatment: a randomized, double-blind clinical trial with nortriptyline and paroxetine." J Psychiatr Res 37: 99-108.	6
Nelson, C., K. Delucchi and L. Schneider (2006). "Effects of Sertraline on Suicidal Thinking and Behavior in Late-Life Depression." 46th Annual NCDEU (New Clinical Drug Evaluation Unit) Meeting; 2006 June 12 - 15; Boca Raton, FL: 246.	3

Nelson, J. C., J. S. Kennedy, B. G. Pollock, F. Laghrissi-Thode, M. Narayan, M. S. Nobler, D. W. Robin, I. Gergel, J. McCafferty and S. Roose (1999). "Treatment of major depression with nortriptyline and paroxetine in patients with ischemic heart disease." <i>Am J Psychiatry</i> 156: 1024-1028.	7
Nelson, J. C., K. Delucchi and L. S. Schneider (2009). "Moderators of antidepressant response in late-life depression trials: A pooled meta-analysis." <i>American Journal of Geriatric Psychiatry</i> 17: A53-A54.	3
Nelson, J. C., K. Delucchi and L. Schneider (2007). "Suicidal thinking and behavior during treatment with sertraline in late-life depression." <i>Am J Geriatr Psychiatry</i> 15: 573-580.	3
Nelson, J. C., M. E. Thase, E. E. Bellocchio, L. M. Rollin, J. M. Eudicone, R. D. McQuade, R. N. Marcus, R. M. Berman and R. A. Baker (2012). "Efficacy of adjunctive aripiprazole in patients with major depressive disorder who showed minimal response to initial antidepressant therapy." <i>Int Clin Psychopharmacol</i> 27: 125-133.	4
Nelson, J. C., M. M. Wohlreich, C. H. Mallinckrodt, M. J. Detke, J. G. Watkin and J. S. Kennedy (2005). "Duloxetine for the treatment of major depressive disorder in older patients." <i>American Journal of Geriatric Psychiatry</i> 13(3): 227-235.	9
Nemeroff, C. B., M. E. Thase and E. S. Group (2007). "A double-blind, placebo-controlled comparison of venlafaxine and fluoxetine treatment in depressed outpatients." <i>J Psychiatr Res</i> 41(3-4): 351-359.	7
Neshkes, R. E., R. Gerner and L. F. Jarvik (1985). "Orthostatic effect of imipramine and doxepin in depressed geriatric outpatients." <i>J Clin Psychopharmacol</i> 5: 102-106.	3
Newhouse, P. A., K. R. Krishnan, P. M. Doraiswamy, E. M. Richter, E. D. Batzar and C. M. Clary (2000). "A double-blind comparison of sertraline and fluoxetine in depressed elderly outpatients." <i>J Clin Psychiatry</i> 61(8): 559-568.	3
Nickel, M. K., C. Lahmann, M. Muehlbacher, C. Nickel, F. Pedrosa Gil, W. Buschmann, N. Rother, R. Fartacek, P. K. Leiberich, K. Tritt, F. O. Mitterlehner, C. Kettler, C. Egger, W. K. Rother and T. H. Loew (2006). "Change in instrumental activities of daily living disability in female senior patients with musculoskeletal pain: a prospective, randomized, controlled trial." <i>Arch Gerontol Geriatr</i> 42(3): 247-255.	3
Nickel, M. K., C. Nickel, C. Lahmann, F. O. Mitterlehner, K. Tritt, P. K. Leiberich, W. K. Rother and T. H. Loew (2005). "Changes in instrumental activities of daily living disability after treatment of depressive symptoms in elderly women with chronic musculoskeletal pain: a double-blind, placebo-controlled trial." <i>Aging Clin Exp Res</i> 17(4): 293-296.	3
Nierenberg, A. A., F. M. Quitkin, C. Kremer, M. B. Keller and M. E. Thase (2004). "Placebo-controlled continuation treatment with mirtazapine: acute pattern of response predicts relapse." <i>Neuropsychopharmacology</i> 29: 1012-1018.	3
Nierenberg, A. A., J. H. Greist, C. H. Mallinckrodt, A. Prakash, A. Sambunaris, G. D. Tollefson and M. M. Wohlreich (2007). "Duloxetine versus escitalopram and placebo in the treatment of patients with major depressive disorder: Onset of antidepressant action, a non-inferiority study." <i>Curr Med Res Opin</i> 23(2): 401-416.	3
Ning, J. and D. Lu (2004). "Observation of the clinical efficacy and safety of mirtazapine on elder depression treatment." <i>Clinical Pharmaceuticals</i> : 63-64.	11

Nobler, M. S., D. P. Devanand, M. K. Kim, L. M. Fitzsimons, T. M. Singer, N. Turret, H. A. Sackeim and S. P. Roose (1996). "Fluoxetine treatment of dysthymia in the elderly." <i>Journal of Clinical Psychiatry</i> 57: 254-256.	3
Nobler, M. S., S. P. Roose, I. Prohovnik, J. R. Moeller, J. Louie, R. L. Heertum and H. A. Sackeim (2000). "Regional cerebral blood flow in mood disorders, V.: Effects of antidepressant medication in late-life depression." <i>Am J Geriatr Psychiatry</i> : 289-296.	4
Nozawa, K., A. Sekine, S. Hozumi and T. Shimizu (2011). "Effect of augmentation with olanzapine in outpatients with depression in partial remission with melancholic features: consecutive case series." <i>Psychiatry Clin Neurosci</i> 65(2): 199-202.	7
Nugent, D. (1979). "A double-blind study of viloxazine (Vivalan) and amitriptyline in depressed geriatric patients." <i>Clinical Trials Journal</i> 16(1): 13-17.	3
Oakes, T. M., C. Katona, P. Liu, M. Robinson, J. Raskin and J. H. Greist (2013). "Safety and tolerability of duloxetine in elderly patients with major depressive disorder: a pooled analysis of two placebo-controlled studies." <i>Int Clin Psychopharmacol</i> 28: 1-11.	3
Oakley, F., N. A. Khin, R. Parks, L. Bauer and T. Sunderland (2002). "Improvement in activities of daily living in elderly following treatment for post-bereavement depression." <i>Acta Psychiatrica Scandinavica</i> : 231-234.	3
O'Connor, C. M., W. Jiang, M. Kuchibhatla, S. G. Silva, M. S. Cuffe, D. D. Callwood, B. Zakhary, W. G. Stough, R. M. Arias, S. K. Rivelli, R. Krishnan and S.-C. Investigators (2010). "Safety and efficacy of sertraline for depression in patients with heart failure: results of the SADHART-CHF (Sertraline Against Depression and Heart Disease in Chronic Heart Failure) trial." <i>J Am Coll Cardiol</i> 56: 692-699.	4
Olie, J. P. and S. Kasper (2007). "Efficacy of agomelatine, a MT1/MT2 receptor agonist with 5-HT2C antagonistic properties, in major depressive disorder." <i>Int J Neuropsychopharmacol</i> 10: 661-673.	3
Olie, J. P., K. P. Gunn and E. Katz (1997). "A double-blind placebo-controlled multicentre study of sertraline in the acute and continuation treatment of major depression." <i>European Psychiatry</i> 12(1): 34-41.	4
Oslin, D. W. (2005). "Treatment of late-life depression complicated by alcohol dependence." <i>Am J Geriatr Psychiatry</i> 13: 491-500.	9
Oslin, D. W., H. Ten, R. T, J. E. Streim, C. J. Datto, D. Weintraub, S. DiFilippo and I. R. Katz (2003). "Probing the safety of medications in the frail elderly: evidence from a randomized clinical trial of sertraline and venlafaxine in depressed nursing home residents." <i>J Clin Psychiatry</i> 64: 875-882.	4
Oslin, D. W., J. E. Streim, I. R. Katz, B. D. Smith, S. D. DiFilippo, H. Ten, R. T and T. Cooper (2000). "Heuristic comparison of sertraline with nortriptyline for the treatment of depression in frail elderly patients." <i>Am J Geriatr Psychiatry</i> 8: 141-149.	3
Osuji, I. J., E. Vera-Bolanos, T. J. Carmody and E. S. Brown (2010). "Pregnenolone for cognition and mood in dual diagnosis patients." <i>Psychiatry Res</i> 178: 309-312.	4
Ottevanger, E. A. (1994). "The efficacy of fluvoxamine in patients with severe depression." <i>Prog Neuropsychopharmacol Biol Psychiatry</i> 18: 731-740.	3
Overo, K. (1997) "Citalopram versus amitriptyline in depressed elderly." 150th Annual Meeting of the American Psychiatric Association; 1997 May 17-22; San Diego, CA.	3

Pae, C. U. (2009). "A review of the safety and tolerability of aripiprazole." <i>Expert Opin Drug Saf</i> 8: 373-386.	3
Paile-Hyvarinen, M., K. Wahlbeck and J. G. Eriksson (2007). "Quality of life and metabolic status in mildly depressed patients with type 2 diabetes treated with paroxetine: a double-blind randomised placebo controlled 6-month trial." <i>BMC Fam Pract</i> 8: 34.	3
Pancheri, P., C. Delle, M. Donnini, S. Seripa, C. Gambino, E. Vicario and L. Trillo (1994). "Effects of moclobemide on depressive symptoms and cognitive performance in a geriatric population: a controlled comparative study versus imipramine." <i>Clin Neuropharmacol</i> 17 Suppl 1: S58-73.	3
Pangallo, B., M. A. Dellva, D. N. D'Souza, B. Essink, J. Russell and C. Goldberger (2011). "A randomized, double-blind study comparing LY2216684 and placebo in the treatment of major depressive disorder." <i>J Psychiatr Res</i> 45: 748-755.	9
Papakostas George, I., C. Shelton Richard, M. Zajecka John, B. Etemad, K. Rickels, A. Clain, L. Baer, D. Dalton Elizabeth, R. Sacco Garret, D. Schoenfeld, M. Pencina, A. Meisner, T. Bottiglieri, E. Nelson, D. Mischoulon, E. Alpert Jonathan, G. Barbee James, S. Zisook and M. Fava (2012). "L-methylfolate as adjunctive therapy for SSRI-resistant major depression: Results of two randomized, double-blind, parallel-sequential trials." <i>Am J Psychiatry</i> 169: 1267-1274.	4
Papakostas George, I., V. Iosifescu Dan, T. Petersen, K. Hamill Sarah, E. Alpert Jonathan, A. Nierenberg Andrew, F. Rosenbaum Jerrold and M. Fava (2004). "Serum Cholesterol in the Continuation Phase of Pharmacotherapy With Fluoxetine in Remitted Major Depressive Disorder." <i>J Clin Psychopharmacol</i> 24: 467-469.	3
Papakostas, G. I., P. McGrath, J. Stewart, D. Charles, Y. Chen, D. Mischoulon, C. Dording and M. Fava (2008). "Psychic and somatic anxiety symptoms as predictors of response to fluoxetine in major depressive disorder." <i>Psychiatry Res</i> 161: 116-120.	3
Parnetti, L., S. Sommacal, L. Morselli, M. A and U. Senin (1993). "Multicentre controlled randomised double-blind placebo study of minaprine in elderly patients suffering from prolonged depressive reaction." <i>Drug Investigation</i> 6: 181-188.	3
Patat, A., C. Dubruc, C. Deschamp, G. Durrieu, S. Trocherie, I. Cimarosti, H. Allain, P. Rosenzweig and J. M. Gandon (1997). "EEG, MAO-A inhibition, pharmacokinetics and safety of befloxatone in the elderly." <i>Hum Psychopharmacol</i> 12(6): 557-571.	6
Patkar, A. A., K. Peindl, R. Mago, P. Mannelli and P. S. Masand (2006). "An open-label, rater-blinded, augmentation study of aripiprazole in treatment-resistant depression." <i>Prim Care Companion J Clin Psychiatry</i> 8(2): 82-87.	3
Patris, M., J. M. Bouchard, T. Bougerol, J. F. Charbonnier, J. F. Chevalier, G. Clerc, C. Cyran, P. Van Amerongen, O. Lemming and H. E. Hopfner Petersen (1996). "Citalopram versus fluoxetine: a double-blind, controlled, multicentre, phase III trial in patients with unipolar major depression treated in general practice." <i>Int Clin Psychopharmacol</i> 11(2): 129-136.	3
Pellicer, Y. and P. Schaeffer (1993). "[Multicenter double-blind study comparing the efficacy and tolerance of paroxetine and clomipramine in reactive depression in the elderly patient]." <i>Encephale</i> 19(3): 257-261.	3 och 11

Pelton, G. H., O. L. Harper, M. H. Tabert, H. A. Sackeim, N. Scarmeas, S. P. Roose and D. P. Devanand (2008). "Randomized double-blind placebo-controlled donepezil augmentation in antidepressant-treated elderly patients with depression and cognitive impairment: a pilot study." <i>Int J Geriatr Psychiatry</i> 23: 670-676.	6
Peniston, J. H., X. Hu, S. L. Potts, M. S. Wieman and D. C. Turk (2012). "Tolerability of concomitant use of selective serotonin reuptake inhibitors or serotonin-norepinephrine reuptake inhibitors and oxymorphone extended release." <i>Postgrad Med</i> 124: 114-122.	6
Perlis, R. H., C. M. Beasley, Jr, J. D. Wines, Jr, R. N. Tamura, C. Cusin, D. Shear, J. Amsterdam, F. Quitkin, R. E. Strong, J. F. Rosenbaum and M. Fava (2007). "Treatment-associated suicidal ideation and adverse effects in an open, multicenter trial of fluoxetine for major depressive episodes." <i>Psychother Psychosom</i> 76: 40-46.	4
Perroud, N., G. Bondolfi, R. Uher, M. Gex-Fabry, J. M. Aubry, G. Bertschy, A. Malafosse and M. Kosel (2011). "Clinical and genetic correlates of suicidal ideation during antidepressant treatment in a depressed outpatient sample." <i>Pharmacogenomics</i> 12(3): 365-377.	4
Perroud, N., R. Uher, A. Marusic, M. Rietschel, O. Mors, N. Henigsberg, J. Hauser, W. Maier, D. Souery, A. Placentino, A. Szczepankiewicz, L. Jorgensen, J. Strohmaier, A. Zobel, C. Giovannini, A. Elkin, C. Gunasinghe, J. Gray, D. Campbell, B. Gupta, A. E. Farmer, P. McGuffin and K. J. Aitchison (2009). "Suicidal ideation during treatment of depression with escitalopram and nortriptyline in genome-based therapeutic drugs for depression (GENDEP): a clinical trial." <i>BMC Med</i> 7: 60.	4
Perroud, N., R. Uher, J. Hauser, M. Rietschel, N. Henigsberg, A. Placentino, D. Kozel, W. Maier, O. Mors, D. Souery, M. Dmitrzak-Weglacz, L. Jorgensen, Z. Kovacic, C. Giovannini, J. Mendlewicz, A. Zobel, J. Strohmaier, P. McGuffin, J. Aitchison Katherine and A. Farmer (2010). "History of suicide attempts among patients with depression in the GENDEP project." <i>J Affect Disord</i> 123: 131-137.	3
Petracca, G. M., E. Chemerinski and S. E. Starkstein (2001). "A double-blind, placebo-controlled study of fluoxetine in depressed patients with Alzheimer's disease." <i>Int Psychogeriatr</i> 13(2): 233-240.	3
Petursson, H. (1993) "A controlled study of moclobemide in elderly depression with cognitive decline." <i>Psychopharmacology (Berl)</i> , B5.	3
Pezzella, G., R. Moslinger-Gehmayr and A. Contu (2001). "Treatment of depression in patients with breast cancer: a comparison between paroxetine and amitriptyline." <i>Breast Cancer Res Treat</i> 70: 1-10.	3
Philipp, M., R. Kohnen and K. O. Hiller (1999). "Hypericum extract versus imipramine or placebo in patients with moderate depression: Randomised multicentre study of treatment for eight weeks." <i>British Medical Journal</i> 319: 1534-1538.	4
Pia, S., D. Giorgio, R. Ciriello, A. Sannino, L. Simone and C. Paoletti (1992) "Double blind controlled study to evaluate the effectiveness and tolerability of fluoxetine versus mianserin in the treatment of depressive disorders among the elderly and their effects on cognitive-behavioral parameters." <i>New Trends in Experimental & Clinical Psychiatry</i> , 139-146.	4

Pigott, T. A., A. Prakash, L. M. Arnold, S. T. Aaronson, C. H. Mallinckrodt and M. M. Wohlrreich (2007). "Duloxetine versus escitalopram and placebo: an 8-month, double-blind trial in patients with major depressive disorder." <i>Curr Med Res Opin</i> 23(6): 1303-1318.	7
piLa, P., D. Giorgio, R. Ciriello, A. Sannino, S. De, C. Paoletti and C. V. Colonna (1992). "Evaluation of the efficacy, tolerability, and therapeutic profile of fluoxetine versus mianserin in the treatment of depressive disorders in the elderly." <i>Current Therapeutic Research - Clinical and Experimental</i> 52: 847-858.	3
Pilotto, A., G. D'Onofrio, F. Panza, M. Copetti, L. Cascavilla, F. Paris, F. Pellegrini, D. Seripa and L. Ferrucci (2012). "Treatment of late-life major depressive disorder with selective serotonin reuptake inhibitors improves the multidimensional prognostic index." <i>J Clin Psychopharmacol</i> 32: 726-729.	7
Pitychoutis, P. M., N. Kokras, D. Sanoudou, C. Dalla and Z. Papadopoulou-Daifoti (2013). "Pharmacogenetic considerations for late life depression therapy." <i>Expert Opin Drug Metab Toxicol</i> 9(8): 989-999.	9
Pizzi, C., S. Mancini, L. Angeloni, F. Fontana, L. Manzoli and G. M. Costa (2009). "Effects of selective serotonin reuptake inhibitor therapy on endothelial function and inflammatory markers in patients with coronary heart disease." <i>Clin Pharmacol Ther</i> 86: 527-532.	7
Poelinger, W. and H. Haber (1989). "Fluoxetine 40 mg vs maprotiline 75 mg in the treatment of out-patients with depressive disorders." <i>Int Clin Psychopharmacol</i> 4 Suppl 1: 47-50.	3
Pollock, B. G., B. H. Mulsant, R. Nebes, M. A. Kirshner, A. E. Begley, S. Mazumdar, C. F. Reynolds and rd (1998). "Serum anticholinergicity in elderly depressed patients treated with paroxetine or nortriptyline." <i>Am J Psychiatry</i> 155: 1110-1112.	3
Pollock, B. G., R. E. Ferrell, B. H. Mulsant, S. Mazumdar, M. Miller, R. A. Sweet, S. Davis, M. A. Kirshner, P. R. Houck, J. A. Stack, C. F. Reynolds and D. J. Kupfer (2000). "Allelic variation in the serotonin transporter promoter affects onset of paroxetine treatment response in late-life depression." <i>Neuropsychopharmacology</i> 23(5): 587-590.	3
Pope, H. G., Jr, G. H. Cohane, G. Kanayama, A. J. Siegel and J. I. Hudson (2003). "Testosterone gel supplementation for men with refractory depression: a randomized, placebo-controlled trial." <i>Am J Psychiatry</i> 160: 105-111.	10
Porter, R. J., A. J. Phipps, P. Gallagher, A. Scott, P. S. Stevenson and J. T. O'Brien (2005). "Effects of acute tryptophan depletion on mood and cognitive functioning in older recovered depressed subjects." <i>Am J Geriatr Psychiatry</i> 13: 607-615.	3
Pritchett, Y. L., M. D. Marciak, P. K. Corey-Lisle, R. A. Berzon, D. Desaiah and M. J. Detke (2007). "Use of effect size to determine optimal dose of duloxetine in major depressive disorder." <i>J Psychiatr Res</i> 41: 311-318.	6
Puchler, K., A. Plenker and H. P. Volz (1997). "Safety, tolerability and efficacy of the reversible monoamine oxidase inhibitor RS-8359 in early clinical trials." <i>Int Clin Psychopharmacol</i> 12: S31-S36.	9
Quante, A., F. Regen, F. Schindler, K. Volkmer, E. Severus, C. Urbanek, A. Luborzewski, S. Roepke and I. Anhelescu (2013). "Quetiapine as combination treatment with citalopram in unipolar depression with prominent somatic symptoms: a randomised, double-blind, placebo-controlled pilot study." <i>Psychiatr Danub</i> 25(3): 214-220.	3

Quilty, L. C., F. De Fruyt, J. P. Rolland, S. H. Kennedy, P. F. Rouillon and R. M. Bagby (2008). "Dimensional personality traits and treatment outcome in patients with major depressive disorder." <i>J Affect Disord</i> 108(3): 241-250.	3
Quitkin, F. M., E. Petkova, P. J. McGrath, B. Taylor, C. Beasley, J. Stewart, J. Amsterdam, M. Fava, J. Rosenbaum, F. Reimherr, J. Fawcett, Y. Chen and D. Klein (2003). "When should a trial of fluoxetine for major depression be declared failed?" <i>Am J Psychiatry</i> 160: 734-740.	3
Quitkin, F. M., P. J. McGrath, J. G. Rabkin, J. W. Stewart, W. Harrison, D. C. Ross, E. Tricamo, J. Fleiss, J. Markowitz and D. F. Klein (1991). "Different types of placebo response in patients receiving antidepressants." <i>Am J Psychiatry</i> 148(2): 197-203.	4
Raby, W. N., E. A. Rubin, F. Garawi, W. Cheng, E. Mason, L. Sanfilippo, S. Lord, A. Bisaga, E. Aharonovich, F. Levin, D. McDowell and E. V. Nunes (2014). "A randomized, double-blind, placebo-controlled trial of venlafaxine for the treatment of depressed cocaine-dependent patients." <i>Am J Addict</i> 23(1): 68-75.	3
Raffaele, R., L. Rampello, I. Vecchio, C. Tornali and M. Malaguarnera (1996). "Trazodone therapy of the post-stroke depression." <i>Arch Gerontol Geriatr</i> 23: 217-220.	3
Rahman, M. K., M. J. Akhtar, N. C. Savla, R. R. Sharma, J. M. Kellett and J. J. Ashford (1991). "A double-blind, randomised comparison of fluvoxamine with dothiepin in the treatment of depression in elderly patients." <i>Br J Clin Pract</i> 45(4): 255-258.	3
Raji, M. A. and S. R. Brady (2001). "Mirtazapine for treatment of depression and comorbidities in Alzheimer disease." <i>Ann Pharmacother</i> 35: 1024-1027.	3
Raji, M. A., P. D. Barnum, J. Freeman and A. B. Markowitz (2007). "Mirtazapine for depression and comorbidities in older patients with cancer [2]." <i>Annals of Pharmacotherapy</i> 41: 1548-1549.	3
Rampello, L., A. Alvano, S. Chiechio, R. Raffaele, I. Vecchio and M. Malaguarnera (2005). "An evaluation of efficacy and safety of reboxetine in elderly patients affected by "retarded" post-stroke depression. A random, placebo-controlled study." <i>Arch Gerontol Geriatr</i> 40: 275-285.	3
Rapaport, M. H., A. A. Nierenberg, R. Howland, C. Dording, P. J. Schettler and D. Mischoulon (2011). "The treatment of minor depression with St. John's Wort or citalopram: failure to show benefit over placebo." <i>J Psychiatr Res</i> 45: 931-941.	3
Rapaport, M. H., A. Bose and H. Zheng (2004). "Escitalopram continuation treatment prevents relapse of depressive episodes." <i>J Clin Psychiatry</i> 65: 44-49.	5
Rapaport, M. H., G. M. Gharabawi, C. M. Canuso, R. A. Mahmoud, M. B. Keller, C. A. Bossie, I. Turkoz, R. A. Lasser, A. Loescher, P. Bouhours, F. Dunbar and C. B. Nemeroff (2006). "Effects of risperidone augmentation in patients with treatment-resistant depression: Results of open-label treatment followed by double-blind continuation." <i>Neuropsychopharmacology</i> 31(11): 2505-2513.	3
Rapaport, M. H., G. M. Gharabawi, C. M. Canuso, R. A. Mahmoud, M. B. Keller, C. A. Bossie, I. Turkoz, R. A. Lasser, A. Loescher, P. Bouhours, F. Dunbar and C. B. Nemeroff (2006). "Effects of risperidone augmentation in patients with treatment-resistant depression: Results of open-label treatment followed by double-blind continuation." <i>Neuropsychopharmacology</i> 31: 2505-2513.	9

Rapaport, M. H., L. S. Schneider, D. L. Dunner, J. T. Davies and C. D. Pitts (2003). "Efficacy of controlled-release paroxetine in the treatment of late-life depression." <i>J Clin Psychiatry</i> 64: 1065-1074.	4
Rapaport, M. H., R. B. Lydiard, C. D. Pitts, D. Schaefer, E. I. Bartolic, M. Iyengar, M. Carfagno and A. Lipschitz (2009). "Low doses of controlled-release paroxetine in the treatment of late-life depression: a randomized, placebo-controlled trial." <i>J Clin Psychiatry</i> 70: 46-57.	7
Raskin, J., C. G. Wiltse, J. J. Dinkel, D. J. Walker, D. Desaiah and C. Katona (2008). "Safety and tolerability of duloxetine at 60 mg once daily in elderly patients with major depressive disorder." <i>J Clin Psychopharmacol</i> 28: 32-38.	10
Raskin, J., J. Y. Xu and D. K. Kajdasz (2008). "Time to response for duloxetine 60 mg once daily versus placebo in elderly patients with major depressive disorder." <i>Int Psychogeriatr</i> 20: 309-327.	3
Raskind, M. A. (1993). "Geriatric psychopharmacology Management of late-life depression and the noncognitive behavioral disturbances of Alzheimer's disease." <i>Psychiatric Clinics of North America</i> : 815-827.	3
Rasmussen, K. G., R. G. Knapp, M. M. Biggs, G. E. Smith, T. A. Rummans, G. Petrides, M. M. Husain, M. K. O'Connor, M. Fink and C. H. Kellner (2007). "Data management and design issues in an unmasked randomized trial of electroconvulsive therapy for relapse prevention of severe depression: the consortium for research in electroconvulsive therapy trial." <i>J ECT</i> 23(4): 244-250.	4
Ratti, E., K. Bellew, P. Bettica, H. Bryson, S. Zamuner, G. Archer, L. Squassante, A. Bye, D. Trist, K. R. Krishnan and S. Fernandes (2011). "Results from 2 randomized, double-blind, placebo-controlled studies of the novel NK ₁ receptor antagonist casopitant in patients with major depressive disorder." <i>J Clin Psychopharmacol</i> 31: 727-733.	4
Ratti, E., P. Bettica, R. Alexander, G. Archer, D. Carpenter, G. Evoniuk, R. Gomeni, E. Lawson, M. Lopez, H. Millns, A. Rabiner Eugenii, D. Trist, M. Trower, S. Zamuner, R. Krishnan and M. Fava (2013). "Full central neurokinin-1 receptor blockade is required for efficacy in depression: Evidence from orvepitant clinical studies." <i>Journal of Psychopharmacology</i> 27: 424-434.	4
Ravindran, A. V., R. Judge, B. N. Hunter, J. Bray and N. H. Morton (1997). "A double-blind, multicenter study in primary care comparing paroxetine and clomipramine in patients with depression and associated anxiety. Paroxetine Study Group." <i>J Clin Psychiatry</i> 58(3): 112-118.	3
Reed, C. R., D. K. Kajdasz, H. Whalen, M. C. Athanasiou, S. Gallipoli and M. E. Thase (2012). "The efficacy profile of vilazodone, a novel antidepressant for the treatment of major depressive disorder." <i>Curr Med Res Opin</i> 28: 27-39.	3
Reimherr Frederick, W., D. Amsterdam Jay, M. Quitkin Frederic, F. Rosenbaum Jerrold, M. Fava, J. Zajecka, M. Beasley Charles, Jr, D. Michelson, P. Roback and K. Sundell (1998). "Optimal length of continuation therapy in depression: A prospective assessment during long-term fluoxetine treatment." <i>Am J Psychiatry</i> 155: 1247-1253.	3
Reimherr, F. W., L. A. Cunningham, S. R. Batey, J. A. Johnston and J. A. Ascher (1998). "A multicenter evaluation of the efficacy and safety of 150 and 300 mg/d sustained-release bupropion tablets versus placebo in depressed outpatients." <i>Clin Ther</i> 20: 505-516.	9

Reimherr, F. W., M. L. Martin, J. M. Eudicone, B. K. Marchant, Q. V. Tran, A. Pikalov, R. N. Marcus, R. M. Berman and B. X. Carlson (2010). "A pooled MADRS/IDS cross-correlation analysis: clinician and patient self-report assessment of improvement in core depressive symptoms with adjunctive aripiprazole." <i>J Clin Psychopharmacol</i> 30: 300-305.	9
Reis, M., A. Aberg-Wistedt, H. Agren, P. Hoglund, A. C. Akerblad and F. Bengtsson (2004). "Serum disposition of sertraline, N-desmethylsertraline and paroxetine: a pharmacokinetic evaluation of repeated drug concentration measurements during 6 months of treatment for major depression." <i>Hum Psychopharmacol</i> 19: 283-291.	3
Rektorova, I., I. Rektor, M. Bares, V. Dostal, E. Ehler, Z. Fanfrdlova, J. Fiedler, H. Klajblová, P. Kulist'ak, P. Ressner, J. Svatova, K. Urbanek and J. Veliskova (2003). "Pramipexole and pergolide in the treatment of depression in Parkinson's disease: a national multicentre prospective randomized study." <i>Eur J Neurol</i> 10: 399-406.	3
Reynolds, C. F., 3rd, E. Frank, J. M. Perel, S. D. Imber, C. Cornes, R. K. Morycz, S. Mazumdar, M. D. Miller, B. G. Pollock, A. H. Rifai and et al. (1992). "Combined pharmacotherapy and psychotherapy in the acute and continuation treatment of elderly patients with recurrent major depression: a preliminary report." <i>Am J Psychiatry</i> 149(12): 1687-1692.	9
Reynolds, C. F., 3rd, G. S. Smith, M. A. Dew, B. H. Mulsant, M. D. Miller, M. Schlernitzauer, J. A. Stack, P. R. Houck and B. G. Pollock (2005). "Accelerating symptom-reduction in late-life depression: a double-blind, randomized, placebo-controlled trial of sleep deprivation." <i>Am J Geriatr Psychiatry</i> 13(5): 353-358.	7
Reynolds, C. F., rd, E. Frank, J. M. Perel, S. Mazumdar, M. A. Dew, A. Begley, P. R. Houck, M. Hall, B. Mulsant, M. K. Shear, M. D. Miller, C. Cornes and D. J. Kupfer (1996). "High relapse rate after discontinuation of adjunctive medication for elderly patients with recurrent major depression." <i>Am J Psychiatry</i> 153: 1418-1422.	7
Reynolds, C. F., rd, J. M. Perel, E. Frank, C. Cornes, M. D. Miller, P. R. Houck, S. Mazumdar, J. A. Stack, B. G. Pollock, M. A. Dew and D. J. Kupfer (1999). "Three-year outcomes of maintenance nortriptyline treatment in late-life depression: a study of two fixed plasma levels." <i>Am J Psychiatry</i> 156: 1177-1181.	4
Reynolds, C. F., rd, M. A. Butters, O. Lopez, B. G. Pollock, M. A. Dew, B. H. Mulsant, E. J. Lenze, M. Holm, J. C. Rogers, S. Mazumdar, P. R. Houck, A. Begley, S. Anderson, J. F. Karp, M. D. Miller, E. M. Whyte, J. Stack, A. Gildengers, K. Szanto, S. Bensasi, D. I. Kaufer, M. I. Kamboh and S. T. DeKosky (2011). "Maintenance treatment of depression in old age: a randomized, double-blind, placebo-controlled evaluation of the efficacy and safety of donepezil combined with antidepressant pharmacotherapy." <i>Arch Gen Psychiatry</i> 68: 51-60.	3
Reynolds, I. and F. C (1992). "Treatment of depression in special populations." <i>Journal of Clinical Psychiatry</i> 53: 45-53.	4
Reynolds, I., F. C, E. Frank, J. M. Perel, M. D. Miller, C. F. Paradis, J. A. Stack, B. G. Pollock, A. H. Rifai, C. Cornes, C. J. George, S. Mazumdar and D. J. Kupfer (1995). "Nortriptyline side effects during double-blind, randomized, placebo- controlled maintenance therapy in older depressed patients." <i>American Journal of Geriatric Psychiatry</i> 3: 170-175.	3

Ried, L. D., M. J. Tueth, E. Handberg, S. Kupfer and C. J. Pepine (2005). "A study of antihypertensive drugs and depressive symptoms (SADD-Sx) in patients treated with a calcium antagonist versus an atenolol hypertension treatment strategy in the International Verapamil SR-Trandolapril Study (INVEST)." <i>Psychosom Med</i> 67: 398-406.	4
Robert, P. and S. A. Montgomery (1995). "Citalopram in doses of 20-60 mg is effective in depression relapse prevention: a placebo-controlled 6 month study." <i>Int Clin Psychopharmacol</i> 10 Suppl 1: 29-35.	4
Robinson, D. S., D. L. Roberts, J. M. Smith, J. C. Stringfellow, S. B. Kaplita, J. A. Seminara and R. N. Marcus (1996). "The safety profile of nefazodone." <i>Journal of Clinical Psychiatry</i> 57: 31-38.	3
Robinson, M. J., T. O. Meyers, J. Raskin, P. Liu, S. Shoemaker and C. J. Nelson (2011). "Duloxetine versus placebo in the long-term treatment of patients with late-life major depression." <i>American Journal of Geriatric Psychiatry</i> 19(3): S103-S104.	9
Robinson, M., T. M. Oakes, J. Raskin, P. Liu, S. Shoemaker and J. C. Nelson (2014). "Acute and long-term treatment of late-life major depressive disorder: duloxetine versus placebo." <i>Am J Geriatr Psychiatry</i> 22(1): 34-45.	10
Robinson, R. G., S. K. Schultz, C. Castillo, T. Kopel, J. T. Kosier, R. M. Newman, K. Curdue, G. Petracca and S. E. Starkstein (2000). "Nortriptyline versus fluoxetine in the treatment of depression and in short-term recovery after stroke: A placebo-controlled, double-blind study." <i>American Journal of Psychiatry</i> 157: 351-359.	3
Rochon, P. and P. E. Lee (2005). "Citalopram did not differ from placebo for treatment of depression in patients greater than or equal to 75 years of age." <i>ACP Journal Club</i> 142(3): 79-79.	3
Roine, R. O., S. Kajaste and M. Kaste (1993). "Neuropsychological sequelae of cardiac arrest." <i>JAMA</i> 269: 237-242.	3
Romera, I., V. Perez, J. M. Menchon, A. Schacht, R. Papen, D. Neuhauser, M. Abbar, P. Svanborg and I. Gilaberte (2012). "Early switch strategy in patients with major depressive disorder: a double-blind, randomized study." <i>J Clin Psychopharmacol</i> 32(4): 479-486.	6
Rondanelli, M., A. Giacosa, A. Opizzi, C. Pelucchi, V. La, G. Montorfano, M. Negroni, B. Berra, P. Politi and A. M. Rizzo (2010). "Effect of omega-3 fatty acids supplementation on depressive symptoms and on health-related quality of life in the treatment of elderly women with depression: a double-blind, placebo-controlled, randomized clinical trial." <i>J Am Coll Nutr</i> 29: 55-64.	3
Rondanelli, M., A. Giacosa, A. Opizzi, C. Pelucchi, V. La, G. Montorfano, M. Negroni, B. Berra, P. Politi and A. M. Rizzo (2011). "Long chain omega 3 polyunsaturated fatty acids supplementation in the treatment of elderly depression: effects on depressive symptoms, on phospholipids fatty acids profile and on health-related quality of life." <i>J Nutr Health Aging</i> 15: 37-44.	3
Roose, S. (2002). "Treatment of depression in the ``old-old'': A randomized, double-blind, placebo-controlled trial of citalopram in patients at least 75 years of age." <i>European Neuropsychopharmacology</i> : S202.	3

Roose, S. P., A. H. Glassman, E. Attia and S. Woodring (1994). "Comparative efficacy of selective serotonin reuptake inhibitors and tricyclics in the treatment of melancholia." <i>Am J Psychiatry</i> 151(12): 1735-1739.	3
Roose, S. P., A. H. Glassman, E. Attia, S. Woodring, E. G. V. Giardina and J. T. Bigger Jr (1998). "Cardiovascular effects of fluoxetine in depressed patients with heart disease." <i>American Journal of Psychiatry</i> 155(5): 660-665.	3
Roose, S. P., J. C. Nelson, C. Salzman, S. B. Hollander, H. Rodrigues, S. Mirtazapine in the Nursing Home and Group (2003). "Open-label study of mirtazapine orally disintegrating tablets in depressed patients in the nursing home." <i>Curr Med Res Opin</i> 19: 737-746.	3
Roscoe, J. A., G. R. Morrow, J. T. Hickok, K. M. Mustian, J. J. Griggs, S. E. Matteson, P. Bushunow, R. Qazi and B. Smith (2005). "Effect of paroxetine hydrochloride (Paxil) on fatigue and depression in breast cancer patients receiving chemotherapy." <i>Breast Cancer Res Treat</i> 89: 243-249.	3
Rosen, J., B. G. Pollock, L. P. Altieri and E. A. Jonas (1993). "Treatment of nortriptyline's side effects in elderly patients: a double-blind study of bethanechol." <i>Am J Psychiatry</i> 150: 1249-1251.	7
Rosen, J., R. Sweet, B. G. Pollock and B. H. Mulsant (1993). "Nortriptyline in the hospitalized elderly: tolerance and side effect reduction." <i>Psychopharmacol Bull</i> 29: 327-331.	4
Rosenberg, C., N. Damsbo, E. Fuglum, L. V. Jacobsen and S. Horsgard (1994). "Citalopram and imipramine in the treatment of depressive patients in general practice. A Nordic multicentre clinical study." <i>Int Clin Psychopharmacol</i> 9 Suppl 1: 41-48.	3
Ross, J. (2008). "Discontinuation of lithium augmentation in geriatric patients with unipolar depression: a systematic review." <i>Can J Psychiatry</i> 53(2): 117-120.	9
Rossini, D., A. Serretti, L. Franchini, L. Mandelli, E. Smeraldi, R. De and R. Zanardi (2005). "Sertraline versus fluvoxamine in the treatment of elderly patients with major depression: a double-blind, randomized trial." <i>J Clin Psychopharmacol</i> 25: 471-475.	4
Rossini, D., L. Franchini, E. Smeraldi and R. Zanardi (2002). "Sertraline versus fluvoxamine in the treatment of elderly patients with major depression: A double-blind, randomised trial." <i>European Neuropsychopharmacology</i> : S250.	6
Roth, M., C. Q. Mountjoy and R. Amrein (1996). "Moclobemide in elderly patients with cognitive decline and depression: an international double-blind, placebo-controlled trial." <i>Br J Psychiatry</i> 168: 149-157.	6
Rouillon, F., D. Serrurier, H. D. Miller and M. J. Gerard (1991). "Prophylactic efficacy of maprotiline on unipolar depression relapse." <i>J Clin Psychiatry</i> 52: 423-431.	7
Ruddell, M., A. Spencer, K. Hill and A. House (2007). "Fluoxetine vs placebo for depressive symptoms after stroke: Failed randomised controlled trial." <i>Int J Geriatr Psychiatry</i> 22: 963-965.	9
Rudolph, R. L. and A. D. Feiger (1999). "A double-blind, randomized, placebo-controlled trial of once-daily venlafaxine extended release (XR) and fluoxetine for the treatment of depression." <i>J Affect Disord</i> 56(2-3): 171-181.	9

Rush, A. J., D. Warden, R. Wisniewski Stephen, M. Fava, H. Trivedi Madhukar, N. Gaynes Bradley and A. Nierenberg Andrew (2009). "STARD: Revising conventional wisdom." CNS Drugs 23: 627-647.	6
Rush, A. J., H. Trivedi Madhukar, J. Carmody Thomas, M. J. Donahue Rafe, L. Houser Trisha, C. Bolden-Watson, R. Batey Sharyn, A. Ascher John and A. Metz (2001). "Response in related to baseline anxiety levels in major depressive disorder treated with bupropion sustained release or sertraline." Neuropsychopharmacology 25: 131-138.	3
Rush, A. J., M. H. Trivedi, J. W. Stewart, A. A. Nierenberg, M. Fava, B. T. Kurian, D. Warden, D. W. Morris, J. F. Luther, M. M. Husain, I. A. Cook, R. C. Shelton, I. M. Lesser, S. G. Kornstein and S. R. Wisniewski (2011). "Combining medications to enhance depression outcomes (COMED): acute and long-term outcomes of a single-blind randomized study." Am J Psychiatry 168(7): 689-701.	6
Rush, A. J., S. R. Wisniewski, D. Warden, J. F. Luther, L. L. Davis, M. Fava, A. A. Nierenberg and M. H. Trivedi (2008). "Selecting among second-step antidepressant medication monotherapies: predictive value of clinical, demographic, or first-step treatment features." Arch Gen Psychiatry 65: 870-880.	9
Russell, J. M., L. M. Koran, J. Rush, R. M. Hirschfeld, W. Harrison, E. S. Friedman, S. Davis and M. Keller (2001). "Effect of concurrent anxiety on response to sertraline and imipramine in patients with chronic depression." Depress Anxiety 13(1): 18-27.	4
Russell, J. M., S. G. Kornstein, M. T. Shea, J. P. McCullough, W. M. Harrison, R. M. Hirschfeld and M. B. Keller (2003). "Chronic depression and comorbid personality disorders: response to sertraline versus imipramine." J Clin Psychiatry 64(5): 554-561.	3
Russell, J., J. Raskin, C. Wiltse, D. Walker and O. Brawman-Mintzer (2007). "Efficacy and tolerability of duloxetine treatment in elderly patients with major depressive disorder and concurrent anxiety symptoms." Psychiatry 4: 33-45.	9
Ruths, S., J. Straand, A. Nygaard Harald and D. Aarsland (2008). "Stopping antipsychotic drug therapy in demented nursing home patients: A randomized, placebo-controlled study-The Bergen District Nursing Home Study (BEDNURS)." Int J Geriatr Psychiatry 23: 889-895.	10
Sackeim, H. A., E. M. Dillingham, J. Prudic, T. Cooper, W. V. McCall, P. Rosenquist, K. Isenberg, K. Garcia, B. H. Mulsant and R. F. Haskett (2009). "Effect of concomitant pharmacotherapy on electroconvulsive therapy outcomes: short-term efficacy and adverse effects." Arch Gen Psychiatry 66(7): 729-737.	4
Saiz-Ruiz, J., J. M. Montes, E. Alvarez, S. Cervera, J. Giner, J. Guerrero, A. Seva, F. Dourdin and J. J. Lopez-Ibor (1998). "Tianeptine therapy for depression in the elderly." Progress in Neuro-Psychopharmacology and Biological Psychiatry 22: 319-329.	3
Sambunaris, A., A. Bose, C. P. Gommoll, C. Chen, W. M. Greenberg and D. V. Sheehan (2014). "A phase III, double-blind, placebo-controlled, flexible-dose study of levomilnacipran extended-release in patients with major depressive disorder." J Clin Psychopharmacol 34(1): 47-56.	3
Sarginson, J. E., L. C. Lazzeroni, H. S. Ryan, A. F. Schatzberg, G. M. Murphy and Jr (2010). "FKBP5 polymorphisms and antidepressant response in geriatric depression." Am J Med Genet B Neuropsychiatr Genet 153B: 554-560.	3

Sarginson, J. E., L. C. Lazzeroni, H. S. Ryan, B. D. Ershoff, A. F. Schatzberg and G. M. Murphy (2010). "ABCB1 (MDR1) polymorphisms and antidepressant response in geriatric depression." <i>Pharmacogenetics and Genomics</i> 20(8): 467-475.	3
Sarris, J., I. P. G, O. Vitolo, M. Fava and D. Mischoulon (2014). "S-adenosyl methionine (SAME) versus escitalopram and placebo in major depression RCT: efficacy and effects of histamine and carnitine as moderators of response." <i>J Affect Disord</i> 164: 76-81.	3
Savaskan, E., S. E. Muller, A. Bohringer, A. Schulz and H. Schachinger (2008). "Antidepressive therapy with escitalopram improves mood, cognitive symptoms, and identity memory for angry faces in elderly depressed patients." <i>International Journal of Neuropsychopharmacology</i> 11(3): 381-388.	4
Schmidt, M. E., M. Fava, J. M. Robinson and R. Judge (2000). "The efficacy and safety of a new enteric-coated formulation of fluoxetine given once weekly during the continuation treatment of major depressive disorder." <i>Journal of Clinical Psychiatry</i> 61: 851-857.	4
Schmidt, U. and H. Sommer (1993). "Extract of St. John's wort in the treatment of depression - Attention and reaction remain unimpaired." <i>Fortschr Med</i> 111: 37-40.	11
Schmidt, U., G. Harrer, U. Kuhn and A. Biller (1999). "Equivalence comparison of the St John's wort extract LoHyp-57 versus fluoxetine HCL." <i>Zeitschrift fur Phytotherapie</i> 20: 89-90.	11
Schneider, L. S. (2004). "Venlafaxine offers no benefit over sertraline and is less well tolerated in depressed nursing home residents." <i>Evidence Based Mental Health</i> 7: 47-47.	3
Schneider, L. S., G. W. Small and C. M. Clary (2001). "Estrogen replacement therapy and antidepressant response to sertraline in older depressed women." <i>American Journal of Geriatric Psychiatry</i> 9: 393-399.	7
Schneider, L. S., G. W. Small, S. H. Hamilton, A. Bystritsky, C. B. Nemeroff and B. S. Meyers (1997). "Estrogen replacement and response to fluoxetine in a multicenter geriatric depression trial. Fluoxetine Collaborative Study Group." <i>Am J Geriatr Psychiatry</i> 5: 97-106.	9
Schneider, L. S., J. C. Nelson, C. M. Clary, P. Newhouse, K. R. Krishnan, T. Shiovitz, K. Weihs and G. Sertraline Elderly Depression Study (2003). "An 8-week multicenter, parallel-group, double-blind, placebo-controlled study of sertraline in elderly outpatients with major depression." <i>Am J Psychiatry</i> 160(7): 1277-1285.	9
Schneider, L. S., T. B. Cooper, J. A. Severson, T. Zemplenyi and R. B. Sloane (1988). "Electrocardiographic changes with nortriptyline and 10-hydroxynortriptyline in elderly depressed outpatients." <i>J Clin Psychopharmacol</i> : 402-408.	6
Schone, W. and M. Ludwig (1994). "[Paroxetine in the treatment of depression in geriatric patients--a double-blind comparative study with fluoxetine]." <i>Fortschr Neurol Psychiatr</i> 62 Suppl 1: 16-18.	11
Schone, W. and M. Ludwig (1994). "Paroxetine in the treatment of geriatric depressed patients - A double-blind comparison with fluoxetine." <i>Fortschritte der Neurologie Psychiatrie</i> 62: 16-18.	3
Schrader, E. (2000). "Equivalence of St John's wort extract (Ze 117) and fluoxetine: a randomized, controlled study in mild-moderate depression." <i>Int Clin Psychopharmacol</i> 15: 61-68.	9

Schutter, D. J. (2009). "Antidepressant efficacy of high-frequency transcranial magnetic stimulation over the left dorsolateral prefrontal cortex in double-blind sham-controlled designs: a meta-analysis." <i>Psychol Med</i> 39(1): 65-75.	3 och 4
Schwartz, T. L., G. S. Nasra, M. Chilton, D. Kang and H. Kumaresan (2007). "Aripiprazole augmentation of selective serotonin or serotonin norepinephrine reuptake inhibitors in the treatment of major depressive disorder." <i>Primary Psychiatry</i> 14(1): 67-69.	3
Schweitzer, I., J. Tiller, K. Maguire and B. Davies (1989). "Treatment of atypical depression with moclobemide: A sequential double controlled study." <i>Int J Clin Pharmacol Res</i> 9: 111-117.	9
Schweizer, E., K. Rickels and H. Hassman (1994) "A double-blind, placebo-controlled comparison of imipramine and buspirone in the treatment of major depression in the elderly in the community." <i>Psychopharmacol Bull</i> , 639.	9
Schweizer, E., K. Rickels, H. Hassman and F. Garcia-Espana (1998). "Buspirone and imipramine for the treatment of major depression in the elderly." <i>J Clin Psychiatry</i> 59(4): 175-183.	7
Septien-Velez, L., B. Pitrosky, S. K. Padmanabhan, J. M. Germain and K. A. Tourian (2007). "A randomized, double-blind, placebo-controlled trial of desvenlafaxine succinate in the treatment of major depressive disorder." <i>Int Clin Psychopharmacol</i> 22: 338-347.	3
Serebruany, V. L., A. H. Glassman, A. I. Malinin, C. B. Nemeroff, D. L. Musselman, Z. Van, T. L, M. S. Finkel, K. R. R. Krishnan, M. Gaffney, W. Harrison, R. M. Califf and C. M. O'Connor (2003). "Platelet/endothelial biomarkers in depressed patients treated with the selective serotonin reuptake inhibitor sertraline after acute coronary events: The Sertraline Antidepressant Heart Attack Randomized Trial (SADHART) platelet substudy." <i>Circulation</i> 108: 939-944.	3
Serrano-Duenas, M. (2002). "[A comparison between low doses of amitriptyline and low doses of fluoxetine used in the control of depression in patients suffering from Parkinson's disease]." <i>Rev Neurol</i> 35(11): 1010-1014.	11
Settle, E. C., S. M. Stahl, S. R. Batey, J. A. Johnston and J. A. Ascher (1999). "Safety profile of sustained-release bupropion in depression: results of three clinical trials." <i>Clin Ther</i> 21: 454-463.	3
Shangina, O. A., V. I. Kostin and L. I. Koturga (2005). "Effects of therapy with various groups of antianginal agents on life quality of the elderly with coronary heart disease and comorbid depression." <i>Klin Med (Mosk)</i> : 34-37.	11
Sheehan, D., G. C. Dunbar and D. L. Fuell (1992). "The effect of paroxetine on anxiety and agitation associated with depression." <i>Psychopharmacol Bull</i> 28: 139-144.	3
Sheffrin, M., H. C. Driscoll, E. J. Lenze, B. H. Mulsant, B. G. Pollock, M. D. Miller, M. A. Butters, M. A. Dew and C. F. Reynolds, 3rd (2009). "Pilot study of augmentation with aripiprazole for incomplete response in late-life depression: getting to remission." <i>J Clin Psychiatry</i> 70(2): 208-213.	3
Sheikh, J. I., E. L. Cassidy, P. M. Doraiswamy, R. M. Salomon, M. Hornig, P. J. Holland, F. S. Mandel, C. M. Clary and T. Burt (2004). "Efficacy, safety, and tolerability of sertraline in patients with late-life depression and comorbid medical illness." <i>J Am Geriatr Soc</i> 52: 86-92.	3
Sheikh, J., P. Newhouse, T. Burt and C. Clary (2002). "Remission rates in elderly depressed patients treated with sertraline." <i>European Neuropsychopharmacology</i> : S232.	3

Shiovitz, T., W. M. Greenberg, C. Chen, G. Forero and C. P. Gommoll (2014). "A Randomized, Double-blind, Placebo-controlled Trial of the Efficacy and Safety of Levomilnacipran ER 40-120mg/day for Prevention of Relapse in Patients with Major Depressive Disorder." <i>Innov Clin Neurosci</i> 11(1-2): 10-22.	3
Shores, M. M., D. R. Kivlahan, T. I. Sadak, E. J. Li and A. M. Matsumoto (2009). "A randomized, double-blind, placebo-controlled study of testosterone treatment in hypogonadal older men with subthreshold depression (dysthymia or minor depression)." <i>J Clin Psychiatry</i> 70: 1009-1016.	7
Shrivastava, R. K., S. P. Shrivastava, J. Landinez, C. D. Pitts, D. M. Schaefer, A. Lipschitz and M. Iyengar (2005). "Efficacy and tolerability of low doses of paroxetine CR in elderly outpatients with major depression." <i>Neuropsychopharmacology</i> : S239.	4
Shu, L., A. H. Sulaiman, Y. S. Huang, C. Fones Soon Leng, V. S. Crutel and Y. S. Kim (2014). "Comparable efficacy and safety of 8 weeks treatment with agomelatine 25-50mg or fluoxetine 20-40mg in Asian out-patients with major depressive disorder." <i>Asian J Psychiatr</i> 8: 26-32.	3
Shulman, K. I. (2006). "Continuing paroxetine treatment reduces recurrence of major depression in the elderly." <i>Evidence Based Mental Health</i> 9(4): 101-101.	7
Siegfried, K. and M. O'Connolly (1986). "Cognitive and psychomotor effects of different antidepressants in the treatment of old age depression." <i>Int Clin Psychopharmacol</i> 1(3): 231-243.	3
Signorovitch, J., K. Ramakrishnan, R. Ben-Hamadi, A. P. Yu, E. Q. Wu, H. Dworak and M. H. Erder (2011). "Remission of major depressive disorder without adverse events: a comparison of escitalopram versus serotonin norepinephrine reuptake inhibitors." <i>Curr Med Res Opin</i> 27: 1089-1096.	3
Silverstone, P. H. and A. Ravindran (1999). "Once-daily venlafaxine extended release (XR) compared with fluoxetine in outpatients with depression and anxiety. Venlafaxine XR 360 Study Group." <i>J Clin Psychiatry</i> 60: 22-28.	4
Silverstone, P. H., R. Entsuah and D. Hackett (2002). "Two items on the Hamilton Depression rating scale are effective predictors of remission: comparison of selective serotonin reuptake inhibitors with the combined serotonin/norepinephrine reuptake inhibitor, venlafaxine." <i>Int Clin Psychopharmacol</i> 17: 273-280.	6
Simon, G. E., D. Revicki, J. Heiligenstein, L. Grothaus, M. VonKorff, W. J. Katon and T. R. Hylan (2000). "Recovery from depression, work productivity, and health care costs among primary care patients." <i>Gen Hosp Psychiatry</i> 22: 153-162.	6
Simon, G. E., J. H. Heiligenstein, L. Grothaus, W. Katon and D. Revicki (1998). "Should anxiety and insomnia influence antidepressant selection: a randomized comparison of fluoxetine and imipramine." <i>J Clin Psychiatry</i> 59: 49-55.	6
Simon, G. E., M. VonKorff, J. H. Heiligenstein, D. A. Revicki, L. Grothaus, W. Katon and E. H. Wagner (1996). "Initial antidepressant choice in primary care: Effectiveness and cost of fluoxetine vs tricyclic antidepressants." <i>Journal of the American Medical Association</i> 275: 1897-1902.	4
Simons, J. A. (1996). "Fluoxetine in Parkinson's disease." <i>Movement Disorders</i> 11: 581-582.	3

Sinclair, J. M., M. R. Walsh, J. C. Valle-Jones and A. A. Schiff (1975). "Treatment of anxiety/depressive conditions in the elderly: a double-blind comparative study of Motival and amitriptyline." <i>Age Ageing</i> : 226-231.	6
Slimmer, L. (2003). "A collaborative care management programme in a primary care setting was effective for older adults with late life depression." <i>Evidence Based Nursing</i> 6(3): 91-91.	3
Small, G. W., L. S. Schneider, S. H. Hamilton, A. Bystritsky, B. S. Meyers and C. B. Nemeroff (1996). "Site variability in a multisite geriatric depression trial." <i>Int J Geriatr Psychiatry</i> 11: 1089-1095.	3
Small, G. W., M. Birkett, B. S. Meyers, L. M. Koran, A. Bystritsky and C. B. Nemeroff (1996). "Impact of physical illness on quality of life and antidepressant response in geriatric major depression. Fluoxetine Collaborative Study Group." <i>J Am Geriatr Soc</i> 44: 1220-1225.	3
Small, G. W., S. H. Hamilton, A. Bystritsky, B. S. Meyers and C. B. Nemeroff (1995). "Clinical response predictors in a double-blind, placebo-controlled trial of fluoxetine for geriatric major depression. Fluoxetine Collaborative Study Group." <i>Int Psychogeriatr</i> 7 Suppl: 41-53.	4
Smeraldi, E. (1998). "Amisulpride versus fluoxetine in patients with dysthymia or major depression in partial remission: a double-blind, comparative study." <i>J Affect Disord</i> 48(1): 47-56.	3
Smith, D. F., E. Stromgren and H. N. Petersen (1984). "Lack of effect of tryptophan treatment in demented gerontopsychiatric patients. A double-blind, crossover-controlled study." <i>Acta Psychiatrica Scandinavica</i> 70: 470-477.	3
Smith, G. S., C. F. Reynolds, 3rd, P. R. Houck, M. A. Dew, J. Ginsberg, Y. Ma, B. H. Mulsant and B. G. Pollock (2009). "Cerebral glucose metabolic response to combined total sleep deprivation and antidepressant treatment in geriatric depression: a randomized, placebo-controlled study." <i>Psychiatry Res</i> 171(1): 1-9.	6
Sneed, J. R., M. E. Culang, J. G. Keilp, B. R. Rutherford, D. P. Devanand and S. P. Roose (2010). "Antidepressant medication and executive dysfunction: a deleterious interaction in late-life depression." <i>Am J Geriatr Psychiatry</i> 18(2): 128-135.	6
Sneed, J. R., S. P. Roose, J. G. Keilp, K. R. Krishnan, G. S. Alexopoulos and H. A. Sackeim (2007). "Response inhibition predicts poor antidepressant treatment response in very old depressed patients." <i>Am J Geriatr Psychiatry</i> 15: 553-563.	7
Soares, C. N., M. E. Thase, A. Clayton, C. J. Guico-Pabia, K. Focht, Q. Jiang, S. G. Kornstein, P. T. Ninan and C. P. Kane (2011). "Open-label treatment with desvenlafaxine in postmenopausal women with major depressive disorder not responding to acute treatment with desvenlafaxine or escitalopram." <i>CNS Drugs</i> 25: 227-238.	3
Solai, L. K., B. G. Pollock, B. H. Mulsant, R. F. Frye, M. D. Miller, R. A. Sweet, M. Kirshner, D. Sorisio, A. Begley, C. F. Reynolds and rd (2002). "Effect of nortriptyline and paroxetine on CYP2D6 activity in depressed elderly patients." <i>J Clin Psychopharmacol</i> 22: 481-486.	3
Sommer, H. and G. Harrer (1994). "Placebo-controlled double-blind study examining the effectiveness of an hypericum preparation in 105 mildly depressed patients." <i>J Geriatr Psychiatry Neurol</i> 7 Suppl 1: S9-11.	4

Sondergaard, M. P., J. O. Jarden, K. Martiny, G. Andersen and P. Bech (2006). "Dose response to adjunctive light therapy in citalopram-treated patients with post-stroke depression. A randomised, double-blind pilot study." <i>Psychother Psychosom</i> 75(4): 244-248.	9
Stage, K. B., A. Danish University and Group (2005). "Orthostatic side effects of clomipramine and moclobemide during treatment for depression." <i>Nord J Psychiatry</i> 59: 298-301.	5
Stassen, H. H., J. Angst and A. Delini-Stula (1999). "Fluoxetine versus moclobemide: cross-comparison between the time courses of improvement." <i>Pharmacopsychiatry</i> 32: 56-60.	3
Steffens, D. C., D. R. McQuoid and K. R. R. Krishnan (2003). "Partial response as a predictor of outcome in geriatric depression." <i>American Journal of Geriatric Psychiatry</i> 11: 340-348.	3
Steffens, D. C., J. C. Nelson, J. M. Eudicone, C. Andersson, H. Yang, Q. V. Tran, R. A. Forbes, B. X. Carlson and R. M. Berman (2011). "Efficacy and safety of adjunctive aripiprazole in major depressive disorder in older patients: a pooled subpopulation analysis." <i>Int J Geriatr Psychiatry</i> 26: 564-572.	3
Steffens, D. C., M. Snowden, M. Y. Fan, H. Hendrie, W. J. Katon, J. Unutzer and I. Investigators (2006). "Cognitive impairment and depression outcomes in the IMPACT study." <i>Am J Geriatr Psychiatry</i> 14(5): 401-409.	3
Steinberg, M., C. A. Munro, Q. Samus, P. V. Rabins, J. Brandt and C. G. Lyketsos (2004). "Patient predictors of response to treatment of depression in Alzheimer's disease: the DIADS study." <i>Int J Geriatr Psychiatry</i> 19: 144-150.	3
Stewart, J. W., P. J. McGrath and F. M. Quitkin (1992). "Can mildly depressed outpatients with atypical depression benefit from antidepressants?" <i>Am J Psychiatry</i> 149(5): 615-619.	3
Stewart, J. W., P. J. McGrath, C. Blondeau, D. A. Deliyannides, D. Hellerstein, S. Norris, J. Amat, D. J. Pilowsky, P. Tessier, L. Laberge, D. O'Shea, Y. Chen, A. Withers, R. Bergeron and P. Blier (2014). "Combination antidepressant therapy for major depressive disorder: speed and probability of remission." <i>J Psychiatr Res</i> 52: 7-14.	3
Stewart, T. D., A. Hatch, K. Largay, J. J. Sheehan, S. V. Marler, R. M. Berman and J. C. Nelson (2014). "Effect of symptom severity on efficacy and safety of aripiprazole adjunctive to antidepressant monotherapy in major depressive disorder: a pooled analysis." <i>J Affect Disord</i> 162: 20-25.	3
Stockler, M. R., R. O'Connell, A. K. Nowak, D. Goldstein, J. Turner, N. R. Wilcken, D. Wyld, E. A. Abdi, A. Glasgow, P. J. Beale, M. Jefford, H. Dhillon, S. Heritier, C. Carter, I. B. Hickie, R. J. Simes, o. Zoloft's Effects, Symptoms, T. survival Time and Group (2007). "Effect of sertraline on symptoms and survival in patients with advanced cancer, but without major depression: a placebo-controlled double-blind randomised trial." <i>Lancet Oncol</i> 8: 603-612.	9
Stoudemire, A., C. D. Hill and R. Morris (1991) "Cognitive outcome following tricyclic and electroconvulsive treatment of major depression in the elderly." <i>American Journal of Psychiatry</i> , 1336-1340.	5
Strauss, D. and K. Solomon (1983). "Psychopharmacologic intervention for depression in the elderly." <i>Clinical Gerontologist</i> 2: 3-29.	3

Streim, J. E., D. W. Oslin, I. R. Katz, B. D. Smith, S. DiFilippo, T. B. Cooper and T. Ten Have (2000). "Drug treatment of depression in frail elderly nursing home residents." <i>Am J Geriatr Psychiatry</i> 8(2): 150-159.	3
Strik, J. J. M. H., A. Honig, E. Klinkenberg, J. Dijkstra and J. Jolles (2006). "Cognitive performance following fluoxetine treatment in depressed patients post myocardial infarction." <i>Acta Neuropsychiatrica</i> 18(1): 1-6.	3
Strik, J. J., A. Honig, R. Lousberg, A. H. Lousberg, E. C. Cheriex, H. G. Tuynman-Qua, P. M. Kuijpers, H. J. Wellens, P. Van and M. H (2000). "Efficacy and safety of fluoxetine in the treatment of patients with major depression after first myocardial infarction: findings from a double-blind, placebo-controlled trial." <i>Psychosom Med</i> 62: 783-789.	3
Strik, J. J., A. Honig, R. Lousberg, E. C. Cheriex and H. M. Van Praag (1998). "Cardiac side-effects of two selective serotonin reuptake inhibitors in middle-aged and elderly depressed patients." <i>Int Clin Psychopharmacol</i> 13(6): 263-267.	3
Stryjer, R., R. D. Strous, G. Shaked, F. Bar, B. Feldman, M. Kotler, L. Polak, S. Rosenzwaig and A. Weizman (2003). "Amantadine as augmentation therapy in the management of treatment-resistant depression." <i>Int Clin Psychopharmacol</i> 18(2): 93-96.	4
Suo, A. Q. and L. Dang (2006). "Treatment of accompanied depression in subcortical arteriosclerotic encephalopathy." <i>Chinese Journal of Clinical Rehabilitation</i> 10: 7-9.	4
Swenson, J. R., C. M. O'Connor, D. Barton, Z. Van, T. L, K. Swedberg, L. M. Forman, M. Gaffney and A. H. Glassman (2003). "Influence of depression and effect of treatment with sertraline on quality of life after hospitalization for acute coronary syndrome." <i>American Journal of Cardiology</i> 92: 1271-1276.	4
Szadoczky, E. and J. Furedi (2002). "Efficacy and acceptability of tianeptine and sertraline in the acute treatment phase of depression." <i>Encephale</i> 28: 343-349.	11
Szanto, K., B. H. Mulsant, P. R. Houck, M. A. Dew, A. Dombrovski, B. G. Pollock, I. Reynolds and F. C (2007). "Emergence, persistence, and resolution of suicidal ideation during treatment of depression in old age." <i>J Affect Disord</i> 98: 153-161.	9
Szegedi, A., A. Hillert, H. Wetzel, E. Klieser, W. Gaebel and O. Benkert (1997). "Pramipexole, a dopamine agonist, in major depression: Antidepressant effects and tolerability in an open-label study with multiple doses." <i>Clin Neuropharmacol</i> 20: S36-S45.	7
Szegedi, A., H. Wetzel, D. Angersbach, G. C. Dunbar, H. Schwarze, M. Philipp and O. Benkert (1997). "A double-blind study comparing paroxetine and maprotiline in depressed outpatients." <i>Pharmacopsychiatry</i> 30(3): 97-105.	3
Szegedi, A., H. Wetzel, D. Angersbach, M. Philipp and O. Benkert (1997). "Response to treatment in minor and major depression: results of a double-blind comparative study with paroxetine and maprotiline." <i>J Affect Disord</i> 45: 167-178.	9
Szegedi, A., M. J. Muller, I. Anghelescu, C. Klawe, R. Kohnen and O. Benkert (2003). "Early improvement under mirtazapine and paroxetine predicts later stable response and remission with high sensitivity in patients with major depression." <i>J Clin Psychiatry</i> 64(4): 413-420.	4

Tadic, A., I. Helmreich, R. Mergl, M. Hautzinger, R. Kohnen, V. Henkel and U. Hegerl (2010). "Early improvement is a predictor of treatment outcome in patients with mild major, minor or subsyndromal depression." <i>J Affect Disord</i> 120(1-3): 86-93.	3
Tajalizadekhoob, Y., F. Sharifi, H. Fakhrzadeh, M. Mirarefin, M. Ghaderpanahi, Z. Badamchizade and S. Azimipour (2011). "The effect of low-dose omega 3 fatty acids on the treatment of mild to moderate depression in the elderly: a double-blind, randomized, placebo-controlled study." <i>Eur Arch Psychiatry Clin Neurosci</i> 261: 539-549.	10
Tammaro, A., M. T. Picceo, P. Gemmellaro and O. Bonaccorso (1977). "Camazepam versus placebo. A double-blind clinical study on geriatric patients suffering from psychic complaints. Short communication." <i>Arzneimittel-Forschung/Drug Research</i> 27(11): 2177-2178.	4
Tan, R. S. (1998). "Prescribing less than recommended doses may be acceptable for geriatric patients." <i>J Am Geriatr Soc</i> 46: 536.	3
Tan, R. S., R. J. Barlow, C. Abel, S. Reddy, A. J. Palmer, A. E. Fletcher, C. G. Nicholl, B. M. Pitt and C. J. Bulpitt (1994). "The effect of low dose lofepramine in depressed elderly patients in general medical wards." <i>Br J Clin Pharmacol</i> 37: 321-324.	3
Taragano, F. E., C. G. Lyketsos, C. A. Mangone, R. F. Allegri and E. Comesana-Diaz (1997). "A double-blind, randomized, fixed-dose trial of fluoxetine vs. amitriptyline in the treatment of major depression complicating Alzheimer's disease." <i>Psychosomatics</i> 38(3): 246-252.	4
Tauscher-Wisniewski, S., D. Disch, J. Plewes, S. Ball, Beasley, Jr. and M. C (2007). "Evaluating suicide-related adverse events in clinical trials of fluoxetine treatment in adults for indications other than major depressive disorder." <i>Psychol Med</i> 37: 1585-1593.	9
Tedeschini, E., Y. Levkovitz and G. Papakostas (2010). "Antidepressants for late-life depression: A meta analysis of placebo-controlled randomized trials." <i>European Neuropsychopharmacology</i> 20: S382-S383.	7
Teper, E. and A. J. Thomas (2006). "Is depression different in older adults?" <i>Aging Health</i> 2: 905-915.	9
Teri, L., B. V. Reifler, R. C. Veith, R. Barnes, E. White, P. McLean and M. Raskind (1991). "Imipramine in the treatment of depressed Alzheimer's patients: impact on cognition." <i>J Gerontol</i> 46: P372-377.	3
Terra, J. L. and S. A. Montgomery (1998). "Fluvoxamine prevents recurrence of depression: results of a long-term, double-blind, placebo-controlled study." <i>Int Clin Psychopharmacol</i> 13: 55-62.	3
Thase, M. E., S. A. Corya, O. Osuntokun, M. Case, D. B. Henley, T. M. Sanger, S. B. Watson and S. Dube (2007). "A randomized, double-blind comparison of olanzapine/fluoxetine combination, olanzapine, and fluoxetine in treatment-resistant major depressive disorder." <i>J Clin Psychiatry</i> 68: 224-236.	3
Thase, M.E., Mahableshwarkar, A. and Dragheim, M., The efficacy of vortioxetine vs placebo in the treatment of adults with major depressive disorder: Patient level data from 10 short-term studies and a meta-analysis, <i>Neuropsychopharmacology</i> , 38 (2013) S363-S364. Accession:	9

Thomas, L., B. H. Mulsant, F. X. Solano, A. M. Black, S. Bensasi, T. Flynn, J. S. Harman, B. L. Rollman, E. P. Post, B. G. Pollock, I. Reynolds and F. C (2002). "Response speed and rate of remission in primary and specialty care of elderly patients with depression." American Journal of Geriatric Psychiatry 10: 583-591.	3
Thompson, L. W., D. Gallagher and R. Czirr (1988). "Personality disorder and outcome in the treatment of late-life depression." J Geriatr Psychiatry 21(2): 133-153.	9
Thompson, T. R., L. E. Smith, F. A. Marsteller, J. L. Woodard, V. L. Phillips, W. V. McCall and W. M. McDonald (2001) "Acute and continuation treatment of depression in the elderly." 154th Annual Meeting of the American Psychiatric Association; 2001 May 5-10; New Orleans LA, Nr834.	3
Tignol, J., J. Pujol-Domenech, J. P. Chartres, J. M. Leger, Y. Pletan, I. Tonelli, A. Tournoux and N. Pezous (1998). "Double-blind study of the efficacy and safety of milnacipran and imipramine in elderly patients with major depressive episode." Acta Psychiatr Scand 97(2): 157-165.	9
Tiller, J. W. G., I. Schweitzer, K. Maguire and B. M. Davies (1988). "Moclobemide and mianserin for depression in the elderly." Pharmacological Research Communications 20: 149.	3
Tiller, J., K. Maguire and B. Davies (1990). "A sequential double-blind controlled study of moclobemide and mianserin in elderly depressed patients." Int J Geriatr Psychiatry 5: 199-204.	3
Tohen, M., M. Case, M. H. Trivedi, M. E. Thase, S. J. Burke and T. M. Durell (2010). "Olanzapine/fluoxetine combination in patients with treatment-resistant depression: rapid onset of therapeutic response and its predictive value for subsequent overall response in a pooled analysis of 5 studies." J Clin Psychiatry 71(4): 451-462.	7
Tollefson, G. D. and S. L. Holman (1993). "Analysis of the Hamilton Depression Rating Scale factors from a double-blind, placebo-controlled trial of fluoxetine in geriatric major depression." Int Clin Psychopharmacol 8(4): 253-259.	3
Tollefson, G. D. and S. W. Andersen (1999). "Should we consider mood disturbance in schizophrenia as an important determinant of quality of life?" J Clin Psychiatry 60 Suppl 5: 23-29; discussion 30.	3
Tollefson, G. D., J. C. Bosomworth, J. H. Heiligenstein, J. H. Potvin and S. Holman (1995). "A double-blind, placebo-controlled clinical trial of fluoxetine in geriatric patients with major depression. The Fluoxetine Collaborative Study Group." Int Psychogeriatr 7(1): 89-104.	3
Tollefson, G., D. Bosomvort, D. Kheiligenstein, D. Potvin and S. Kholman (1997) "A double-blind, placebo-controlled clinical trial of fluoxetine in geriatric patients with major depression." Sotsialnaia i Klinicheskaiia Psichatriia, 79-92.	3
Tourigny-Rivard, M. F., V. K. Nair and P. Vincent (1996). "Fluvoxamine vs desipramine in elderly depressed patients." Xth World Congress of Psychiatry, Madrid, Spain.	3
Trivedi, M. H., A. J. Rush, T. J. Carmody, R. M. J. Donahue, C. Bolden-Watson, T. L. Houser and A. Metz (2001). "Do bupropion SR and sertraline differ in their effects on anxiety in depressed patients?" Journal of Clinical Psychiatry 62(10): 776-781.	3
Truffinet, P., F. Rouillon, D. Serrurier and R. Phillips (1990). "Relapses of unipolar depression in the elderly and the efficacy of maprotiline." Psychologie Medicale: 769-777.	11

Uher, R., A. Farmer, N. Henigsberg, M. Rietschel, O. Mors, W. Maier, D. Kozel, J. Hauser, D. Souery, A. Placentino, J. Strohmaier, N. Perroud, A. Zobel, A. Rajewska-Rager, M. Z. Dernovsek, E. R. Larsen, P. Kalember, C. Giovannini, M. Barreto, P. McGuffin and K. J. Aitchison (2009). "Adverse reactions to antidepressants." <i>Br J Psychiatry</i> 195(3): 202-210.	7
Uher, R., W. Maier, J. Hauser, A. Marusic, C. Schmael, O. Mors, N. Henigsberg, D. Souery, A. Placentino, M. Rietschel, A. Zobel, M. Dmitrzak-Weglacz, A. Petrovic, L. Jorgensen, P. Kalember, C. Giovannini, M. Barreto, A. Elkin, S. Landau, A. Farmer, K. J. Aitchison and P. McGuffin (2009). "Differential efficacy of escitalopram and nortriptyline on dimensional measures of depression." <i>Br J Psychiatry</i> 194(3): 252-259.	4
Wade, A. G., T. E. Schlaepfer, H. F. Andersen and C. D. Kilts (2009). "Clinical milestones predict symptom remission over 6-month and choice of treatment of patients with major depressive disorder (MDD)." <i>J Psychiatr Res</i> 43(5): 568-575.	3
Wade, A., G. M. Crawford, M. Angus, R. Wilson and L. Hamilton (2003). "A randomized, double-blind, 24-week study comparing the efficacy and tolerability of mirtazapine and paroxetine in depressed patients in primary care." <i>Int Clin Psychopharmacol</i> 18: 133-141.	4
Wade, A., K. Gembert and I. Florea (2007). "A comparative study of the efficacy of acute and continuation treatment with escitalopram versus duloxetine in patients with major depressive disorder." <i>Curr Med Res Opin</i> 23(7): 1605-1614.	3
Wade, A., L. Michael and H. Bang (2002). "Escitalopram 10 mg/day is effective and well tolerated in a placebo-controlled study in depression in primary care." <i>Int Clin Psychopharmacol</i> 17: 95-102.	7
Wade, A., N. Despiegel and E. H. Reines (2006). "Escitalopram in the long-term treatment of major depressive disorder." <i>Annals of Clinical Psychiatry</i> 18: 83-89.	3
Wakelin Jennifer, S. (1986). "Fluvoxamine in the treatment of the older depressed patient; double-blind, placebo-controlled data." <i>Int Clin Psychopharmacol</i> 1: 221-230.	3
Valenstein, M., H. M. Kim, D. Ganoczy, D. Eisenberg, P. N. Pfeiffer, K. Downing, K. Hoggatt, M. Ilgen, K. L. Austin, K. Zivin, F. C. Blow and J. F. McCarthy (2012). "Antidepressant agents and suicide death among US Department of Veterans Affairs patients in depression treatment." <i>J Clin Psychopharmacol</i> 32(3): 346-353.	6
Wallace, A. E., L. L. Kofoed and A. N. West (1995). "Double-blind, placebo-controlled trial of methylphenidate in older, depressed, medically ill patients." <i>Am J Psychiatry</i> 152(6): 929-931.	9
Walters, G., C. F. Reynolds, 3rd, B. H. Mulsant and B. G. Pollock (1999). "Continuation and maintenance pharmacotherapy in geriatric depression: an open-trial comparison of paroxetine and nortriptyline in patients older than 70 years." <i>J Clin Psychiatry</i> 60 Suppl 20: 21-25.	3
van de Rest, O., J. M. Geleijnse, F. J. Kok, W. A. van Staveren, W. H. Hoefnagels, A. T. Beekman and L. C. de Groot (2008). "Effect of fish-oil supplementation on mental well-being in older subjects: a randomized, double-blind, placebo-controlled trial." <i>Am J Clin Nutr</i> 88(3): 706-713.	3

Van Den, B., W. W, T. K. Birkenhager, P. G. H. Mulder, J. A. Bruijn and P. Moleman (2006). "Imipramine is effective in preventing relapse in electroconvulsive therapy-responsive depressed inpatients with prior pharmacotherapy treatment failure: A randomized, placebo-controlled trial." <i>Journal of Clinical Psychiatry</i> 67: 263-268.	6
van Iersel, M. B., S. U. Zuidema, R. T. C. M. Koopmans, F. R. J. Verhey and M. G. M. O. Rikkert (2005). "Antipsychotics for Behavioural and Psychological Problems in Elderly People with Dementia: A Systematic Review of Adverse Events." <i>Drugs Aging</i> 22(10): 845-858.	4
van, H. and M. Zivkov (1996). "Pharmacological treatment of depression in cancer patients. A placebo-controlled study of mianserin." <i>Br J Psychiatry</i> 169: 440-443.	6
van, K., A. L, R. J. Laheij, N. Aparicio, B. De, A. W, H. Van den, A. C. Tan, B. J. Witteman and J. B. Jansen (2008). "Effect of the antidepressant venlafaxine in functional dyspepsia: a randomized, double-blind, placebo-controlled trial." <i>Clin Gastroenterol Hepatol</i> 6: 746-752; quiz 718.	4
van, M., P. J, J. de, A. Honig, A. H. Schene, A. M. Kuyper, H. J. Crijns, A. Schins, D. Tulner, B. van den, P. M, J. Ormel and M.-I. investigators (2007). "Effects of antidepressant treatment following myocardial infarction." <i>Br J Psychiatry</i> 190: 460-466.	5
van, Z., T. L, F. Lesperance, N. Frasure-Smith, A. I. Malinin, D. Atar, M. A. Laliberte and V. L. Serebruany (2009). "Platelet and endothelial activity in comorbid major depression and coronary artery disease patients treated with citalopram: the Canadian Cardiac Randomized Evaluation of Antidepressant and Psychotherapy Efficacy Trial (CREATE) biomarker sub-study." <i>J Thromb Thrombolysis</i> 27: 48-56.	5
Wang, X., Z. Tan, Z. Wu, J. Gao and M. Feng (2003). "The effects of anti-depression therapy on post-stroke depression and neurologic rehabilitation in the elderly patients." <i>Chinese Journal of Geriatrics</i> : 270-273.	11
Vanna, M., J. Kummer, A. Agnoli, P. Gentili, A. Lorizio and R. Anand (1990). "Moclobemide compared with second-generation antidepressants in elderly people." <i>Acta Psychiatr Scand Suppl</i> : 64-66.	3
Varia, I., S. Venkataraman, C. Hellegers, K. Gersing and P. M. Doraiswamy (2007). "Effect of mirtazapine orally disintegrating tablets on health-related quality of life in elderly depressed patients with comorbid medical disorders: a pilot study." <i>Psychopharmacol Bull</i> 40: 47-56.	4
Wassertheil-Smoller, S., W. B. Applegate, K. Berge, C. J. Chang, B. R. Davis, R. Grimm, Jr., J. Kostis, S. Pressel and E. Schron (1996). "Change in depression as a precursor of cardiovascular events. SHEP Cooperative Research Group (Systolic Hypertension in the elderly)." <i>Arch Intern Med</i> 156(5): 553-561.	3
Weber, E., J. Stack, B. G. Pollock, B. Mulsant, A. Begley, S. Mazumdar and C. F. r. Reynolds (2000). "Weight change in older depressed patients during acute pharmacotherapy with paroxetine and nortriptyline: a double-blind randomized trial." <i>Am J Geriatr Psychiatry</i> 8: 245-250.	3
Wehmeier, P. M., M. Kluge, A. Maras, D. Riemann, M. Berger, R. Kohnen, R. W. Dittmann and W. F. Gattaz (2005). "Fluoxetine versus trimipramine in the treatment of depression in geriatric patients." <i>Pharmacopsychiatry</i> 38: 13-16.	3

Weihs, K. L., E. C. Settle, Jr., S. R. Batey, T. L. Houser, R. M. Donahue and J. A. Ascher (2000). "Bupropion sustained release versus paroxetine for the treatment of depression in the elderly." <i>J Clin Psychiatry</i> 61(3): 196-202.	3
Weintraub, D., J. E. Streim, C. J. Datto, I. R. Katz, S. D. DiFilippo and D. W. Oslin (2003). "Effect of Increasing the Dose and Duration of Sertraline Trial in the Treatment of Depressed Nursing Home Residents." <i>J Geriatr Psychiatry Neurol</i> 16: 109-111.	7
Weintraub, D., P. B. Rosenberg, L. T. Drye, B. K. Martin, C. Frangakis, J. E. Mintzer, A. P. Porsteinsson, L. S. Schneider, P. V. Rabins, C. A. Munro, C. L. Meinert, C. G. Lyketsos and R. Group Diads (2010). "Sertraline for the treatment of depression in Alzheimer disease: week-24 outcomes." <i>Am J Geriatr Psychiatry</i> 18: 332-340.	3
Weissman, J., A. Flint, B. Meyers, S. Ghosh, B. Mulsant, A. Rothschild and E. Whyte (2012). "Factors associated with non-completion in a double-blind randomized controlled trial of olanzapine plus sertraline versus olanzapine plus placebo for psychotic depression." <i>Psychiatry Res</i> 197: 221-226.	3
Weissman, M. M., B. Prusoff, A. J. Sholomskas and S. Greenwald (1992). "A double-blind clinical trial of alprazolam, imipramine, or placebo in the depressed elderly." <i>J Clin Psychopharmacol</i> 12(3): 175-182.	9
Ventura, D., E. P. Armstrong, G. H. Skrepnek and M. H. Erder (2007). "Escitalopram versus sertraline in the treatment of major depressive disorder: A randomized clinical trial." <i>Curr Med Res Opin</i> 23: 245-250.	7
Vermeiden, M., P. G. Mulder, W. W. van den Broek, J. A. Bruijn and T. K. Birkenhager (2013). "A double-blind randomized study comparing plasma level-targeted dose imipramine and high-dose venlafaxine in depressed inpatients." <i>J Psychiatr Res</i> 47(10): 1337-1342.	3
Wermuth, L., P. S. Sørensen, S. Timm, B. Christensen, N. P. Utzon, J. Boas, E. Dupont, E. Hansen, I. Magnussen, B. Mikkelsen, J. Worm-Petersen, L. Lauritzen, L. Bayer and P. Bech (1998). "Depression in idiopathic Parkinson's disease treated with citalopram: A placebo-controlled trial." <i>Nord J Psychiatry</i> 52(2): 163-169.	3
Versiani, M. (2000). "Reboxetine, the first selective noradrenaline reuptake inhibitor antidepressant: Efficacy and tolerability in 2613 patients." <i>Int J Psychiatry Clin Pract</i> 4: 201-208.	3
Versiani, M., R. Moreno, C. J. Ramakers-van Moorsel, A. J. Schutte and G. Comparative Efficacy Antidepressants Study (2005). "Comparison of the effects of mirtazapine and fluoxetine in severely depressed patients." <i>CNS Drugs</i> 19(2): 137-146.	3
Versiani, M., U. Oggero, P. Alterwain, R. Capponi, F. Dajas, G. Heinze-Martin, C. A. Marquez, M. A. Polo, L. E. Rivero-Almanzor, L. Rossel and et al. (1989). "A double-blind comparative trial of moclobemide v. imipramine and placebo in major depressive episodes." <i>Br J Psychiatry Suppl</i> (6): 72-77.	3
Whitehead, A. M. and J. J. Ashford (1992). "Fluvoxamine in the treatment of depressive illness. A series of double-blind hospital based comparative studies carried out in the UK." <i>Br J Clin Pract</i> 46: 21-23.	3

Whitmyer, V. G., D. L. Dunner, S. G. Kornstein, A. L. Meyers, C. H. Mallinckrodt, M. M. Wohlrreich, J. S. Gonzales and J. H. Greist (2007). "A comparison of initial duloxetine dosing strategies in patients with major depressive disorder." <i>J Clin Psychiatry</i> 68(12): 1921-1930.	3
Whyte, E. M., J. Basinski, P. Farhi, M. A. Dew, A. Begley, B. H. Mulsant and C. F. Reynolds (2004). "Geriatric depression treatment in nonresponders to selective serotonin reuptake inhibitors." <i>Journal of Clinical Psychiatry</i> : 1634-1641.	4
Wiart, L., H. Petit, P. A. Joseph, J. M. Mazaux and M. Barat (2000). "Fluoxetine in early poststroke depression: a double-blind placebo-controlled study." <i>Stroke</i> 31: 1829-1832.	3
Vieta, E., M. Bauer, S. Montgomery, R. S. McIntyre, J. Szamosi, W. R. Earley and H. Eriksson (2013). "Pooled analysis of sustained response rates for extended release quetiapine fumarate as monotherapy or adjunct to antidepressant therapy in patients with major depressive disorder." <i>J Affect Disord</i> 150(2): 639-643.	3
Wiethoff, K., M. Bauer, T. C. Baghai, H. J. Moller, R. Fisher, D. Hollinde, J. Kiermeir, I. Hauth, G. Laux, J. Cordes, P. Brieger, K. T. Kronmuller, J. Zeiler and M. Adli (2010). "Prevalence and treatment outcome in anxious versus nonanxious depression: results from the German Algorithm Project." <i>J Clin Psychiatry</i> 71: 1047-1054.	3
Viktrup, L. and I. Yalcin (2007). "Duloxetine treatment of stress urinary incontinence in women: effects of demographics, obesity, chronic lung disease, hypoestrogenism, diabetes mellitus, and depression on efficacy." <i>Eur J Obstet Gynecol Reprod Biol</i> 133(1): 105-113.	3
Wiles, N. J., J. Mulligan, T. J. Peters, P. J. Cowen, V. Mason, D. Nutt, D. Sharp, D. Tallon, L. Thomas, M. C. O'Donovan and G. Lewis (2012). "Severity of depression and response to antidepressants: GENPOD randomised controlled trial." <i>Br J Psychiatry</i> 200: 130-136.	4
Willem's, J. L., R. J. Willem's, G. M. Willem's, A. E. R. Arnold, W. Van de and M. Verstraete (1990). "Significance of initial ST segment elevation and depression for the management of thrombolytic therapy in acute myocardial infarction." <i>Circulation</i> 82: 1147-1158.	4
Williams, J. W., Jr, J. Barrett, T. Oxman, E. Frank, W. Katon, M. Sullivan, J. Cornell and A. Sengupta (2000). "Treatment of dysthymia and minor depression in primary care: A randomized controlled trial in older adults." <i>JAMA</i> 284: 1519-1526.	4
Williams, J. W., Jr, W. Katon, E. H. B. Lin, P. H. Nöel, J. Worchel, J. Cornell, L. Harpole, B. A. Fultz, E. Hunkeler, V. S. Mika and J. Unützer (2004). "Improving patient care. The effectiveness of depression care management on diabetes-related outcomes in older patients." <i>Ann Intern Med</i> 140: 1015.	4
Williams, L. S., K. Kroenke, T. Bakas, L. D. Plue, E. Brizendine, W. Tu and H. Hendrie (2007). "Care management of poststroke depression: a randomized, controlled trial." <i>Stroke</i> 38(3): 998-1003.	4
Williams, M. M., R. E. Clouse, B. D. Nix, E. H. Rubin, G. S. Sayuk, J. B. McGill, A. J. Gelenberg, P. S. Ciechanowski, I. B. Hirsch and P. J. Lustman (2007). "Efficacy of sertraline in prevention of depression recurrence in older versus younger adults with diabetes." <i>Diabetes Care</i> 30: 801-806.	4
Volkers, A. C., J. H. Tulen, B. van den, W. W, J. A. Bruyn, J. Passchier and L. Pepplinkhuizen (2004). "Effects of imipramine, fluvoxamine and depressive mood on autonomic cardiac functioning in major depressive disorder." <i>Pharmacopsychiatry</i> 37: 18-25.	7

Volz, H. P., H. Muller and H. J. Moller (1995). "Are there any differences in the safety and efficacy of brofaromine and imipramine between non-elderly and elderly patients with major depression?" <i>Neuropsychobiology</i> 32(1): 23-30.	9
Volz, H. P., H. Muller, Y. Sturm, B. Preussler and H. J. Moller (1995). "Effect of initial treatment with antidepressants as a predictor of outcome after 8 weeks." <i>Psychiatry Res</i> 58: 107-115.	3
Volz, H.P., Treatment of cognitive deficits in MDD, <i>International Journal of Psychiatry in Clinical Practice</i> , 17 (2013) 7-8. Accession:	9
Von Knorring, L. (1980). "A double-blind trial: Vivalan against placebo in depressed elderly patients." <i>Journal of International Medical Research</i> 8(1): 18-21.	4
Von, K., W. Katon, T. Bush, E. H. Lin, G. E. Simon, K. Saunders, E. Ludman, E. Walker and J. Unutzer (1998). "Treatment costs, cost offset, and cost-effectiveness of collaborative management of depression." <i>Psychosom Med</i> 60: 143-149.	9
Vorbach, E. U., K. H. Arnoldt and W. D. Hubner (1997). "Efficacy and tolerability of St. John's wort extract LI 160 versus imipramine in patients with severe depressive episodes according to ICD-10." <i>Pharmacopsychiatry</i> 30 Suppl 2: 81-85.	9
Vorbach, E. U., W. D. Hubner and K. H. Arnold (1993). "Effectiveness and tolerance of the hypericum extract LI 160 in comparison with imipramine. Randomized double blind study with 135 out-patients." <i>Nervenheilkunde</i> 12: 290-296.	3
Wu, E., P. E. Greenberg, E. Yang, A. Yu and M. H. Erder (2008) "Comparison of escitalopram versus citalopram for the treatment of major depressive disorder in a geriatric population." <i>Curr Med Res Opin</i> , 2587-2595.	3
Xie, R., J. Liu and H. Quan (2005). "A prospective random clinical contrast study of treatment with sertraline in elderly patients with post-stroke depression." <i>Chinese Journal of Clinical Neuroscience</i> : 294-297.	11
Xiong, G. L., M. Fiuzat, M. Kuchibhatla, R. Krishnan, C. M. O'Connor, W. Jiang and S.-C. Investigators (2012). "Health status and depression remission in patients with chronic heart failure: patient-reported outcomes from the SADHART-CHF trial." <i>Circ Heart Fail</i> 5: 688-692.	4
Xu, S. Q., J. Cao and W. W. Huang (2010). "Comparison of escitalopram with fluoxetine in old depressive patients." <i>Chinese Journal of New Drugs</i> 19(3): 207-209.	11
Yalamanchili, V. and J. C. Gallagher (2012). "Treatment with hormone therapy and calcitriol did not affect depression in older postmenopausal women: no interaction with estrogen and vitamin D receptor genotype polymorphisms." <i>Menopause</i> 19: 697-703.	3
Yang, H., L. Sinicropi-Yao, S. Chuizi, S. J. Youn, A. Clain, L. Baer, Y. Chen, P. J. McGrath, M. Fava and G. I. Papakostas (2010). "Residual sleep disturbance and risk of relapse during the continuation/maintenance phase treatment of major depressive disorder with the selective serotonin reuptake inhibitor fluoxetine." <i>Annals of General Psychiatry</i> 9.	3
Yang, J., L. Zhao and X. Mai (2005) "A comparative study of Taoist cognitive psychotherapy from China and mianserin in the treatment of depression in late life." <i>Chinese Journal of Nervous and Mental Diseases</i> , 333-335.	11

Yang, Q. H., L. C. Shen, L. R. Jing, A. L. Zhu and X. W. Yu (1995) "A double-blind controlled trial of maprotryline and amitriptyline in 52 cases elderly depression." <i>Sichuan Mental Health</i> , 97-100.	9
Yargic, L. I., A. Corapcioglu, N. Kocabasoglu, A. Erdogan, G. Koroglu and D. Yilmaz (2004). "A prospective randomized single-blind, multicenter trial comparing the efficacy and safety of paroxetine with and without quetiapine therapy in depression associated with anxiety." <i>Int J Psychiatry Clin Pract</i> 8(4): 205-211.	3
Ye, L. X., Y. D. Wang, H. Wang and D. S. Liang (2006). "Effect of Paxil and berhomine on poststroke anxiety-depression and neurological recovery." <i>Chinese Journal of Clinical Rehabilitation</i> 10: 153-155.	3
Yohannes, A. M., M. J. Connolly and R. C. Baldwin (2001). "A feasibility study of antidepressant drug therapy in depressed elderly patients with chronic obstructive pulmonary disease." <i>Int J Geriatr Psychiatry</i> 16: 451-454.	9
Yuen, G. S., F. Gunning-Dixon, E. Woods, M. Hoptman and G. S. Alexopoulos (2012). "Neuroanatomical correlates of apathy in late life depression and antidepressant treatment response." <i>Neuropsychopharmacology</i> 38: S96.	9
Zajecka, J., A. Schatzberg, S. Stahl, A. Shah, A. Caputo and A. Post (2010). "Efficacy and safety of agomelatine in the treatment of major depressive disorder: a multicenter, randomized, double-blind, placebo-controlled trial." <i>J Clin Psychopharmacol</i> 30: 135-144.	9
Zerjav, S., G. Tse and M. J. W. Scott (2009). "Review of duloxetine and venlafaxine in depression." <i>Canadian Pharmacists Journal</i> 142: 144-152.	9
Zhang, G. and X. Zhou (2000) "A controlled study of mianserin in the treatment of depression in elder." <i>Journal of Clinical Psychological Medicine</i> , 268-269.	11
Zhang, G. Y. and Y. H. Pan (2007). "The efficacy of mirtazapine for acute poststroke depression and its effect on post stroke rehabilitation." <i>Chinese Journal of Cerebrovascular Diseases</i> 4: 164-167.	11
Zheng, A. L., W. H. Qi, D. Y. Hu, N. S. Cai, J. B. Ge, W. H. Fan, Y. F. Ni, G. P. Lu, F. R. Zhang, M. Wei, B. He, S. Y. Wu, B. G. Sun, Z. G. Wu, H. G. Jin and Y. Huang (2006). "[Effects of antidepressant therapy in patients with suspected "angina pectoris" and negative coronary angiogram complicating comorbid depression]." <i>Zhonghua Xin Xue Guan Bing Za Zhi</i> 34: 1097-1100.	11
Zhou, M. and L. Yao (2005). "The efficacy and tolerability of venlafaxine and fluoxetine in treatment of elderly patients with first-episode depression." <i>Chinese Journal of Psychiatry</i> : 157-160.	11
Zhu, J., X. Jiang, D. Zhou and F. Zhang (2005) "Comparative study of mirtazapine vs. fluoxetine in treatment of elderly depressive patients." <i>Journal of Clinical Psychological Medicine</i> , 277-278.	3
Zisook, S., K. Ganadjian, C. Moutier, R. Prather and S. Rao (2008). "Sequenced Treatment Alternatives to Relieve Depression (STARD): Lessons learned." <i>Journal of Clinical Psychiatry</i> 69: 1184-1185.	3
Zohar, J., H. Keegstra and L. Barrelet (2003). "Fluvoxamine as effective as clomipramine against symptoms of severe depression: Results from a multicentre, double-blind study." <i>Hum Psychopharmacol</i> 18: 113-119.	3

Zourkova, A. (2001). "Effect of mirtazapine and paroxetine on residual symptoms of depressive disorders and their effect on P450 CYP 2D6 activity." Homeostasis in Health and Disease 41: 242-249.

3

Psykologisk behandling

Referens	Exklusions- orsak
2004-Home-based programme significantly reduces depressive symptoms and improves health status in chronically ill older adults with minor depression or dysthymia.	9
2005-Cognitive therapy for depression. Bandolier 2005;12:1-2.	3
2010-Cognitive behavioral therapy effective for depressed elderly. J Natl Med Assoc 2010;102:358.	9
Afonso-2010-[Reminiscence with different types of autobiographical memories: Effects on the reduction of depressive symptomatology in old age].	11
Afonso R, Bueno B. [Reminiscence with different types of autobiographical memories: Effects on the reduction of depressive symptomatology in old age]. Psicothema 2010;22:213-20.	7
Afonso Rosa-2011-Reminiscence, Psychological Well-Being, and Ego Integrity in Portuguese Elderly People.	6
Al-Otaiba-2012-Age-based differences in treatment outcome among alcohol-dependent women.	3
Anderson-Hanley-2003-The effects of a control-enhancing intervention for nursing home residents: cognition and locus of control as moderators.	3
Ando M, Morita T, Akechi T, Okamoto T, Japanese Task Force for S, Care. Efficacy of short-term life-review interviews on the spiritual well-being of terminally ill cancer patients. J Pain Symptom Manage 2010;39:993-1002.	3
Apostolo JL, Cardoso DF, Rosa AI, Paul C. The effect of cognitive stimulation on nursing home elders:	3
Appels A, van E, Bar F, van der P, Erdman RA, Assman M, et al. Effects of a behavioural intervention on quality of life and related variables in angioplasty patients: results of the EXhaustion Intervention Trial. J Psychosom Res 2006;61:1-7; discussion 9.	3
Arean-1993-Comparative effectiveness of social problem-solving therapy and reminiscence therapy as treatments for depression in older adults.	3
Arean-2010-Problem-solving therapy and supportive therapy in older adults with major depression and executive dysfunction.	3
Axford J, Heron C, Ross F, Victor CR. Management of knee osteoarthritis in primary care: pain and depression are the major obstacles. J Psychosom Res 2008;64:461-7.	3
Baker-2010-Randomized controlled trial of cognitive-behavioural therapy for coexisting depression and alcohol problems: short-term outcome.	3
Bakker-2011-Integrative psychotherapeutic nursing home program to reduce multiple psychiatric symptoms of cognitively impaired symptoms of cognitively impaired patients and caregiver burden: Randomized controlled trial.	3
Beary-2013-Living with the 'black dog': depression in care homes.	9

Beckie TM, Beckstead JW, Schocken DD, Evans ME, Fletcher GF. The effects of a tailored cardiac rehabilitation program on depressive symptoms in women: A randomized clinical trial. <i>Int J Nurs Stud</i> 2011;48:3-12.	3
Bedi-2000-Assessing effectiveness of treatment of depression in primary care. Partially randomised preference trial.	3
Berger-2010-Patterns of circadian activity rhythms and their relationships with fatigue and anxiety/depression in women treated with breast cancer adjuvant chemotherapy.	6
Berman RL, Iris MA, Bode R, Drengenberg C. The effectiveness of an online mind-body intervention for older adults with chronic pain. <i>J Pain</i> 2009;10:68-79.	3
Bichescu D, Neuner F, Schauer M, Elbert T. Narrative exposure therapy for political imprisonment-related chronic posttraumatic stress disorder and depression. <i>Behav Res Ther</i> 2007;45:2212-20.	3
Bilotta C, Bergamaschini L, Spreafico S, Vergani C. Day care centre attendance and quality of life in depressed older adults living in the community. <i>European Journal of Ageing</i> 2010;7:29-35.	7
Black-1998-Effect of intervention for psychological distress on rehospitalization rates in cardiac rehabilitation patients.	3
Bohlmeijer-2009-The effects of integrative reminiscence on depressive symptomatology and mastery of older adults.	7
Bohlmeijer-2008-The effects of integrative reminiscence on meaning in life: results of a quasi-experimental study.	7
Bolton-2003-Group Interpersonal Psychotherapy for Depression in Rural Uganda: A Randomized Controlled Trial.	3
Bond-2010-The effects of a web-based intervention on psychosocial well-being among adults aged 60 and older with diabetes: a randomized trial.	3
Bosma-2011-Disparities by education level in outcomes of a self-management intervention: the DELTA trial in The Netherlands.	7
Bosmans-2012-Cost-effectiveness of problem-solving treatment in comparison with usual care for primary care patients with mental health problems: a randomized trial.	3
Bosmans-2007-Cost-effectiveness of interpersonal psychotherapy for elderly primary care patients with major depression.	3
Breitbart-2012-Pilot randomized controlled trial of individual meaning-centered psychotherapy for patients with advanced cancer.	3
Brodaty H, Draper BM, Millar J, Low LF, Lie D, Sharah S, et al. Randomized controlled trial of different models of care for nursing home residents with dementia complicated by depression or psychosis. <i>J Clin Psychiatry</i> 2003;64:63-72.	3
Broderick JE, Junghaenel DU, Schneider S, Bruckenthal P, Keefe FJ. Treatment expectation for pain coping skills training: relationship to osteoarthritis patients' baseline psychosocial characteristics. <i>Clin J Pain</i> 2011;27:315-22.	3
Brody-2006-Age-related macular degeneration: self-management and reduction of depressive symptoms in a randomized, controlled study.	3

Buckwalter KC, Gerdner L, Kohout F, Hall GR, Kelly A, Richards B, et al. A nursing intervention to decrease depression in family caregivers of persons with dementia. <i>Arch Psychiatr Nurs</i> 1999;13:80-8.	3
Buist-Bouwman-2005-Collaborative care management improves physical functioning in older people with depression.	9
Burns A, Banerjee S, Morris J, Woodward Y, Baldwin R, Proctor R, et al. Treatment and prevention of depression after surgery for hip fracture in older people: randomized, controlled trials. <i>J Am Geriatr Soc</i> 2007;55:75-80.	3
Burns DS. The effect of the bonny method of guided imagery and music on the mood and life quality of cancer patients. <i>J Music Ther</i> 2001;38:51-65.	3
Carreira-2008-A controlled evaluation of monthly maintenance interpersonal psychotherapy in late-life depression with varying levels of cognitive function.	9
Cartwright-2013-Effect of telehealth on quality of life and psychological outcomes over 12 months (Whole Systems Demonstrator telehealth questionnaire study): nested study of patient reported outcomes in a pragmatic, cluster randomised controlled trial.	4
Catalan J, Gath DH, Anastasiades P, Bond SA, Day A, Hall L. Evaluation of a brief psychological treatment for emotional disorders in primary care. <i>Psychol Med</i> 1991;21:1013-8.	3
Cavanagh-2011-The implementation of computerized cognitive behavioural therapies in a service user-led, third sector self help clinic.	3
Cavanagh-2006-The effectiveness of computerized cognitive behavioural therapy in routine care.	3
Cernin PA, Lichtenberg PA. Behavioral treatment for depressed mood: a pleasant events intervention for seniors residing in assisted living. <i>Clinical Gerontologist</i> 2009;32:324-331.	3
Chan-2010-Effects of music on depression and sleep quality in elderly people: A randomised controlled trial.	3
Chan-2009-Effect of music on depression levels and physiological responses in community-based older adults.	3
Chan, M. F., K. S. Leong, B. L. Heng, B. K. Mathew, S. B. Khan, S. S. Lourdusamy, M. Nagapan, S. F. Woo, W. Y. Chee, R. C. Ho and B. J. Taylor (2014). "Reducing depression among community-dwelling older adults using life-story review: a pilot study." <i>Geriatr Nurs</i> 35(2): 105-110.	3
Chan-2013-A randomised controlled study to explore the effect of life story review on depression in older Chinese in Singapore.	3
Chan MF, Tien A, Khan SBAL, Ho RCM, Thayala J. A Randomised Controlled Study to Explore the Effect of Life-story Review o	3
Chan-2012-Effects of music on depression in older people: a randomised controlled trial.	3
Chang BL. Cognitive-behavioral intervention for homebound caregivers of persons with dementia. <i>Nurs Res</i> 1999;48:173-82.	3
Chew-Graham-2007-A randomised controlled trial to test the feasibility of a collaborative care model for the management of depression in older people.	3

Chiang-2010-The effects of reminiscence therapy on psychological well-being, depression, and loneliness among the institutionalized aged.	3
Chippendale-2012- Effect of life review writing on depressive symptoms in older adults: a randomized controlled trial.	3
Choi I, Zou J, Titov N, Dear BF, Li S, Johnston L, et al. Culturally attuned internet treatment for depression amongst Chinese Australians: A randomised controlled trial. <i>J Affect Disord</i> 2012;136:459-468.	3
Choi NG, Hegel MT, Sirrianni L, Marinucci ML, Bruce ML. Passive coping response to depressive symptoms among low-income homebound older adults: does it affect depression severity and treatment outcome? <i>Behav Res Ther</i> 2012;50:668-74.	7
Choi NG, Marti CN, Bruce ML, Hegel MT. Depression in homebound older adults: problem-solving therapy and personal and social resourcefulness.	3
Choi NG, Marti CN, Bruce ML, Hegel MT, Wilson NL, Kunik ME. SIX-MONTH POSTINTERVENTION DEPRESSION AND DISABILITY OUTCOMES OF IN-HOME	3
Chou KR, Lin Y. The effectiveness of group music therapy to improve depression and cognition status in elderly persons with dementia. <i>European Psychiatry</i> 2012;27.	9
Chu-2013-The Impact of Group Music Therapy on Depression and Cognition in Elderly Persons With Dementia: A Randomized Controlled Study.	3
Chung LJ, Tsai PS, Liu BY, Chou KR, Lin WH, Shyu YK, et al. Home-based deep breathing for depression in patients with coronary heart disease: a randomised controlled trial. <i>Int J Nurs Stud</i> 2010;47:1346-53.	3
Cole Martin G, Fenton Fred R, Engelsmann F, Mansouri I. Effectiveness of geriatric psychiatry consultation in an acute care hospital: A randomized clinical trial. <i>J Am Geriatr Soc</i> 1991;39:1183-1188.	4
Constantino-2001-Group intervention for widowed survivors of suicide.	3
Coon-2003-Anger and depression management: psychoeducational skill training interventions for women caregivers of a relative with dementia.	3
Dakin-2013-Patient perspectives on the benefits of psychotherapy for late-life depression.	7
Damush TM, Wu J, Bair MJ, Sutherland JM, Kroenke K. Self-management practices among primary care patients with musculoskeletal pain and depression. <i>J Behav Med</i> 2008;31:301-7.	3
Dao TK, Youssef NA, Armsworth M, Wear E, Papathopoulos KN, Gopaldas R. Randomized controlled trial of brief cognitive behavioral intervention for depression and anxiety symptoms preoperatively in patients undergoing coronary artery bypass graft surgery. <i>J Thorac Cardiovasc Surg</i> 2011;142:e109-15.	3
Davis MC. Life review therapy as an intervention to manage depression and enhance life satisfaction in individuals with right hemisphere cerebral vascular accidents; 1997.	3
Davison BJ, Degner LF. Empowerment of men newly diagnosed with prostate cancer. <i>Cancer Nurs</i> 1997;20:187-96.	3

de G, L E, Hollon SD, Huibers MJ. Predicting outcome in computerized cognitive behavioral therapy for depression in primary care: A randomized trial. <i>J Consult Clin Psychol</i> 2010;78:184-9.	3
de G, L E, Huibers MJ, Riper H, Gerhards SA, Arntz A. Use and acceptability of unsupported online computerized cognitive behavioral therapy for depression and associations with clinical outcome. <i>J Affect Disord</i> 2009;116:227-31.	3
de M, Saskia, Dekker J, Schoevers R, van A, Gerda, et al. Short psychodynamic supportive psychotherapy, antidepressants, and their combination in the treatment of major depression: A mega-analysis based on three randomized clinical trials. <i>Depress Anxiety</i> 2008;25:565-574.	9
DeBerry S. The effects of mediation-relaxation on anxiety and depression in a geriatric population. <i>Psychotherapy</i> 1982;19:512-521.	3
Dechamps-2010-Individualized Cognition-Action intervention to prevent behavioral disturbances and functional decline in institutionalized older adults: a randomized pilot trial.	3
Delaney C, Apostolidis B. Pilot testing of a multicomponent home care intervention for older adults with heart failure: an academic clinical partnership. <i>J Cardiovasc Nurs</i> 2010;E27-40.	7
Dobkin-2011-Cognitive-behavioral therapy for depression in Parkinson's disease: a randomized, controlled trial.	3
Dobkin RD, Menza M, Allen LA, Tiu J, Friedman J, Bienfait KL, et al. Telephone-based cognitive-behavioral therapy for depression in Parkinson disease. <i>J Geriatr Psychiatry Neurol</i> 2011;24:206-14.	3
Dobkin RD, Troster AI, Rubino JT, Allen LA, Gara MA, Mark MH, et al. Neuropsychological outcomes after psychosocial intervention for depression	3
Doering LV, Chen B, Bodán RC, Magsarili MC, Nyamathi A, Irwin MR. Early cognitive behavioral therapy for depression after cardiac surgery	3
Dolbeault-2009-The effectiveness of a psycho-educational group after early-stage breast cancer treatment: results of a randomized French study.	3
Dombrovski-2007-Maintenance treatment for old-age depression preserves health-related quality of life: a randomized, controlled trial of paroxetine and interpersonal psychotherapy.	3
Donkin L, Hickie IB, Christensen H, Naismith SL, Neal B, Cockayne NL, et al. Sampling bias in an internet treatment trial for depression. <i>Transl Psychiatry</i> 2012;2:e174.	7
Donnan P, Hutchinson A, Paxton R, Grant B, Firth M. Self-help materials for anxiety: a randomized controlled trial in general practice. <i>Br J Gen Pract</i> 1990;40:498-501.	3
Doorenbos-2006-Physical functioning: effect of behavioral intervention for symptoms among individuals with cancer.	3
Downe-Wamboldt BL, Butler LJ, Melanson PM, Coulter LA, Singleton JF, Keefe JM, et al. The effects and expense of augmenting usual cancer clinic care with telephone problem-solving counseling. <i>Cancer Nurs</i> 2007;30:441-53.	3
Edelman-1999-A group cognitive behaviour therapy programme with metastatic breast cancer patients.	3

Eller LS, Kirksey KM, Nicholas PK, Corless IB, Holzemer WL, Wantland DJ, et al. A randomized controlled trial of an HIV/AIDS Symptom Management Manual for depressive symptoms. AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV 2013;25:391-399.	3
Ernst S, Welke J, Heintze C, Gabriel R, Zollner A, Kiehne S, et al. Effects of mindfulness-based stress reduction on quality of life in nursing home residents: a feasibility study. Forsch Komplementmed 2008;15:74-81.	7
Ersek-2008-Results of a randomized controlled trial to examine the efficacy of a chronic pain self-management group for older adults [ISRCTN11899548].	3
Escobar-2007-Effectiveness of a time-limited cognitive behavior therapy type intervention among primary care patients with medically unexplained symptoms.	3
Fan XY. Effect of psychologic interference on gastric power of sufferers from functional indigestion and depression. Chinese Journal of Clinical Rehabilitation 2006;10:22-24.	11
Fernandez-Calvo-2011-[Efficacy of cognitive training programs based on new software technologies in patients with Alzheimer-type dementia].	11
Fledderus-2012-Acceptance and commitment therapy as guided self-help for psychological distress and positive mental health: A randomized controlled trial.	3
Flemming N, Kjeld A, Lolk A. Unipolar depression in elderly treated with home visits: Follow-up study. Neuroepidemiology 2012;39:225-226.	7
Floyd M, Rohen N, Shackelford JA, Hubbard KL, Parnell MB, Scogin F, et al. Two-year follow-up of bibliotherapy and individual cognitive therapy for depressed older adults. Behav Modif 2006;281-94.	3
Floyd-2004-Cognitive therapy for depression: A comparison of individual psychotherapy and bibliotherapy for depressed older adults.	3
Foley-2010-Mindfulness-based cognitive therapy for individuals whose lives have been affected by cancer: a randomized controlled trial.	3
Freedland-2012-Effect of obstructive sleep apnea on response to cognitive behavior therapy for depression after an acute myocardial infarction.	9
Freedland KE, Skala JA, Carney RM, Rubin EH, Lustman PJ, Davila-Roman VG, et al. Treatment of depression after coronary artery bypass surgery: a randomized controlled trial. Arch Gen Psychiatry 2009;66:387-96.	3
Fry PS. Structured and unstructured reminiscence training and depression among the elderly. Clinical Gerontologist 1983:15-37.	3
Furukawa TA. A stepped-care prevention programme decreased depression and anxiety disorders in elderly patients. Evidence Based Medicine 2009;14:146-146.	9
Furze-2009-"Prehabilitation" prior to CABG surgery improves physical functioning and depression.	3
Galfin-2012-A brief guided self-help intervention for psychological distress in palliative care patients: a randomised controlled trial.	3
Gallagher DE. Comparative effectiveness of group psychotherapies for reduction of depression in elderly outpatients. In: Dissertation Abstracts International; 1979. p 5550-1.	9

Gallagher DE, Thompson LW. Treatment of major depressive disorder in older adult outpatients with brief psychotherapies. <i>Psychotherapy (Chicago, Ill.)</i> 1982;482-490.	9
Gallagher DE, Thompson LW. Effectiveness of psychotherapy for both endogenous and nonendogenous depression in older adult outpatients. <i>Journals of Gerontology</i> 1983;707-712.	3
Gallagher-Thompson-1990-Maintenance of gains versus relapse following brief psychotherapy for depression.	3
Gallagher-Thompson-1994-Comparative effects of cognitive-behavioral and brief psychodynamic psychotherapies for depressed family caregivers.	3
Gallegos AM, Hoerger M, Talbot NL, Moynihan JA, Duberstein PR. Emotional benefits of mindfulness-based stress reduction in older adults	3
Gander F, Proyer RT, Ruch W, Wyss T. Strength-based positive interventions:	3
Gask-2004-A pragmatic cluster randomized controlled trial of an educational intervention for GPs in the assessment and management of depression.	4
Gatz M, Fiske A, Fox Lauren S, Kaskie B, Kasl-Godley Julia E, McCallum Todd J, et al. Empirically validated psychological treatments for older adults. <i>Journal of Mental Health and Aging</i> 1998;4:9-46.	9
Gaugler-2008-Can counseling and support reduce burden and depressive symptoms in caregivers of people with Alzheimer's disease during the transition to institutionalization? Results from the New York University caregiver intervention study.	3
Gellis-2008-Randomized controlled trial of problem-solving therapy for minor depression in home care.	10
Gensichen J, Petersen JJ, Karroum T, Rauck S, Ludman E, Konig J, et al. Positive impact of a family practice-based depression case management on patient's self-management. <i>Gen Hosp Psychiatry</i> 2011;33:23-8.	3
Geron SM, Keefe B. Moving evidence-based interventions to populations: a case study using social workers in primary care. <i>Home Health Care Serv Q</i> 2006;25:95-113.	9
Giaquinto-2006-Effects of music-based therapy on distress following knee arthroplasty.	3
Gilbody S. Depression in older adults: collaborative care model seems effective. <i>Evidence Based Mental Health</i> 2008;11:44-44.	9
Gitlin-2012-A community-integrated home based depression intervention for older African Americans: [corrected] description of the Beat the Blues randomized trial and intervention costs.	9
Glueckauf RL, Davis WS, Willis F, Sharma D, Gustafson DJ, Hayes J, et al. Telephone-based, cognitive-behavioral therapy for African American dementia caregivers with depression: initial findings. <i>Rehabil Psychol</i> 2012;57:124-39.	3
Goerling U, Foerg A, Sander S, Schramm N, Schlag PM. The impact of short-term psycho-oncological interventions on the psychological outcome of cancer patients of a surgical-oncology department - a randomised controlled study. <i>Eur J Cancer</i> 2011;47:2009-14.	3
Griffiths-2012-The Effectiveness of an Online Support Group for Members of the Community with Depression: A Randomised Controlled Trial.	3

Guan WB, Gao DJ, Li AM, Ouyang S, Dai ZC. Recent effects of early psychological intervention for patients with post-stroke depression. Chinese Journal of Clinical Rehabilitation 2004;8:6832-6833.	11
Guetin S, Ginies P, Siou DK, Picot MC, Pommie C, Guldner E, et al. The effects of music intervention in the management of chronic pain: a single-blind, randomized, controlled trial. Clin J Pain 2012;28:329-37.	3
Guetin-2009-Effect of music therapy on anxiety and depression in patients with Alzheimer's type dementia: randomised, controlled study.	3
Gum-2007-Low-income depressed older adults with psychiatric comorbidity: secondary analyses of response to psychotherapy and case management.	9
Haight BK. The therapeutic role of a structured life review process in homebound elderly subjects. J Gerontol 1988;P40-4.	3
Haight BK, Michel Y, Hendrix S. Life review: preventing despair in newly relocated nursing home residents short- and long-term effects. Int J Aging Hum Dev 1998;47:119-42.	3
Haley-2008-Long-term effects of bereavement and caregiver intervention on dementia caregiver depressive symptoms.	3
Hanser SB, Thompson LW. Effects of a music therapy strategy on depressed older adults. J Gerontol 1994;49:P265-9.	3
Hansson M, Bodlund O, Chotai J. Patient education and group counselling to improve the treatment of depression in primary care: a randomized controlled trial. J Affect Disord 2008;105:235-40.	9
Haringsma R, Engels GI, Cuijpers P, Spinhoven P. Effectiveness of the Coping With Depression (CWD) course for older adults provided by the community-based mental health care system in the Netherlands: a randomized controlled field trial. Int Psychogeriatr 2006;18:307-25.	3
Haringsma-2006-Predictors of response to the 'Coping with Depression' course for older adults. A field study.	7
Harrison M, Reeves D, Harkness E, Valderas J, Kennedy A, Rogers A, et al. A secondary analysis of the moderating effects of depression and multimorbidity on the effectiveness of a chronic disease self-management programme. Patient Educ Couns 2012;87:67-73.	7
Hautzinger M. Behavior therapy for depression in the elderly. In: Verhaltenstherapie; 1992. p 217-21.	11
Hautzinger M, Welz S. Cognitive behavioral therapy for depressed elderly outpatients - A controlled randomized trial. In: Verhaltenstherapie & Psychosoziale Praxis; 2004. p 789-98.	11
Hautzinger M, Welz S. [Cognitive behavioral therapy for depressed older outpatients--a controlled, randomized trial]. Z Gerontol Geriatr 2004;37:427-35.	11
Hautzinger M, Welz S. Kognitive Verhaltenstherapie bei Depressionen im Alter: Ergebnisse einer kontrollierten Vergleichsstudie unter ambulanten Bedingungen an Depressionen mittleren Schweregrads. Z Gerontol Geriatr 2004;37:427-435.	11

Hautzinger M, Welz S. Kurz- und längerfristige wirksamkeit psychologischer interventionen bei depressionen im alter. Zeitschrift für Klinische Psychologie und Psychotherapie: Forschung und Praxis 2008;37:52-60.	11
Hautzinger M, Welz S. Short- and long-term efficacy of psychological intervention for depression in older adults. [German]. [References]. Zeitschrift für Klinische Psychologie und Psychotherapie: Forschung und Praxis 2008;52-60.	11
Heckman-2011-A randomized clinical trial of a coping improvement group intervention for HIV-infected older adults.	3
Hirsch RD, Junglas K, Konradt B, Jonitz MF. [Humor therapy in the depressed elderly: results of an empirical study]. Zeitschrift für Gerontologie und Geriatrie : Organ der Deutschen Gesellschaft für Gerontologie und Geriatrie 2010;42-52.	11
Hoffman-2012-Effectiveness of mindfulness-based stress reduction in mood, breast- and endocrine-related quality of life, and well-being in stage 0 to III breast cancer: a randomized, controlled trial.	3
Holman-2011-Cost-effectiveness of cognitive behaviour therapy versus talking and usual care for depressed older people in primary care.	9
Hsiao-2012-The effects of psychotherapy on psychological well-being and diurnal cortisol patterns in breast cancer survivors.	3
Hsieh CJ, Chang C, Su SF, Hsiao YL, Shih YW, Han WH, et al. Reminiscence group therapy on depression and apathy in nursing home residents with mild-to-moderate dementia. Journal of Experimental and Clinical Medicine 2010;72-8.	3
Hsu CT, Weng CY, Kuo CS, Lin CL, Jong MC, Kuo SY, et al. Effects of a cognitive-behavioral group program for community-dwelling elderly with minor depression. Int J Geriatr Psychiatry 2010;654-5.	7
Hu Z, Hu Y, Lu Q. Impact of early rehabilitation therapy on post stroke depression. Chinese Journal of Clinical Rehabilitation 2003;7:849.	3
Hummel J, Weisbrod C, Oster P, Kopf D. The effect of cognitive behavioral therapy	9
Hussain RA, Lawrence PS. Social reinforcement of activity and problem-solving training in the treatment of depressed institutionalized elderly patients. Cognitive Therapy and Research 1981;5:57-69.	3
Hynnninen-2010-A randomized controlled trial of cognitive behavioral therapy for anxiety and depression in COPD.	3
Jarrett-2012-Cognitive reactivity, dysfunctional attitudes, and depressive relapse and recurrence in cognitive therapy responders.	3
Jarvik-1982-Treating geriatric depression: A 26-week interim analysis.	3
Jerant-2008-Perceived control moderated the self-efficacy-enhancing effects of a chronic illness self-management intervention.	3
Johns SA, Kroenke K, Theobald DE, Wu J, Tu W. Telecare management of pain and depression in patients with cancer: patient satisfaction and predictors of use. J Ambul Care Manage 2011;34:126-39.	3

Jones-2003-Reminiscence therapy for older women with depression. Effects of nursing intervention classification in assisted-living long-term care.	7
Jonkers-2012-The effectiveness of a minimal psychological intervention on self-management beliefs and behaviors in depressed chronically ill elderly persons: a randomized trial.	3
Jonkers-2009-Economic evaluation of a minimal psychological intervention in chronically ill elderly patients with minor or mild to moderate depression: a randomized trial (the DELTA-study).	9
Julião M, Oliveira F, Nunes B, Vaz Carneiro A, Barbosa A. Efficacy of Dignity Therapy on Depression and Anxiety in Portuguese Terminally Ill Patients:	3
Karimi H, Dolatshahee B, Momeni K, Khodabakhshi A, Rezaei M, Kamrani AA. Effectiveness of integrative and instrumental reminiscence therapies on depression symptoms reduction in institutionalized older adults: An empirical study. Aging Ment Health 2010;881-7.	3
Katon-2001-A randomized trial of relapse prevention of depression in primary care.	3
Kay-Lambkin-2011-Clinician-assisted computerised versus therapist-delivered treatment for depressive and addictive disorders: A randomised controlled trial.	3
Kessler D, Lewis G, Kaur S, Wiles N, King M, Weich S, et al. Therapist-delivered internet psychotherapy for depression in primary care: a randomised controlled trial. The Lancet 2009;374:628-634.	3
King-2002-Effectiveness of teaching general practitioners skills in brief cognitive behaviour therapy to treat patients with depression: randomised controlled trial.	3
Kitsumban V, Thapinta D, Sirindharo PB, Anders RL. Effect of cognitive mindfulness practice program on depression among elderly Thai women. Thai Journal of Nursing Research 2009;13:95-107.	3
Klausner-1998-Late-life depression and functional disability: the role of goal-focused group psychotherapy.	3
Klein DN, Leon AC, Li C, D'Zurilla TJ, Black SR, Vivian D, et al. Social problem solving and depressive symptoms over time: a randomized clinical trial of cognitive-behavioral analysis system of psychotherapy, brief supportive psychotherapy, and pharmacotherapy. J Consult Clin Psychol 2011;79:342-52.	3
Klemm P. Effects of online support group format (moderated vs peer-led) on depressive symptoms and extent of participation in women with breast cancer. Comput Inform Nurs 2012;30:9-18.	3
Knapp-2000-Non-drug strategies to resolve psycho-social difficulties after stroke.	9
Ko-2011-Effects of laughter therapy on depression, cognition and sleep among the community-dwelling elderly	3
Koertge-2008-Effects of a stress management program on vital exhaustion and depression in women with coronary heart disease: a randomized controlled intervention study.	3

Kohen-2011-Response to psychosocial treatment in poststroke depression is associated with serotonin transporter polymorphisms.	9
Kohn-2000-The effect of psychological intervention on patients' long-term adjustment to the ICD: a prospective study.	3
Kolanowski AM, Litaker M, Buettner L. Efficacy of theory-based activities for behavioral symptoms of dementia. <i>Nurs Res</i> 2005;54:219-28.	3
Kong Y, Dong WL, Liu CF. Fluoxetine for poststroke depression: A randomized placebo controlled clinical trial. <i>Neural Regeneration Research</i> 2007;2:162-165.	3
Konnert-2009-The prevention of depression in nursing home residents: a randomized clinical trial of cognitive-behavioral therapy.	3
Kootker-2012-The effectiveness of an augmented cognitive behavioural intervention for post-stroke depression with or without anxiety (PSDA): the Restore4Stroke-PSDA trial.	9
Korte J, Bohlmeijer ET, Cappeliez P, Smit F, Westerhof GJ. Life review therapy for older adults with moderate depressive symptomatology: a pragmatic randomized controlled trial. <i>Psychol Med</i> 2012;42:1163-73.	7
Korte, J., P. Cappeliez, T. Bohlmeijer Ernst and J. Westerhof Gerben (2012). "Meaning in life and mastery mediate the relationship of negative reminiscence with psychological distress among older adults with mild to moderate depressive symptoms." <i>European Journal of Ageing</i> 9: 343-351.	6
Kramer-2005-Evaluation and treatment of depression (part I): benefits for patients, providers, and payors.	9
Krishnamurthy-2007-Assessing depression following two ancient Indian interventions: effects of yoga and ayurveda on older adults in a residential home.	3
Kroenke-2010-Effect of telecare management on pain and depression in patients with cancer: a randomized trial.	3
Kuharek JM. Behavioral group therapy on depressed inpatient elderly: A comparative study. In: <i>Dissertation Abstracts International</i> ; 1991. p 265-6.	9
Kunik-2001-One session cognitive behavioural therapy for elderly patients with chronic obstructive pulmonary disease.	3
Kurtz-2005-A randomized, controlled trial of a patient/caregiver symptom control intervention: effects on depressive symptomatology of caregivers of cancer patients.	3
Lai Claudia KY, Chi I, Kayser-Jones J. A randomized controlled trial of a specific reminiscence approach to promote the well-being of nursing home residents with dementia. <i>International Psychogeriatrics</i> 2004;16:33-49.	3
Laidlaw K. A randomised controlled trial of cognitive behaviour therapy versus treatment ad usual in the treatment of mild to moderate late life depression. <i>National Research Register</i> 2002.	9
Lamers F, Jonkers CC, Bosma H, Chavannes NH, Knottnerus JA, van E, et al. Improving quality of life in depressed COPD patients: effectiveness of a minimal psychological intervention. <i>COPD</i> 2010;7:315-22.	3

Lamers-2006-Effectiveness and cost-effectiveness of a minimal psychological intervention to reduce non-severe depression in chronically ill elderly patients: the design of a randomised controlled trial [ISRCTN92331982]	9
Lamers-2010-A minimal psychological intervention in chronically ill elderly patients with depression: a randomized trial.	3
Lamers F, Jonkers CC, Bosma H, Knottnerus JA, van E, J T. Treating depression in diabetes patients: does a nurse-administered minimal psychological intervention affect diabetes-specific quality of life and glycaemic control? A randomized controlled trial. <i>J Adv Nurs</i> 2011;67:788-99.	3
Landreville P, Bissonnette L. Effects of cognitive bibliotherapy for depressed older adults with a disability. <i>Clinical Gerontologist</i> 1997;17:35-55.	3
Lang-2003-Brief intervention for co-occurring anxiety and depression in primary care: a pilot study.	3
Latour D, Cappeliez P. Pretherapy training for group cognitive therapy with depressed older adults. <i>Canadian Journal on Aging</i> 1994;13:221-235.	4
Lavretsky-2013-A pilot study of yogic meditation for family dementia caregivers with depressive symptoms: effects on mental health, cognition, and telomerase activity.	3
Lenze-2002-Combined pharmacotherapy and psychotherapy as maintenance treatment for late-life depression: effects on social adjustment.	3
Levesque DA, Van M, D F, Schneider RJ, Bauer MR, Goldberg DN, et al. Randomized trial of a computer-tailored intervention for patients with depression. <i>Am J Health Promot</i> 2011;26:77-89.	3
Little-2004-Randomised controlled trial of effect of leaflets to empower patients in consultations in primary care	3
Liu SK, Zhao XM, Xi ZM. [Incidence rate and acupuncture-moxibustion treatment of post-stroke depression]. <i>Zhongguo Zhen Jiu</i> 2006;26:472-4.	11
Liu X, Niu X, Feng Q, Liu Y. Effects of five-element music therapy on elderly people with seasonal affective disorder in a Chinese nursing home.	4
Lopez-2008-Analysis of the efficacy of a psychotherapeutic program to improve the emotional status of caregivers of elderly dependent relatives.	3
Lovejoy T. Telephone-delivered motivational interviewing targeting sexual risk behavior reduces depression, anxiety, and stress in HIV-positive older adults. <i>Annals of Behavioral Medicine</i> 2012;416-21.	3
Ludman Evette J, Peterson D, Katon Wayne J, Von K, Michael, Ciechanowski P, et al. Improving Confidence for Self Care in Patients with Depression and Chronic Illnesses. <i>Behavioral Medicine</i> 2013;39:1-6.	3
Lynch TR. Treatment of elderly depression with personality disorder comorbidity using dialectical behavior therapy. <i>Cognitive and Behavioral Practice</i> 2000:468-77.	9
Lynch-2003-Dialectical behavior therapy for depressed older adults: a randomized pilot study.	3

Lyness-2006-Outcomes of minor and subsyndromal depression among elderly patients in primary care settings.	9
Manber-2003-Patient's therapeutic skill acquisition and response to psychotherapy, alone or in combination with medication.	3
Mantovani-1996-Impact of psychosocial intervention on the quality of life of elderly cancer patients.	3
Markle-Reid-2006-Health promotion for frail older home care clients.	3
Marmar CR, Gaston L, Gallagher D, Thompson LW. Alliance and outcome in late-life depression. <i>J Nerv Ment Dis</i> 1989;177:464-72.	6
Martensson-2005-Patients with heart failure in primary health care: effects of a nurse-led intervention on health-related quality of life and depression.	3
Mastel-Smith B. An intervention to improve the health of people 60 years and older; 2004.	9
Mastel-Smith B, Binder B, Malecha A, Hersch G, Symes L, McFarlane J. Testing therapeutic life review offered by home care workers to decrease depression among home-dwelling older women. <i>Issues Ment Health Nurs</i> 2006;27:1037-1049.	7
Mastel-Smith-2007-Geropsychiatry. Improving depressive symptoms in community-dwelling older adults.	3
Mastel-Smith-2007-Improving depressive symptoms in community-dwelling older adults: a psychosocial intervention using life review and writing.	3
Matlock JA. The impact of supportive therapy with geriatric patients in the first thirty days of nursing home admission; 1995.	9
Maynard CK. Comparison of effectiveness of group interventions for depression in women. <i>Arch Psychiatr Nurs</i> 1993;7:277-83.	3
McCaffrey R. The effect of healing gardens and art therapy on older adults with mild to moderate depression. <i>Holistic Nursing Practice</i> 2007;21:79-84.	6
McCusker J, Cole M, Keller E, Bellavance F, Berard A, Jacoby R. Review: Heterocyclic antidepressants and rational psychological therapies reduce depression in older ambulatory patients. <i>Evidence-Based Medicine</i> 1998;3:152.	9
McGeoch GR, Willsman KJ, Dowson CA, Town GI, Frampton CM, McCartin FJ, et al. Self-management plans in the primary care of patients with chronic obstructive pulmonary disease. <i>Respirology</i> 2006;11:611-8.	3
McLaughlin-20085-Improving psychologic adjustment to chronic illness in cardiac patients. The role of depression and anxiety.	3
McMillan D, Gilbody S. A stepped care intervention is effective for elderly people with subthreshold depression or anxiety. <i>Evidence Based Mental Health</i> 2009;12:115-115.	9
McMurchie-2013-Computerised cognitive behavioural therapy for depression and anxiety with older people: A pilot study to examine patient acceptability and treatment outcome.	7

McPaul J. Focus on research... Home visit versus telephone follow-up in Phase II cardiac rehabilitation following myocardial infarction: effects on anxiety, depression, attendance at Phase III and visits to A&E or readmission. British Journal of Occupational Therapy 2008;71:320-320.	4
Meeks-2007-Activity participation and affect among nursing home residents: Support for a behavioural model of depression.	3
Merrill Kari A, Tolbert Valerie E, Wade Wendy A. Effectiveness of cognitive therapy for depression in a community mental health center: A benchmarking study. Journal of Consulting and Clinical Psychology 2003;71:404-409.	3
Miller MD, Frank E, Cornes C, Houck PR, Reynolds CF, rd. The value of maintenance interpersonal psychotherapy (IPT) in older adults with different IPT foci. Am J Geriatr Psychiatry 2003;11:97-102.	7
Mittelman-2004-Sustained benefit of supportive intervention for depressive symptoms in caregivers of patients with Alzheimer's disease	3
Mohammadi Ali Z, Shahabi T, Panah Fereshteh M. An evaluation of the effect of group music therapy on stress, anxiety, and depression levels in nursing home residents. Canadian Journal of Music Therapy 2011;17:55-68.	3
Mohr DC, Goodkin DE, Islar J, Hauser SL, Genain CP. Treatment of depression is associated with suppression of nonspecific and antigen-specific TH1 responses in multiple sclerosis. Archives of Neurology 2001;58:1081-1086.	3
Moleiro C, Beutler LE. Clinically significant change in psychotherapy for depressive disorders. J Affect Disord 2009;115:220-4.	3
Moon Fai C, P. Leong KS, Boon Ling H, Koottappal Mathew B, Banu A. L. Khan S, Sagayamary Lourdusamy S, et al. Reducing depression among community-dwelling older	3
Moorey-2009-A cluster randomized controlled trial of cognitive behaviour therapy for common mental disorders in patients with advanced cancer.	3
Morgan-2012-Email-based promotion of self-help for subthreshold depression: Mood Memos randomised controlled trial.	3
Morgan AJ, Mackinnon AJ, Jorm AF. Behavior change through automated e-mails: Mediation analysis of self-help strategy use for depressive symptoms. Behav Res Ther 2013;51:57-62.	7
Morgan MAJ, Coates MJ, Dunbar JA, Reddy P, Schlicht K, Fuller J. The TrueBlue model of collaborative care using practice nurses as case managers for depression alongside diabetes or heart disease: A randomised trial. BMJ Open 2013;3.	3
Mossey-1996-Effectiveness of a psychosocial intervention, interpersonal counseling, for subdysthymic depression in medically ill elderly.	3
Moynihan-1998-Evaluation of adjuvant psychological therapy in patients with testicular cancer: randomised controlled trial.	3
Mozley-2007-The care home activity project: does introducing an occupational therapy programme reduce depression in care homes?	3
Mundt-2007-Anchoring perceptions of clinical change on accurate recollection of the past: Memory enhanced retrospective evaluation of treatment (MERET®).	3

Murphy-2012-An evaluation of the effectiveness and cost effectiveness of the National Exercise Referral Scheme in Wales, UK: a randomised controlled trial of a public health policy initiative.	3
Nance2012-Pains, joys, and secrets: nurse-led group therapy for older adults with depression.	3
Ng-2013-The effect of life story review on depression of older community-dwelling Chinese adults in Singapore: A preliminary result.	3
Nomura-2009-[Individual reminiscence therapy improves self-esteem for Japanese community-dwelling older adults].	11
Norris CM, Patterson L, Galbraith D, Hegadoren KM. All you have to do is call; a pilot study to improve the outcomes of patients with coronary artery disease. <i>Applied Nursing Research</i> 2009;22:133-137.	3
Nygaard-2009-Effect of combination therapy with thyroxine (T4) and 3,5,3'-triiodothyronine versus T4 monotherapy in patients with hypothyroidism, a double-blind, randomised cross-over study.'	4
Okumiya-2005-Effects of group work programs on community-dwelling elderly people with age-associated cognitive decline and/or mild depressive moods: a Kahoku Longitudinal Aging Study.	3
O'Leary-2003-Cork Older Adult Intervention Project: outcomes of a gestalt therapy group with older adults.	3
Oranta-2010-The outcomes of interpersonal counselling on depressive symptoms and distress after myocardial infarction.	3
Oslin-2003-Disease management for depression and at-risk drinking via telephone in an older population of veterans.	3
Oxman-2001-Social support and treatment response in older depressed primary care patients	7
Oyama H, Ono Y, Watanabe N, Tanaka E, Kudoh S, Sakashita T, et al. Local community intervention through depression screening and group activity for elderly suicide prevention. <i>Psychiatry Clin Neurosci</i> 2006;110-4.	7
Pariser-2005-Effects of telephone intervention on arthritis self-efficacy, depression, pain, and fatigue in older adults with arthritis.	3
Pariser D, O'Hanlon AM, Speaker RS, Espinoza L. The effects of telephone intervention on arthritis self-efficacy, depression, pain and fatigue in older adults. <i>Journal of Geriatric Physical Therapy</i> 2003;26:45-45.	3
Patel-2010-Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial	3
Penckofer SM, Ferrans C, Mumby P, Byrn M, Emanuele MA, Harrison PR, et al. A psychoeducational intervention (SWEEP) for depressed women with diabetes. <i>Ann Behav Med</i> 2012;44:192-206.	3

Pickett YR, Kennedy GJ, Freeman K, Cummings J, Woolis W. The effect of telephone-facilitated depression care on older, medically ill patients. <i>T</i>	3
Popa-Velea-2010-Influence of personalized therapeutic approach on quality of life and psychiatric comorbidity in patients with advanced colonic cancer requiring palliative care.	3
Post EP, Miller MD, Schulberg HC. Using interpersonal therapy (IPT) to treat depression in older primary care patients. <i>Geriatrics</i> 2008;63:18.	9
Pot-2010-The impact of life review on depression in older adults: a randomized controlled trial.	3
Raglio A, Bellelli G, Traficante D, Gianotti M, Ubezio Maria C, Villani D, et al. Efficacy of music therapy in the treatment of behavioral and psychiatric symptoms of dementia. <i>Alzheimer Dis Assoc Disord</i> 2008;22:158-162.	3
Raglio A, Oasi O, Gianotti M, Manzoni V, Bolis S, Ubezio MC, et al. Effects of music therapy on psychological symptoms and heart rate variability in patients with dementia. A pilot study. <i>Curr Aging Sci</i> 2010;3:242-6.	3
Raue PJ, Morales KH, Post EP, Bogner HR, Have TT, Bruce ML. The wish to die and 5-year mortality in elderly primary care patients. <i>Am J Geriatr Psychiatry</i> 2010;18:341-50.	7
Rawl-2002-Intervention to improve psychological functioning for newly diagnosed patients with cancer.	3
Reynolds-1996-Treatment outcome in recurrent major depression: a post hoc comparison of elderly ("young old") and midlife patients.	3
Reynolds-1999-Nortriptyline and interpersonal psychotherapy as maintenance therapies for recurrent major depression: a randomized controlled trial in patients older than 59 years.	3
Reynolds-1999-Treatment of bereavement-related major depressive episodes in later life: a controlled study of acute and continuation treatment with nortriptyline and interpersonal psychotherapy.	3
Riegel B, Carlson B, Glaser D, Romero T. Randomized controlled trial of telephone case management in Hispanics of Mexican origin with heart failure. <i>J Card Fail</i> 2006;12:211-9.	3
Risos-Rio GER. The effects of a structured live music and arts program on depression, pain control and mood/behavior patterns among nursing home residents. <i>UPNAI Nursing Journal</i> 2009;5:9-15.	3
Rohricht F, Papadopoulos N, Priebe S. An exploratory randomized controlled trial of body psychotherapy for patients	3
Rokke-1999-The role of client choice and target selection in self-management therapy for depression in older adults.	3
Rokke PD, Tomhave JA, Jocic Z. Self-management therapy and educational group therapy for depressed elders. <i>Cognitive Therapy & Research</i> 2000;99-119.	3
Romeo R, Knapp M, Banerjee S, Morris J, Baldwin R, Tarrier N, et al. Treatment and prevention of depression after surgery for hip fracture in older people: cost-effectiveness analysis. <i>J Affect Disord</i> 2011;128:211-9.	6
Rosenthal MZ, Cheavens JS, Compton JS, Thorp SR, Lynch TR. Thought suppression and treatment outcome in late-life depression. <i>Aging and Mental Health</i> 2005;9:35-39.	7

Ross-2008-A randomized controlled trial of a close monitoring program for minor depression and distress.	3
Rucci P, Frank E, Scocco P, Calugi S, Miniati M, Fagiolini A, et al. Treatment-emergent suicidal ideation during 4 months of acute management of unipolar major depression with SSRI pharmacotherapy or interpersonal psychotherapy in a randomized clinical trial. <i>Depress Anxiety</i> 2011;28:303-309.	3
Safren SA, Gonzalez JS, Wexler DJ, Psaros C, Delahanty LM, Blashill AJ, et al. A randomized controlled trial of cognitive behavioral therapy	3
Salminen-2005-Effects of a health advocacy, counselling, and activation programme on depressive symptoms in older coronary heart disease patients.	3
Scazufca M, Matsuda CMCB. Review of the efficacy of psychotherapy vs. pharmacotherapy for depression treatment in old age. <i>Revista Brasileira de Psiquiatria</i> 2002;24:64-69.	9
Schramm E, Zobel I, Dykier P, Kech S, Brakemeier EL, Kulz A, et al. Cognitive behavioral analysis system of psychotherapy versus interpersonal psychotherapy for early-onset chronic depression: a randomized pilot study. <i>J Affect Disord</i> 2011;129:109-16.	3
Scogin F, Hamblin D, Beutler L. Bibliotherapy for depressed older adults: a self-help alternative. In: <i>Gerontologist</i> ; 1987. p 383-7.	3
Scogin F, Jamison C, Davis N. Two-year follow-up of bibliotherapy for depression in older adults. <i>Journal of Consulting and Clinical Psychology</i> 1990:665-7.	3
Scogin-1989-Comparative efficacy of cognitive and behavioral bibliotherapy for mildly and moderately depressed older adults	3
Scogin FR, Moss K, Harris GM, Presnell AH. Treatment of depressive symptoms in diverse, rural, and vulnerable older adults.	3
Scott-2000-Effects of cognitive therapy on psychological symptoms and social functioning in residual depression.	3
Sebregts EH, Falger PR, Appels A, Kester AD, Bar FW. Psychological effects of a short behavior modification program in patients with acute myocardial infarction or coronary artery bypass grafting. A randomized controlled trial. <i>J Psychosom Res</i> 2005;58:417-24.	3
Serfaty M, Csipke E, Haworth D, Murad S, King M. A talking control for use in evaluating the effectiveness of cognitive-behavioral therapy. <i>Behav Res Ther</i> 2011;49:433-40.	7
Serfaty MA, Deborah H, Buszewicz M, Blanchard M, Murad S, King M. The clinical effectiveness of individual cognitive behaviour therapy for depressed older people in primary care and the use of a talking control (TC). <i>European Psychiatry</i> 2011;26.	9
Serrano JP, Latorre JM, Montan~es J. [Life review therapy based on retrieval of specific autobiographical memories in elderly individuals with depression]. <i>Rev Esp Geriatr Gerontol</i> 2005;220-7.	11
Shellman-2009-The effects of integrative reminiscence on depressive symptoms in older African Americans.	3
Sherrill-1997-Psychoeducational workshops for elderly patients with recurrent major depression and their families.	4
Simon-2011-Randomized trial of depression follow-up care by online messaging.	3

Simpson GK, Tate RL, Whiting DL, Cotter RE. Suicide prevention after traumatic brain injury: a randomized controlled trial of a program for the psychological treatment of hopelessness. <i>J Head Trauma Rehabil</i> 2011;26:290-300.	3
Simson-2008-Psychotherapy intervention to reduce depressive symptoms in patients with diabetic foot syndrome.	3
Sloman R. Relaxation and imagery for anxiety and depression control in community patients with advanced cancer. <i>Cancer Nurs</i> 2002;25:432-5.	3
Snarski M, Scogin F, DiNapoli E, Presnell A, McAlpine J, Marcinak J. The effects of behavioral activation therapy with inpatient geriatric psychiatry patients. <i>Behav Ther</i> 2011;42:100-8.	3
Spek2008-One-year follow-up results of a randomized controlled clinical trial on internet-based cognitive behavioural therapy for subthreshold depression in people over 50 years.	3
Spek-2007-Internet-based cognitive behavioural therapy for subthreshold depression in people over 50 years old: a randomized controlled clinical trial.	3
Stasi MF, Amati D, Costa C, Resta D, Senepa G, Scarafioiti C, et al. Pet-therapy: a trial for institutionalized frail elderly patients. <i>Archives of gerontology and geriatrics. Supplement</i> 2004:407-12.	3
Steidtmann-2012-Patient treatment preference as a predictor of response and attrition in treatment for chronic depression.	3
Steuer-1984-Cognitive-behavioral and psychodynamic group psychotherapy in treatment of geriatric depression.	7
Stevens-Ratchford-1993-The effect of life review reminiscence activities on depression and self-esteem in older adults.	3
Stinson-2006-Structured reminiscence: an intervention to decrease depression and increase self-transcendence in older women	3
Stinson CK, Young EA, Kirk E, Walker R. Use of a structured reminiscence protocol to decrease depression in older women. <i>J Psychiatr Ment Health Nurs</i> 2010;17:665-73.	3
Strachowski D, Khaylis A, Conrad A, Neri E, Spiegel D, Taylor CB. The effects of cognitive behavior therapy on depression in older patients with cardiovascular risk. <i>Depress Anxiety</i> 2008:E1-10.	3
Strong-2008-Management of depression for people with cancer (SMaRT oncology 1): a randomised trial.	3
Su-2012-The prevalence of dementia and depression in Taiwanese institutionalized leprosy patients, and the effectiveness evaluation of reminiscence therapy--a longitudinal, single-blind, randomized control study.	3
Su XL, Xiao XC. Effect of psychotherapy on the motor functional rehabilitation in patients with post-stroke depression. <i>Chinese Journal of Clinical Rehabilitation</i> 2004;8:3720-3721.	11
Tang-2009-Personality change during depression treatment: a placebo-controlled trial.	3
Taveira-2011-Pharmacist-led group medical appointments for the management of type 2 diabetes with comorbid depression in older adults.	3

Taylor-Price C. The efficacy of structured reminiscence group psychotherapy as an intervention to decrease depression and increase psychological well-being in female nursing home residents. US; 1996.	3
Thielke-2007-Pain limits the effectiveness of collaborative care for depression.	9
Thomas-2006-Factors relating to depression after stroke.	7
Thompson-1988-Personality disorder and outcome in the treatment of late-life depression	3
Thompson-2001-Comparison of desipramine and cognitive/behavioral therapy in the treatment of elderly outpatients with mild-to-moderate depression.	3
Thompson LW, Gallagher D. Efficacy of psychotherapy in the treatment of late-life depression. Advances in Behaviour Research & Therapy 1984;127-39.	3
Thompson-1987-Comparative effectiveness of psychotherapies for depressed elders.	3
Tsai HH, Tsai YF. Changes in depressive symptoms, social support, and loneliness over 1 year after a minimum 3-month videoconference program for older nursing home resident. In: J Med Internet Res; 2011. p 373-84.	3
Tsai H-H, Tsai Y-F, Wang H-H, Chang Y-C, Chu Hao H. Videoconference program enhances social support, loneliness, and depressive status of elderly nursing home residents. Aging Ment Health 2010;14:947-954.	3
Turner-2013-Randomised controlled trial of group cognitive behaviour therapy versus brief intervention for depression in cardiac patients.	3
Turner A, Murphy BM, Higgins RO, Elliott PC, Le Grande MR, Goble AJ, et al. An integrated secondary prevention group programme reduces	3
Tyler P, Jones V, Thompson S, Catalan J, Schmidt U, Davidson K, et al. Service variation in baseline variables and prediction of risk in a randomised controlled trial of psychological treatment in repeated parasuicide: The Popmact study. International Journal of Social Psychiatry 2003;49:58-69.	3
Unutzer-2008-Long-term cost effects of collaborative care for late-life depression.	6
Van Der-2013-One-year effect of a nurse-led psychosocial intervention on depressive symptoms in patients with head and neck cancer: A randomized controlled trial.	3
van-2007-Interpersonal psychotherapy (IPT) for late-life depression in general practice: uptake and satisfaction by patients, therapists and physicians.	6
van-2006-Interpersonal psychotherapy for elderly patients in primary care.	3
van 't V-T, P J, Cuijpers P, Beekman AJ. [Prevention of depression and anxiety in older people]. Tijdschr Psychiatr 2011;53:579-84.	11
Wang-2005-The effects of reminiscence on depressive symptoms and mood status of older institutionalized adults in Taiwan.	3
Wang-2007-Group reminiscence therapy for cognitive and affective function of demented elderly in Taiwan.	3
Wang JJ, Hsu YC, Cheng SF. The effects of reminiscence in promoting mental health of Taiwanese elderly. Int J Nurs Stud 2005;42:31-6.	3

Wang-2011-Effects of gerotranscendence support group on gerotranscendence perspective, depression, and life satisfaction of institutionalized elders.	3
Wang-2003-Role of educational intervention in the management of comorbid depression and hypertension.	3
Warber SL, Ingerman S, Moura VL, Wunder J, Northrop A, Gillespie BW, et al. Healing the heart: a randomized pilot study of a spiritual retreat for depression in acute coronary syndrome patients. <i>Explore (NY)</i> 2011;7:222-33.	3
Waterreus-1994-Community psychiatric nurses for the elderly: well tolerated, few side-effects and effective in the treatment of depression.	4
Watt-2000-Integrative and instrumental reminiscence therapies for depression in older adults: intervention strategies and treatment effectiveness.	3
Weber-The impact of dyadic social support on self-efficacy and depression after radical prostatectomy.	3
Verkaik-2011-The effects of a nursing guideline on depression in psychogeriatric nursing home residents with dementia.	7
Werneck AL, Rosso AL, Vincent MB. The use of an antagonist 5-HT2a/c for depression and motor function in Parkinson' disease. <i>Arq Neuropsiquiatr</i> 2009;67:407-12.	3
Westerhof GJ, Bohlmeijer ET, van B, I M, Pot AM. Improvement in personal meaning mediates the effects of a life review intervention on depressive symptoms in a randomized controlled trial. <i>Gerontologist</i> 2010;50:541-9.	3
Wetherell-2009-Modular psychotherapy for anxiety in older primary care patients.	3
White-2002-A randomized controlled trial of the psychosocial impact of providing internet training and access to older adults.	3
Wilkinson-2009-A pilot randomised controlled trial of a brief cognitive behavioural group intervention to reduce recurrence rates in late life depression.	4
Willemse-2004-Minimal-contact psychotherapy for sub-threshold depression in primary care. Randomised trial.	3
Wolinsky-2009-The effect of speed-of-processing training on depressive symptoms in ACTIVE.	3
Worley MJ, Trim RS, Tate SR, Hall JE, Brown SA. Service utilization during and after outpatient treatment for comorbid substance use disorder and depression. <i>J Subst Abuse Treat</i> 2010;39:124-31.	3
Vos PJ, Visser AP, Garssen B, Duivenvoorden HJ, de H, H C. Effectiveness of group psychotherapy compared to social support groups in patients with primary, non-metastatic breast cancer. <i>J Psychosoc Oncol</i> 2007;25:37-60.	3
Wu HS, Xu Y, Xu YF, Zhu HX, Zhu LP, Xue JJ. Effectiveness of case management of chronic disease for elderly depression in community [Chinese]. <i>Chinese Mental Health Journal</i> 2010:245-9.	11

Wurtzen H, Dalton SO, Elsass P, Sumbundu AD, Steding-Jensen M, Karlsen RV, et al. Mindfulness significantly reduces self-reported levels of anxiety and depression: Results of a randomised controlled trial among 336 Danish women treated for stage I-III breast cancer. European Journal of Cancer 2013;49:1365-1373.	3
Wuthrich VM, Rapee RM. Randomised controlled trial of group cognitive behavioural therapy for comorbid anxiety	3
Young Laura-2010-Mindfulness-based stress reduction: Effect on emotional distress in older adults.	3
Youssef FA. The impact of group reminiscence counseling on a depressed elderly population. Nurse Pract 1990;15:32, 35-8.	7
Zauszniewski-2007-Effects of teaching resourcefulness and acceptance on affect, behavior, and cognition of chronically ill elders.	3
Zerhusen JD, Boyle K, Wilson W. Out of the darkness: group cognitive therapy for depressed elderly. Journal of psychosocial nursing and mental health services 1991;16:21.	3
Zhou-2011-Effects of music therapy on depression and duration of hospital stay of breast cancer patients after radical mastectomy.	3
Zhou W, He G, Gao J, Yuan Q, Feng H, Zhang CK. The effects of group reminiscence therapy on depression, self-esteem, and affect balance of Chinese community-dwelling elderly. Arch Gerontol Geriatr 2012;54:e440-7.	3

Fysisk aktivitet

Referens	Exklusions- orsak
Alentorn-Geli, E., J. Padilla, G. Moras, H. Lazaro and J. Fernandez-Sola (2008). "Six weeks of whole-body vibration exercise improves pain and fatigue in women with fibromyalgia." <i>J Altern Complement Med</i> 14: 975-981.	3
Antunes, H. K., S. G. Stella, R. F. Santos, O. F. Bueno and M. T. de Mello (2005). "Depression, anxiety and quality of life scores in seniors after an endurance exercise program." <i>Rev Bras Psiquiatr</i> 27(4): 266-271.	3
Berke, E. M., L. M. Gottlieb, A. V. Moudon and E. B. Larson (2007). "Protective association between neighborhood walkability and depression in older men." <i>J Am Geriatr Soc</i> 55: 526-533.	3
Blumenthal, J. A., C. F. Emery, D. J. Madden, S. Schniebold, M. Walsh-Riddle, L. K. George, D. C. McKee, M. B. Higginbotham, F. R. Cobb and R. E. Coleman (1991). "Long-term effects of exercise on psychological functioning in older men and women." <i>J Gerontol</i> 46: P352-361.	3
Blumenthal, J. A., M. A. Babyak, K. A. Moore, W. E. Craighead, S. Herman, P. Khatri, R. Waugh, M. A. Napolitano, L. M. Forman, M. Appelbaum, P. M. Doraiswamy and K. R. Krishnan (1999). "Effects of exercise training on older patients with major depression." <i>Arch Intern Med</i> 159: 2349-2356.	3
Bosnak-Guclu, M., H. Arikan, S. Savci, D. Inal-Ince, E. Tulumen, K. Aytemir and L. Tokgozoglu (2011). "Effects of inspiratory muscle training in patients with heart failure." <i>Respir Med</i> 105: 1671-1681.	3
Brittle, N., S. Patel, C. Wright, S. Baral, P. Versfeld and C. Sackley (2009). "An exploratory cluster randomized controlled trial of group exercise on mobility and depression in care home residents." <i>Clin Rehabil</i> 23: 146-154.	3
Brown, M. A., J. Goldstein-Shirley, J. Robinson and S. Casey (2001). "The effects of a multi-modal intervention trial of light, exercise, and vitamins on women's mood." <i>Women Health</i> 34: 93-112.	3
Cakar, E., U. Dincer, M. Z. Kiralp, D. B. Cakar, O. Durmus, H. Kilac, F. C. Soydan, S. Sevinc and C. Alper (2010). "Jumping combined exercise programs reduce fall risk and improve balance and life quality of elderly people who live in a long-term care facility." <i>Eur J Phys Rehabil Med</i> 46: 59-67.	3
Chaiyawat, P. and K. Kulkantrakorn (2012). "Randomized controlled trial of home rehabilitation for patients with ischemic stroke: impact upon disability and elderly depression." <i>Psychogeriatrics</i> 12: 193-199.	3
Chan, W., M. A. Immink and S. Hillier (2012). "Yoga and exercise for symptoms of depression and anxiety in people with poststroke disability: a randomized, controlled pilot trial." <i>Altern Ther Health Med</i> 18: 34-43.	3
Chen, K. M., M. H. Chen, M. H. Lin, J. T. Fan, H. S. Lin and C. H. Li (2010). "Effects of yoga on sleep quality and depression in elders in assisted living facilities." <i>J Nurs Res</i> : 53-61.	3

Chin, A. P., J. M. M, P. Van, N. M. M, J. W. R. Twisk and M. Van (2004). "Effects of resistance and all-round, functional training on quality of life, vitality and depression of older adults living in long-term care facilities: A 'randomized' controlled trial [ISRCTN87177281]." BMC Geriatr 4: 1-9.	3
Cho, K. L. (2008). "Effect of Tai Chi on depressive symptoms amongst Chinese older patients with major depression: the role of social support." Med Sport Sci 52: 146-154.	3
Chou, K. L., P. W. Lee, E. C. Yu, D. Macfarlane, Y. H. Cheng, S. S. Chan and I. Chi (2004). "Effect of Tai Chi on depressive symptoms amongst Chinese older patients with depressive disorders: a randomized clinical trial." Int J Geriatr Psychiatry 19: 1105-1107.	3
Clark, N. M., N. K. Janz, J. A. Dodge, M. A. Schork, T. E. Fingerlin, J. R. Wheeler, J. Liang, S. J. Keteyian and J. T. Santinga (2000). "Changes in functional health status of older women with heart disease: evaluation of a program based on self-regulation." J Gerontol B Psychol Sci Soc Sci 55(2): S117-126.	3
Conradsson, M., H. Littbrand, N. Lindelof, Y. Gustafson and E. Rosendahl (2010). "Effects of a high-intensity functional exercise programme on depressive symptoms and psychological well-being among older people living in residential care facilities: A cluster-randomized controlled trial." Aging Ment Health 14: 565-576.	3
Crotty, M., J. Prendergast, M. W. Battersby, D. Rowett, S. E. Graves, G. Leach and L. C. Giles (2009). "Self-management and peer support among people with arthritis on a hospital joint replacement waiting list: a randomised controlled trial." Osteoarthritis Cartilage 17: 1428-1433.	3
Deslandes, A. C., H. Moraes, H. Alves, F. A. M. S. Pompeu, H. Silveira and R. Mouta (2010) "Effect of aerobic training on EEG alpha asymmetry and depressive symptoms in the elderly: A 1-year follow-up study." Brazilian journal of medical & biological research, 585-592.	7
Dougherty, C. M., F. M. Lewis, E. A. Thompson, J. D. Baer and W. Kim (2004). "Short-term efficacy of a telephone intervention by expert nurses after an implantable cardioverter defibrillator." Pacing Clin Electrophysiol 27: 1594-1602.	3
Eggermont, L. H., D. L. Knol, E. M. Hol, D. F. Swaab and E. J. Scherder (2009). "Hand motor activity, cognition, mood, and the rest-activity rhythm in dementia: a clustered RCT." Behav Brain Res 196: 271-278.	3
Emery, C. F., R. L. Schein, E. R. Hauck and N. R. MacIntyre (1998). "Psychological and cognitive outcomes of a randomized trial of exercise among patients with chronic obstructive pulmonary disease." Health Psychol 17: 232-240.	3
Eyigor, S., H. Karapolat, H. Yesil, R. Uslu and B. Durmaz (2010). "Effects of pilates exercises on functional capacity, flexibility, fatigue, depression and quality of life in female breast cancer patients: a randomized controlled study." Eur J Phys Rehabil Med 46: 481-487.	3
Fitzsimmons, S. (2001). "Easy rider wheelchair biking. A nursing-recreation therapy clinical trial for the treatment of depression." J Gerontol Nurs 27(5): 14-23.	4
Fox, K. (2012). "Intervention design using self-determination theory to promote activity in depressed patients, older adults, and slimmers." Journal of Science and Medicine in Sport 15: S4.	9
Gary, R. (2006). "Exercise self-efficacy in older women with diastolic heart failure: results of a walking program and education intervention." J Gerontol Nurs 32: 31-39; quiz 40.	3

Gary, R. A., C. A. Sueta, M. Dougherty, B. Rosenberg, D. Cheek, J. Preisser, V. Neelon and R. McMurray (2004). "Home-based exercise improves functional performance and quality of life in women with diastolic heart failure." <i>Heart Lung</i> 33: 210-218.	3
Gary, R. and S. Y. Lee (2007). "Physical function and quality of life in older women with diastolic heart failure: effects of a progressive walking program on sleep patterns." <i>Prog Cardiovasc Nurs</i> 22: 72-80.	3
Gusi, N., M. C. Reyes, J. L. Gonzalez-Guerrero, E. Herrera and J. M. Garcia (2008). "Cost-utility of a walking programme for moderately depressed, obese, or overweight elderly women in primary care: a randomised controlled trial." <i>BMC Public Health</i> 8: 231.	3
Haboush, A., M. Floyd, J. Caron, M. LaSota and K. Alvarez (2006). "Ballroom dance lessons for geriatric depression: An exploratory study." <i>Arts in Psychotherapy</i> 33(2): 89-97.	3
Heckman, T. G., B. D. Heckman, T. Anderson, T. I. Lovejoy, D. Mohr, M. Sutton, J. A. Bianco and J. T. Gau (2013). "Supportive-expressive and coping group teletherapies for HIV-infected older adults: a randomized clinical trial." <i>AIDS Behav</i> 17(9): 3034-3044.	3
Herman, S., J. A. Blumenthal, M. Babyak, P. Khatri, W. E. Craighead, K. R. Krishnan and P. M. Doraiswamy (2002). "Exercise therapy for depression in middle-aged and older adults: predictors of early dropout and treatment failure." <i>Health Psychol</i> 21: 553-563.	3
Hoffman, B. M., J. A. Blumenthal, M. A. Babyak, P. J. Smith, S. D. Rogers, P. M. Doraiswamy and A. Sherwood (2008). "Exercise fails to improve neurocognition in depressed middle-aged and older adults." <i>Med Sci Sports Exerc</i> 40: 1344-1352.	3
Holmgren, E., G. Gosman-Hedstrom, B. Lindstrom and P. Wester (2010). "What is the benefit of a high-intensive exercise program on health-related quality of life and depression after stroke? A randomized controlled trial." <i>Advances in Physiotherapy</i> 12: 125-133.	3
Horowitz, A., J. P. Reinhardt and K. Boerner (2005). "The effect of rehabilitation on depression among visually disabled older adults." <i>Aging Ment Health</i> 9: 563-570.	3
Jeon, M. Y., E. S. Bark, E. G. Lee, J. S. Im, B. S. Jeong and E. S. Choe (2005). "[The effects of a Korean traditional dance movement program in elderly women]." <i>Taehan Kanho Hakhoe Chi</i> : 1268-1276.	3
Justine, M. and T. A. Hamid (2010). "A multicomponent exercise program for institutionalized older adults: effects on depression and quality of life." <i>J Gerontol Nurs</i> 36: 32-41.	3
Kerse, N., K. Falloon, S. A. Moyes, K. J. Hayman, T. Dowell, G. S. Kolt, C. R. Elley, S. Hatcher, K. Peri, S. Keeling, E. Robinson, J. Parsons, J. Wiles and B. Arroll (2008). "DeLLITE depression in late life: an intervention trial of exercise. Design and recruitment of a randomised controlled trial." <i>BMC Geriatr</i> 8: 12.	8
Kerse, N., K. J. Hayman, S. A. Moyes, K. Peri, E. Robinson, A. Dowell, G. S. Kolt, C. R. Elley, S. Hatcher, L. Kiata, J. Wiles, S. Keeling, J. Parsons and B. Arroll (2010). "Home-based activity program for older people with depressive symptoms: DeLLITE--a randomized controlled trial." <i>Ann Fam Med</i> 8: 214-223.	3
Kerse, N., K. Peri, E. Robinson, T. Wilkinson, M. von Radow, L. Kiata, J. Parsons, N. Latham, M. Parsons, J. Willingale, P. Brown and B. Arroll (2008). "Does a functional activity programme improve function, quality of life, and falls for residents in long term care? Cluster randomised controlled trial." <i>BMJ</i> 337: a1445.	3

Khatri, P., J. A. Blumenthal, M. A. Babyak, W. E. Craighead, S. Herman, T. Baldewicz, D. J. Madden, M. Doraiswamy, R. Waugh and K. R. Krishnan (2001). "Effects of exercise training on cognitive functioning among depressed older men and women." <i>J Aging Phys Act</i> 9: 43-57.	3
Lai, S. M., S. Studenski, L. Richards, S. Perera, D. Reker, S. Rigler and P. W. Duncan (2006). "Therapeutic exercise and depressive symptoms after stroke." <i>J Am Geriatr Soc</i> 54: 240-247.	3
Lavretsky-2011-Complementary use of tai chi chih augments escitalopram treatment of geriatric depression: a randomized controlled trial.	3
Legrand, F. D. and C. R. Mille (2009). "The effects of 60 minutes of supervised weekly walking (in a single vs. 3-5 session format) on depressive symptoms among older women: Findings from a pilot randomized trial." <i>Mental Health and Physical Activity</i> 2: 71-75.	5
Lim, Y. M. and G. R. Hong (2010) "Effect of 16-week Kouk-Sun-Do exercise on physical fitness, emotional state, and immunoglobulin A in community-dwelling elders in Korea." <i>Appl Nurs Res</i> , 91-100 DOI: 10.1016/j.apnr.2008.05.004.	3
Lincoln, A. K., A. Shepherd, P. L. Johnson and C. Castaneda-Sceppa (2011). "The impact of resistance exercise training on the mental health of older Puerto Rican adults with type 2 diabetes." <i>J Gerontol B Psychol Sci Soc Sci</i> 66: 567-570.	3
Maci, T., F. L. Pira, G. Quattrocchi, S. D. Nuovo, V. Perciavalle and M. Zappia (2012). "Physical and cognitive stimulation in Alzheimer Disease. the GAIA Project: a pilot study." <i>Am J Alzheimers Dis Other Demen</i> 27: 107-113.	3
Maki, Y., C. Ura, T. Yamaguchi, R. Takahashi and H. Yamaguchi (2012). "Intervention using a community-based walking program is effective for elderly adults with depressive tendencies." <i>J Am Geriatr Soc</i> 60: 1590-1591.	3
Martin, C. K., T. S. Church, A. M. Thompson, C. P. Earnest and S. N. Blair (2009). "Exercise dose and quality of life: a randomized controlled trial." <i>Arch Intern Med</i> 169: 269-278.	3
Mather, A. S., C. Rodriguez, M. F. Guthrie, A. M. McHarg, I. C. Reid and M. E. McMurdo (2002). "Effects of exercise on depressive symptoms in older adults with poorly responsive depressive disorder: randomised controlled trial." <i>Br J Psychiatry</i> 180: 411-415.	3
McCaffrey, R., P. Liehr, T. Gregersen and R. Nishioka (2011). "Garden Walking and Art Therapy for Depression in Older Adults." <i>Res Gerontol Nurs</i> 4: 237-242.	7
McMurdo, M. E. and L. Burnett (1992). "Randomised controlled trial of exercise in the elderly." <i>Gerontology</i> 38: 292-298.	3
McNeil, J. K., E. M. LeBlanc and M. Joyner (1991). "The effect of exercise on depressive symptoms in the moderately depressed elderly." <i>Psychol Aging</i> 6(3): 487-488.	3
Minor, M. A. and J. D. Brown (1993). "Exercise maintenance of persons with arthritis after participation in a class experience." <i>Health Educ Q</i> 20: 83-95.	3
Miser, W. F. (2000). "Exercise as an effective treatment option for major depression in older adults." <i>J Fam Pract</i> 49: 109-110.	10
Modabbernia, A., M. Shahidi, M. Mojtabah and A. Mojtabah (2012). "Reply to the Letter to the Editor Shahidi M, Mojtabah A, Modabbernia A., et al. 2011. Laughter yoga versus group exercise program in elderly depressed women: A randomized controlled trial. <i>Int J Geriatr Psychiatr</i> 26: 322-327. First things first! Caveats in research on "laughter yoga"." <i>Int J Geriatr Psychiatry</i> 27: 875-876.	10

Molloy, G. J., D. W. Johnston, C. Gao, M. D. Witham, J. M. Gray, I. S. Argo, A. D. Struthers and M. E. McMurdo (2006). "Effects of an exercise intervention for older heart failure patients on caregiver burden and emotional distress." <i>Eur J Cardiovasc Prev Rehabil</i> 13: 381-387.	3
Morgan, A. J., A. F. Jorm and A. J. Mackinnon (2012). "Usage and reported helpfulness of self-help strategies by adults with sub-threshold depression." <i>J Affect Disord</i> 136: 393-397.	3
Motl, R. W., J. F. Konopack, E. McAuley, S. Elavsky, G. J. Jerome and D. X. Marquez (2005). "Depressive symptoms among older adults: long-term reduction after a physical activity intervention." <i>J Behav Med</i> 28(4): 385-394.	3
Netz, Y., A. Yaretzki, I. Salganik, T. Jacob, B. Finkelov and E. Argov (1994). "The effect of supervised physical activity on cognitive and affective state of geriatric and psychogeriatric inpatients." <i>Clinical Gerontologist</i> 15: 47-56.	3
Neuberger, G. B., L. S. Aaronson, B. Gajewski, S. E. Embretson, P. E. Cagle, J. K. Loudon and P. A. Miller (2007). "Predictors of exercise and effects of exercise on symptoms, function, aerobic fitness, and disease outcomes of rheumatoid arthritis." <i>Arthritis Rheum</i> 57: 943-952.	3
O'Brien, M. E. (1999). "Progressive resistive exercise: effective antidepressant in the elderly..." <i>Complementary Medicine for the Physician</i> 4: 2-3.	9
Ohira, T., K. H. Schmitz, R. L. Ahmed and D. Yee (2006). "Effects of weight training on quality of life in recent breast cancer survivors: the Weight Training for Breast Cancer Survivors (WTBS) study." <i>Cancer</i> 106: 2076-2083.	3
Pakkala, I., S. Read, R. Leinonen, M. Hirvensalo, T. Lintunen and T. Rantanen (2008). "The effects of physical activity counseling on mood among 75- to 81-year-old people: a randomized controlled trial." <i>Prev Med</i> 46: 412-418.	3
Penninx, B. W., W. J. Rejeski, J. Pandya, M. E. Miller, B. Di, W. B. Applegate and M. Pahor (2002). "Exercise and depressive symptoms: a comparison of aerobic and resistance exercise effects on emotional and physical function in older persons with high and low depressive symptomatology." <i>J Gerontol B Psychol Sci Soc Sci</i> 57: P124-132.	3
Pereira, D. S., Q. Zille de, A. S. Miranda, N. P. Rocha, D. C. Felicio, E. C. Mateo, M. Favero, F. M. Coelho, F. Jesus-Moraleida, P. Gomes, A. D, A. L. Teixeira, P. Maximo and S. L (2013). "Effects of physical exercise on plasma levels of brain-derived neurotrophic factor and depressive symptoms in elderly women - a randomized clinical trial." <i>Arch Phys Med Rehabil</i> .	3
Pfaff, J. J., H. Alfonso, R. U. Newton, M. Sim, L. Flicker and O. P. Almeida (2014). "ACTIVEDEP: a randomised, controlled trial of a home-based exercise intervention to alleviate depression in middle-aged and older adults." <i>Br J Sports Med</i> 48(3): 226-232.	3
Pinniger, R., F. Brown Rhonda, B. Thorsteinsson Einar and P. McKinley (2013). "Tango programme for individuals with age-related macular degeneration." <i>British Journal of Visual Impairment</i> 31: 47-59.	3
Posadzki, P. (2013). "Tai chi chih improves inflammatory markers, cognition and quality of life scores in elderly depressed patients." <i>Focus on Alternative & Complementary Therapies</i> 18: 39-40.	9
Prakhinkit, S., S. Suppapitiporn, H. Tanaka and D. Suksom (2014). "Effects of Buddhism walking meditation on depression, functional fitness, and endothelium-dependent vasodilation in depressed elderly." <i>J Altern Complement Med</i> 20(5): 411-416.	3

Price, J. R. (2006). "A structured activity programme reduces depressive symptoms in moderately depressed older men with coronary heart disease, but not women." <i>Evidence Based Mental Health</i> 9: 17-17.	9
Proyer, R. T., W. Ruch and F. A. Rodden (2012). "Letter on Shahidi et al. (2011): "Laughter Yoga versus group exercise program in elderly depressed women: a randomized controlled trial" I -- first things first! Caveats in research on "laughter yoga"." <i>Int J Geriatr Psychiatry</i> 27: 873-874; author reply 875.	9
Rolland, Y., F. Pillard, A. Klapouszczak, E. Reynish, D. Thomas, S. Andrieu, D. Riviere and B. Vellas (2007). "Exercise program for nursing home residents with Alzheimer's disease: a 1-year randomized, controlled trial." <i>J Am Geriatr Soc</i> 55: 158-165.	3
Rosenberg, D., C. A. Depp, I. V. Vahia, J. Reichstadt, B. W. Palmer, J. Kerr, G. Norman and D. V. Jeste (2010). "Exergames for subsyndromal depression in older adults: a pilot study of a novel intervention." <i>Am J Geriatr Psychiatry</i> 18: 221-226.	7
Roux, M. C. and R. Kemp (2009) "Effect of a companion dog on depression and anxiety levels of elderly residents in a long-term care facility." <i>Psychogeriatrics</i> , 23-26.	3
Sam Yeol, W., K. Jong Ho and J. Jun Hyeok (2013). "Clinical Feasibility of Exercise Game for Depression Treatment in Older Women with Osteoarthritis: a Pilot Study." <i>Journal of Physical Therapy Science</i> 25(2): 165-167.	3
Scholz, U., N. Knoll, F. F. Sniehotta and R. Schwarzer (2006). "Physical activity and depressive symptoms in cardiac rehabilitation: long-term effects of a self-management intervention." <i>Soc Sci Med</i> 62: 3109-3120.	3
Shahidi, M., A. Mojtabed, A. Modabbernia, M. Mojtabed, A. Shafiabady, A. Delavar and H. Honari (2011). "Laughter yoga versus group exercise program in elderly depressed women: a randomized controlled trial." <i>Int J Geriatr Psychiatry</i> 26: 322-327.	3
Shaughnessy, A. F. (2013). "Exercise effective for depression in older patients." <i>American Family Physician</i> 88(12).	9
Shin, K. R. and J. S. Kim (2005). "[The effects of exercise program on knowledge and attitude of exercise and depression in low-income elderly women]." <i>Taehan Kanho Hakhoe Chi</i> : 1144-1152.	7
Sims, J., M. Galea, N. Taylor, K. Dodd, S. Jespersen, L. Joubert and J. Joubert (2009). "Regenerate: assessing the feasibility of a strength-training program to enhance the physical and mental health of chronic post stroke patients with depression." <i>Int J Geriatr Psychiatry</i> 24: 76-83.	3
Singh, N. A., K. M. Clements and M. A. Fiatarone (1997). "A randomized controlled trial of progressive resistance training in depressed elders." <i>J Gerontol A Biol Sci Med Sci</i> 52: M27-35.	3
Singh, N. A., K. M. Clements and M. A. Fiatarone (1997). "A randomized controlled trial of the effect of exercise on sleep." <i>Sleep</i> 20: 95-101.	3
Singh, N. A., K. M. Clements and M. A. Singh (2001). "The efficacy of exercise as a long-term antidepressant in elderly subjects: a randomized, controlled trial." <i>J Gerontol A Biol Sci Med Sci</i> 56: M497-504.	3
Singh, N. A., T. M. Stavrinou, Y. Scarbek, G. Galambos, C. Liber, S. Fiatarone and A. M (2005). "A randomized controlled trial of high versus low intensity weight training versus general practitioner care for clinical depression in older adults." <i>J Gerontol A Biol Sci Med Sci</i> 60: 768-776.	3

Singh, N. A., T. M. Stavrinos, Y. Skarbek, G. Galambos and M. A. Fiatarone-Singh (2000). "The effectiveness and appropriate intensity of progressive resistance training required to treat clinical depression in the elderly: a randomised controlled trial." <i>Australian and New Zealand Journal of Medicine</i> : 304.	9
Singh, N., K. Clements and M. Fiatarone (1996). "A randomised controlled trial of exercise in depressed elders." <i>Australian & New Zealand Journal of Medicine</i> : 599.	9
Srinivasan, S., L. P. Reagan, J. W. Hardin, M. Matthews, E. Leaphart, C. A. Grillo, M. Neese, A. Johnson, J. Gossard, S. Chapman and T. Brinkley (2014). "Adjunctive tai chi in geriatric depression with comorbid arthritis: A randomized, controlled trial." <i>American Journal of Geriatric Psychiatry</i> 22(3): S135-S136.	9
Steffens, D. C. (2013). "Exercise for late-life depression? It depends." <i>Lancet</i> 382(9886): 4-5.	9
Sung, K. (2009). "The effects of 16-week group exercise program on physical function and mental health of elderly Korean women in long-term assisted living facility." <i>J Cardiovasc Nurs</i> 24: 344-351.	3
Teri, L., L. E. Gibbons, S. M. McCurry, R. G. Logsdon, D. M. Buchner, W. E. Barlow, W. A. Kukull, A. Z. LaCroix, W. McCormick and E. B. Larson (2003). "Exercise plus behavioral management in patients with Alzheimer disease: a randomized controlled trial." <i>JAMA</i> 290: 2015-2022.	3
Timonen, L., T. Rantanen, T. E. Timonen and R. Sulkava (2002). "Effects of a group-based exercise program on the mood state of frail older women after discharge from hospital." <i>Int J Geriatr Psychiatry</i> 17: 1106-1111.	3
Tsang, H. W. H., C. K. Mok, Y. Au, T. Y and S. Y. C. Chan (2003). "The effect of qigong on general and psychosocial health of depressed elderly with chronic physical illnesses: a randomized clinical trial." <i>International Psychogeriatrics</i> .	3
Tsang, H. W., C. K. Mok, Y. Au, T. Y and S. Y. Chan (2003). "The effect of Qigong on general and psychosocial health of elderly with chronic physical illnesses: a randomized clinical trial." <i>Int J Geriatr Psychiatry</i> 18: 441-449.	3
Tsutsumi, T., B. M. Don, L. D. Zaichkowsky, K. Takenaka, K. Oka and T. Ohno (1998). "Comparison of high and moderate intensity of strength training on mood and anxiety in older adults." <i>Percept Mot Skills</i> 87(3 Pt 1): 1003-1011.	3
Underwood, M., S. E. Lamb, S. Eldridge, B. Sheehan, A. M. Slowther, A. Spencer, M. Thorogood, N. Atherton, S. A. Bremner, A. Devine, K. Diaz-Ordaz, D. R. Ellard, R. Potter, K. Spanjers and S. J. Taylor (2013). "Exercise for depression in elderly residents of care homes: a cluster-randomised controlled trial." <i>Lancet</i> .	3
Underwood, M., S. Lamb, B. Sheehan, D. R. Ellard and S. J. Taylor (2013). "Exercise for depression in elderly people - Authors' reply." <i>Lancet</i> 382(9901): 1325.	9
Underwood, M., S. Lamb, S. Eldridge, B. Sheehan, A. Slowther, A. Spencer, M. Thorogood, N. Atherton, S. Bremner, A. Devine, K. Diaz-Ordaz, D. Ellard, R. Potter, K. Spanjers and S. Taylor (2013). "Exercise for depression in care home residents: a randomised controlled trial with cost-effectiveness analysis (OPERA)." <i>Health Technol Assess</i> 17: 1-281.	3
Walker, J. G., A. J. Mackinnon, P. Batterham, A. F. Jorm, I. Hickie, A. McCarthy, M. Fenech and H. Christensen (2010). "Mental health literacy, folic acid and vitamin B12, and physical activity for the prevention of depression in older adults: randomised controlled trial." <i>Br J Psychiatry</i> 197: 45-54.	3

Vankova, H., I. Holmerova, K. Machacova, L. Volicer, P. Veleta and A. M. Celko (2014). "The effect of dance on depressive symptoms in nursing home residents." <i>J Am Med Dir Assoc</i> 15(8): 582-587.	3
Vanková, H., I. Holmerová, R. Andel, P. Veleta and H. Janecková (2008). "Functional status and depressive symptoms among older adults from residential care facilities in the Czech Republic." <i>Int J Geriatr Psychiatry</i> 23: 466-471.	7
Vasques, P. E., H. Moraes, H. Silveira, A. C. Deslandes and J. Laks (2011) "Acute exercise improves cognition in the depressed elderly: the effect of dual-tasks." <i>Clinics (Sao Paulo, Brazil)</i> , 1553-1557.	7
Villaverde, G., L. Torres, M. Abalos, M. G, C. Argente del, J. M, I. M. Guisado, B. Guisado and R. Ramirez (2012). "Influence of exercise on mood in postmenopausal women." <i>J Clin Nurs</i> 21: 923-928.	3
Yang, K. H., Y. H. Kim and M. S. Lee (2005). "Efficacy of Qi-therapy (external Qigong) for elderly people with chronic pain." <i>Int J Neurosci</i> 115: 949-963.	3

Övriga behandlingar

Kombination

Referens	Exklusions- orsak
(2003). "Erratum: Effect of Improving Depression Care on Pain and Functional Outcomes among Older Adults with Arthritis: A Randomized Controlled Trial: (Journal of the American Medical Association (November 12, 2003) 290 (2428-2434))." <i>Journal of the American Medical Association</i> 290: 2803.	9
Aburizik, A., L. Dindo, P. Kaboli, M. Charlton, K. Dawn and C. Turvey (2013). "A pilot randomized controlled trial of a depression and disease management program delivered by phone." <i>J Affect Disord</i> 151(2): 769-774.	3
Akerblad, A. C., F. Bengtsson, L. Ekselius and K. von (2003). "Effects of an educational compliance enhancement programme and therapeutic drug monitoring on treatment adherence in depressed patients managed by general practitioners." <i>Int Clin Psychopharmacol</i> 18: 347-354.	3
Albus, C., B. Bjarnson-Wehrens, D. B. Gysan, G. Herold, C. A. Schneider, E. zu, H. G. Predel and S. PraFord (2012). "[The effects of multimodal intervention for the primary prevention of cardiovascular diseases on depression, anxiety, and Type-D pattern: initial results of the randomized controlled PreFord trial]." <i>Herz</i> 37: 59-62.	3
Alexopoulos, G. S., C. F. Reynolds, rd, M. L. Bruce, I. R. Katz, P. J. Raue, B. H. Mulsant, D. W. Oslin, H. Ten and P. Group (2009). "Reducing suicidal ideation and depression in older primary care patients: 24-month outcomes of the PROSPECT study." <i>Am J Psychiatry</i> 166: 882-890.	3
Andohek, K. M. and S. Banerjee (1997). "Psychogeriatric home care led to more elderly patients recovering from depression." <i>Evidence-Based Medicine</i> 2: 86.	8
Apil, S. R., E. Hoencamp, H. Judith, M. P and P. Spinhoven (2012). "A stepped care relapse prevention program for depression in older people: a randomized controlled trial." <i>Int J Geriatr Psychiatry</i> 27: 583-591.	3
Araya, R., G. Rojas, R. Fritsch, J. Gaete, M. Rojas, G. Simon and T. J. Peters (2003). "Treating depression in primary care in low-income women in Santiago, Chile: A randomised controlled trial." <i>Lancet</i> 361: 995-1000.	3
Arean, P. A., L. Ayalon, E. Hunkeler, E. H. Lin, L. Tang, L. Harpole, H. Hendrie, J. W. Williams, Jr, J. Unutzer and I. Investigators (2005). "Improving depression care for older, minority patients in primary care." <i>Med Care</i> 43: 381-390.	3
Arean, P. A., A. M. Gum, L. Tang and J. Unutzer (2007). "Service use and outcomes among elderly persons with low incomes being treated for depression." <i>Psychiatr Serv</i> 58: 1057-1064.	7
Arean, P. A., C. E. McCulloch, A. Bostrom, A. Gum, D. Gallagher-Thompson and L. Thompson (2005). "Treatment of depression in low-income older adults." <i>Psychol Aging</i> 20: 601-609.	3
Arnow, B. A., R. Manber, C. Blasey, J. A. Blalock, B. O. Rothbaum, M. E. Thase, D. Vivian, D. N. Klein, J. C. Markowitz, A. J. Rush, L. P. Riso, McCullough, Jr., P. J and M. B. Keller (2003). "Therapeutic Reactance as a Predictor of Outcome in the Treatment of Chronic Depression." <i>Journal of Consulting and Clinical Psychology</i> 71: 1025-1035.	3

Arthur, A. J., C. Jagger, J. Lindesay and R. J. Matthews (2002). "Evaluating a mental health assessment for older people with depressive symptoms in general practice: a randomised controlled trial." <i>Br J Gen Pract</i> 52: 202-207.	4
Azar, A. R., M. P. Chopra, L. Y. Cho, E. Coakley and J. L. Rudolph (2011). "Remission in major depression: results from a geriatric primary care population." <i>Int J Geriatr Psychiatry</i> 26: 48-55.	7
Banerjee, S., A. Mann, K. Shamash and A. MacDonald (1997). "Effect of psychogeriatric team on depression in frail elderly people at home [letter; author's reply]." <i>BMJ</i> : 378.	9
Banerjee, S., K. Shamash, A. J. Macdonald and A. H. Mann (1996). "Randomised controlled trial of effect of intervention by psychogeriatric team on depression in frail elderly people at home." <i>BMJ</i> 313: 1058-1061.	4
Bares, M., M. Kopecek, T. Novak, P. Stopkova, P. Sos, J. Kozeny, M. Brunovsky and C. Hoschl (2009). "Low frequency (1-Hz), right prefrontal repetitive transcranial magnetic stimulation (rTMS) compared with venlafaxine ER in the treatment of resistant depression: A double-blind, single-centre, randomized study." <i>J Affect Disord</i> 118: 94-100.	4
Belle, S. H., L. Burgio, R. Burns, D. Coon, S. J. Czaja, D. Gallagher-Thompson, L. N. Gitlin, J. Klinger, K. M. Koepke, C. C. Lee, J. Martindale-Adams, L. Nichols, R. Schulz, S. Stahl, A. Stevens, L. Winter, S. Zhang, H. Resources for Enhancing Alzheimer's Caregiver and I. I. Investigators (2006). "Enhancing the quality of life of dementia caregivers from different ethnic or racial groups: a randomized, controlled trial." <i>Ann Intern Med</i> 145: 727-738.	3
Beutler, L. E., F. Scogin, P. Kirkish, D. Schretlen, A. Corbishley, D. Hamblin, K. Meredith, R. Potter, C. R. Bamford and A. I. Levenson (1987). "Group cognitive therapy and alprazolam in the treatment of depression in older adults." <i>Journal of Consulting and Clinical Psychology</i> : 550-556.	4
Birkenhager, T. K., B. van den, W. W, P. Moleman and J. A. Bruijn (2006). "Outcome of a 4-step treatment algorithm for depressed inpatients." <i>J Clin Psychiatry</i> 67: 1266-1271.	3
Blanchard, M. R., A. Waterreus and A. H. Mann (1995). "The effect of primary care nurse intervention upon older people screened as depressed." <i>Int J Geriatr Psychiatry</i> 10: 289-298.	4
Blumenthal, J. A., A. Sherwood, M. A. Babyak, L. L. Watkins, P. J. Smith, B. M. Hoffman, C. V. O'Hayer, S. Mabe, J. Johnson, P. M. Doraiswamy, W. Jiang, D. D. Schocken and A. L. Hinderliter (2012). "Exercise and pharmacological treatment of depressive symptoms in patients with coronary heart disease: results from the UPBEAT (Understanding the Prognostic Benefits of Exercise and Antidepressant Therapy) study." <i>J Am Coll Cardiol</i> 60: 1053-1063.	3
Boggio, P. S., F. Fregni, F. Bermpohl, C. G. Mansur, M. Rosa, D. O. Rumi, E. R. Barbosa, R. Odebrecht, A. Pascual-Leone, S. P. Rigonatti, M. A. Marcolin, S. Araujo and T. M (2005). "Effect of repetitive TMS and fluoxetine on cognitive function in patients with Parkinson's disease and concurrent depression." <i>Mov Disord</i> 20: 1178-1184.	4
Bogner, H. R., M. L. Bruce, C. F. Reynolds, rd, B. H. Mulsant, M. S. Cary, K. Morales, G. S. Alexopoulos and P. Group (2007). "The effects of memory, attention, and executive dysfunction on outcomes of depression in a primary care intervention trial: the PROSPECT study." <i>Int J Geriatr Psychiatry</i> 22: 922-929.	4
Bogner, H. R., M. S. Cary, M. L. Bruce, C. F. Reynolds, rd, B. Mulsant, H. Ten and G. S. Alexopoulos (2005). "The role of medical comorbidity in outcome of major depression in primary care: the PROSPECT study." <i>Am J Geriatr Psychiatry</i> 13: 861-868.	7

Bogner, H. R., V. de and F. H (2008). "Integration of depression and hypertension treatment: a pilot, randomized controlled trial." <i>Ann Fam Med</i> 6: 295-301.	3
Bogner, H. R., V. de and F. H (2010). "Integrating type 2 diabetes mellitus and depression treatment among African Americans: a randomized controlled pilot trial." <i>Diabetes Educ</i> 36: 284-292.	3
Bogner, H. R., K. H. Morales, V. de, F. H and A. R. Cappola (2012). "Integrated management of type 2 diabetes mellitus and depression treatment to improve medication adherence: a randomized controlled trial." <i>Ann Fam Med</i> 10: 15-22.	3
Bogner, H. R., K. H. Morales, C. F. Reynolds, M. S. Cary and M. L. Bruce (2012). "Prognostic factors, course, and outcome of depression among older primary care patients: the PROSPECT study." <i>Aging Ment Health</i> 16: 452-461.	7
Bosmans, J., M. de Bruijne, H. van Hout, H. van Marwijk, A. Beekman, L. Bouter, W. Stalman and M. van Tulder (2006). "Cost-effectiveness of a disease management program for major depression in elderly primary care patients." <i>J Gen Intern Med</i> 21(10): 1020-1026.	6
Bradshaw David, H., K. Kumpfer, S. Trevithick and J. Stoddard Gregory (2010). "Outcome evaluation of the Veterans Affairs Salt Lake City Integrative Health Clinic for chronic pain and stress-related depression, anxiety, and post-traumatic stress disorder." <i>Journal of Alternative & Complementary Medicine</i> 16: 823-835.	3
Brandon, S. T., B. Bair and R. Kathol (2011). "Understanding the importance of time in delivering evidence-based major depression treatment to geriatric patients in the primary care setting." <i>Clinical Geriatrics</i> 19: 40-43.	9
Browne, G., M. Steiner, J. Roberts, A. Gafni, C. Byrne, E. Dunn, B. Bell, M. Mills, L. Chalklin, D. Wallik and J. Kraemer (2002). "Sertraline and/or interpersonal psychotherapy for patients with dysthymic disorder in primary care: 6-month comparison with longitudinal 2-year follow-up of effectiveness and costs." <i>J Affect Disord</i> 68: 317-330.	3
Bruce, M. L., H. Ten, R. T, C. F. Reynolds, rd, Katz, II, H. C. Schulberg, B. H. Mulsant, G. K. Brown, G. J. McAvay, J. L. Pearson and G. S. Alexopoulos (2004). "Reducing suicidal ideation and depressive symptoms in depressed older primary care patients: a randomized controlled trial." <i>JAMA</i> 291: 1081-1091.	3
Buettner, L. L. and S. Fitzsimmons (2002). "AD-venture program: therapeutic biking for the treatment of depression in long-term care residents with dementia." <i>Am J Alzheimers Dis Other Demen</i> 17(2): 121-127.	4
Buyse, D. J., C. F. Reynolds, rd, C. C. Hoch, P. R. Houck, D. J. Kupfer, S. Mazumdar and E. Frank (1996). "Longitudinal effects of nortriptyline on EEG sleep and the likelihood of recurrence in elderly depressed patients." <i>Neuropsychopharmacology</i> 14: 243-252.	6
Callahan, C. M., H. C. Hendrie, R. S. Dittus, D. C. Brater, S. L. Hui and W. M. Tierney (1994). "Improving treatment of late life depression in primary care: a randomized clinical trial." <i>J Am Geriatr Soc</i> 42: 839-846.	3
Callahan, C. M., K. Kroenke, S. R. Counsell, H. C. Hendrie, A. J. Perkins, W. Katon, P. Hitchcock, L. Harpole, E. M. Hunkeler and J. Unutzer (2005). "Treatment of depression improves physical functioning in older adults." <i>J Am Geriatr Soc</i> 53: 367-373.	3

Campbell, N. C., J. Thain, H. G. Deans, L. D. Ritchie, J. M. Rawles and J. L. Squair (1998). "Secondary prevention clinics for coronary heart disease: randomised trial of effect on health." <i>BMJ</i> 316: 1434-1437.	3
Cardoso, E. F., F. Fregni, M. Martins, P. S. Boggio, M. Luis, K. Coracini, V. Lopes, L. M. Melo, J. R. Sato, M. Antonio, S. P. Rigonatti, A. C. Cruz, Jr, B. Reis, E. Amaro and Jr (2008). "rTMS treatment for depression in Parkinson's disease increases BOLD responses in the left prefrontal cortex." <i>Int J Neuropsychopharmacol</i> 11: 173-183.	4
Carroll, B. J. (2004). "Erratum: Sertraline and the cheshire cat in geriatric depression (American Journal of Psychiatry (April 2004) 161 (759))." <i>American Journal of Psychiatry</i> 161: 1145-1146.	9
Chalder, M., N. J. Wiles, J. Campbell, S. P. Hollinghurst, A. M. Haase, A. H. Taylor, K. R. Fox, C. Costelloe, A. Searle, H. Baxter, R. Winder, C. Wright, K. M. Turner, M. Calnan, D. A. Lawlor, T. J. Peters, D. J. Sharp, A. A. Montgomery and G. Lewis (2012). "Facilitated physical activity as a treatment for depressed adults: randomised controlled trial." <i>BMJ</i> 344: e2758.	3
Chan, D., M. Y. Fan and J. Unutzer (2011). "Long-term effectiveness of collaborative depression care in older primary care patients with and without PTSD symptoms." <i>Int J Geriatr Psychiatry</i> 26: 758-764.	6
Chapman, D. G. and R. W. Toseland (2007). "Effectiveness of advanced illness care teams for nursing home residents with dementia." <i>Soc Work</i> 52: 321-329.	3
Chen, H., E. H. Coakley, K. Cheal, J. Maxwell, G. Costantino, D. D. Krahn, R. G. Malgady, U. N. Durai, L. M. Quijano, S. Zaman, C. J. Miller, J. H. Ware, H. Chung, C. Aoyama, S. Van, W. W and S. E. Levkoff (2006). "Satisfaction with mental health services in older primary care patients." <i>Am J Geriatr Psychiatry</i> 14: 371-379.	3
Chen, X. (1992). "[Active music therapy for senile depression]." <i>Zhonghua Shen Jing Jing Shen Ke Za Zhi</i> 25(4): 208-210, 252-203.	11
Chenderes-2009-Treatment of late life depression associated with mild cognitive impairment.	Ej läst
Ciarambino, T., P. Castellino, G. Paolisso, L. Coppola, N. Ferrara, G. Signoriello and M. Giordano (2012). "Long term effects of low protein diet on depressive symptoms and quality of life in elderly Type 2 diabetic patients." <i>Clin Nephrol</i> 78: 122-128.	3
Ciechanowski, P., E. Wagner, K. Schmaling, S. Schwartz, B. Williams, P. Diehr, J. Kulzer, S. Gray, C. Collier and J. LoGerfo (2004). "Community-integrated home-based depression treatment in older adults: a randomized controlled trial." <i>JAMA</i> 291(13): 1569-1577.	3
Cole, M. G., J. McCusker, M. Elie, N. Dendukuri, E. Latimer and E. Belzile (2006). "Systematic detection and multidisciplinary care of depression in older medical inpatients: a randomized trial." <i>CMAJ</i> 174: 38-44.	4
Coleman, E. A., L. C. Grothaus, N. Sandhu and E. H. Wagner (1999). "Chronic care clinics: A randomized controlled trial of a new model of primary care for frail older adults." <i>J Am Geriatr Soc</i> 47(7): 775-783.	3
Conn, D. K. (2005). "Collaborative care depression management for older adults: level of comorbidity does not affect outcome." <i>Evidence Based Mental Health</i> 8(4): 105-105.	9

Conradi, H. J., J. de, H. Kluiter, A. Smit, M. van der, J. A. Jenner, O. van, W. T, P. M. Emmelkamp and J. Ormel (2007). "Enhanced treatment for depression in primary care: long-term outcomes of a psycho-educational prevention program alone and enriched with psychiatric consultation or cognitive behavioral therapy." <i>Psychol Med</i> 37: 849-862.	3
Coon, D. W. and L. W. Thompson (2003). "The relationship between homework compliance and treatment outcomes among older adult outpatients with mild-to-moderate depression." <i>Am J Geriatr Psychiatry</i> 11: 53-61.	7
Cowan, M. J., K. E. Freedland, M. M. Burg, P. G. Saab, M. E. Youngblood, C. E. Cornell, L. H. Powell, S. M. Czajkowski and E. Investigators (2008). "Predictors of treatment response for depression and inadequate social support--the ENRICHD randomized clinical trial." <i>Psychother Psychosom</i> 77: 27-37.	7
Coyne, J. C., J. Koppel, C. C. Colenda, M. L. Bruce, G. S. Alexopoulos, H. C. Schulberg, H. Ten, I. R. Katz, C. F. Reynolds and Iii (2004). "Interventions for treatment of depression in primary care... Bruce ML, Ten Have TR, Reynolds CF III et al. Reducing suicidal ideation and depressive symptoms in depressed older primary care patients: a randomized controlled trial. <i>JAMA</i> . 2004;291:1081-1091." <i>JAMA: Journal of the American Medical Association</i> 291: 2814-2816.	9
Cuijpers, P. and P. Lammeren (2001). "Secondary prevention of depressive symptoms in elderly inhabitants of residential homes." <i>Int J Geriatr Psychiatry</i> : 702-708.	7
Cullum, S., S. Tucker, C. Todd and C. Brayne (2007). "Effectiveness of liaison psychiatric nursing in older medical inpatients with depression: a randomised controlled trial." <i>Age Ageing</i> 36: 436-442.	4
Czajkowski, S. M. (2003). "Effects of Treating Depression and Low Perceived Social Support on Clinical Events after Myocardial Infarction: The Enhancing Recovery in Coronary Heart Disease Patients (ENRICHD) Randomized Trial." <i>Journal of the American Medical Association</i> 289: 3106-3116.	3
Davidson, K. W., N. Rieckmann, L. Clemow, J. E. Schwartz, D. Shimbo, V. Medina, G. Albanese, I. Kronish, M. Hegel and M. M. Burg (2010). "Enhanced depression care for patients with acute coronary syndrome and persistent depressive symptoms: coronary psychosocial evaluation studies randomized controlled trial." <i>Arch Intern Med</i> 170: 600-608.	3
Davison Tanya, E., G. Karantzas, D. Mellor, P. McCabe Marita and D. Mrkic (2013). "Staff-focused interventions to increase referrals for depression in aged care facilities: A cluster randomized controlled trial." <i>Aging Ment Health</i> 17: 449-455.	6
de, J., M. HendrickSEN, A. van, S. Kool, V. Peen, R. Van, E. van den and J. Dekker (2004). "Psychotherapy alone and combined with pharmacotherapy in the treatment of depression." <i>Br J Psychiatry</i> 185: 37-45.	3
Deen, T. L., J. C. Fortney and J. M. Pyne (2011). "Relationship between satisfaction, patient-centered care, adherence and outcomes among patients in a collaborative care trial for depression." <i>Adm Policy Ment Health</i> 38: 345-355.	3
Denman, A. M. (2004). "Enhanced depression care improved arthritis pain and function in older patients." <i>ACP Journal Club</i> 141: 14-14.	3
Dietrich, A. J., T. E. Oxman, Williams, Jr., W. J, H. C. Schulberg, M. L. Bruce, P. W. Lee, S. Barry, P. J. Raue, J. J. Lefever, M. Heo, K. Rost, K. Kroenke, M. Gerrity and P. A. Nutting (2004). "Re-engineering systems for the treatment of depression in primary care: Cluster randomised controlled trial." <i>British Medical Journal</i> 329: 602-605.	3

Dombrovski, A. Y., B. H. Mulsant, P. R. Houck, S. Mazumdar, E. J. Lenze, C. Andreescu, J. M. Cyranowski, C. F. Reynolds and rd (2007). "Residual symptoms and recurrence during maintenance treatment of late-life depression." <i>J Affect Disord</i> 103: 77-82.	7
Dozeman, E., D. J. van Schaik, A. T. Beekman, W. A. Stalman, J. E. Bosmans and H. W. van Marwijk (2007). "Depression and anxiety, an indicated prevention (DIP) protocol in homes for the elderly: feasibility and (cost) effectiveness of a stepped care programme." <i>BMC Geriatr</i> 7: 6.	8
Ell, K., M. P. Aranda, B. Xie, P.-J. Lee and C.-P. Chou (2010). "Collaborative depression treatment in older and younger adults with physical illness: Pooled comparative analysis of three randomized clinical trials." <i>The American Journal of Geriatric Psychiatry</i> 18(6): 520-530.	9
Ell, K., W. Katon, B. Xie, P. J. Lee, S. Kapetanovic, J. Guterman and C. P. Chou (2011). "One-year postcollaborative depression care trial outcomes among predominantly Hispanic diabetes safety net patients." <i>Gen Hosp Psychiatry</i> 33: 436-442.	3
Ell, K., J. Unutzer, M. Aranda, N. E. Gibbs, P. J. Lee and B. Xie (2007). "Managing depression in home health care: a randomized clinical trial." <i>Home Health Care Serv Q</i> 26(3): 81-104.	4
Fann, J. R., M. Y. Fan and J. Unutzer (2009). "Improving primary care for older adults with cancer and depression." <i>J Gen Intern Med</i> 24 Suppl 2: S417-424.	3
Feng, S. Z., M. Y. Zhang and Z. H. Dai (2005). "Impacts of rehabilitative therapy on post-stroke depression and the ability of daily life." <i>Chinese Journal of Clinical Rehabilitation</i> 9: 154-155.	3
Fernandez, H. and R. M (1994). "The efficacy of psychological intervention for depression in chronic hemodialysis patients." <i>Psoriasis</i> 15: 39-42.	10
Finkenzeller, W., I. Zobel, S. Rietz, E. Schramm and M. Berger (2009). "[Interpersonal psychotherapy and pharmacotherapy for post-stroke depression. Feasibility and effectiveness]." <i>Nervenarzt</i> 80: 805-812.	10
Gallo, J. J., H. R. Bogner, K. H. Morales, E. P. Post, J. Y. Lin and M. L. Bruce (2007). "The effect of a primary care practice-based depression intervention on mortality in older adults: a randomized trial." <i>Ann Intern Med</i> 146(10): 689-698.	3
Gallo, J. J., K. H. Morales, H. R. Bogner, P. J. Raue, J. Zee, M. L. Bruce and C. F. Reynolds, 3rd (2013). "Long term effect of depression care management on mortality in older adults: follow-up of cluster randomized clinical trial in primary care." <i>Bmj</i> 346: f2570.	4
Gensichen, J. (2006). "IMPACT collaborative care improves depression in elderly patients in primary care in the longer term." <i>Evidence Based Mental Health</i> 9: 76-76.	9
Gilbody, S. M. (2007). "IMPACT collaborative care programme reduces suicide ideation in depressed older adults." <i>Evidence Based Mental Health</i> 10: 51-51.	9
Gottfries, C. G., I. Karlsson and A. L. Nyth (1992). "Treatment of depression in elderly patients with and without dementia disorders." <i>Int Clin Psychopharmacol</i> : 55-64.	9
Hargus, E., C. Crane, T. Barnhofer and J. M. G. Williams (2010). "Effects of Mindfulness on Meta-Awareness and Specificity of Describing Prodromal Symptoms in Suicidal Depression." <i>Emotion</i> 10: 34-42.	3
Harpole, L. H., J. W. Williams, Jr, M. K. Olsen, K. M. Stechuchak, E. Oddone, C. M. Callahan, W. J. Katon, E. H. Lin, L. M. Grypma and J. Unutzer (2005). "Improving depression outcomes in older adults with comorbid medical illness." <i>Gen Hosp Psychiatry</i> 27: 4-12.	7

Hedrick, S. C., E. F. Chaney, B. Felker, C. F. Liu, N. Hasenberg, P. Heagerty, J. Buchanan, R. Bagala, D. Greenberg, G. Paden, S. D. Fihn and W. Katon (2003). "Effectiveness of collaborative care depression treatment in Veterans' Affairs primary care." <i>J Gen Intern Med</i> 18: 9-16.	3
Hegel, M. T., J. Unutzer, L. Tang, P. A. Arean, W. Katon, P. H. Noel, J. W. Williams, Jr and E. H. Lin (2005). "Impact of comorbid panic and posttraumatic stress disorder on outcomes of collaborative care for late-life depression in primary care." <i>Am J Geriatr Psychiatry</i> 13: 48-58.	7
Hegerl, U., M. Hautzinger, R. Mergl, R. Kohnen, M. Schutze, W. Scheunemann, A. K. Allgaier, J. Coyne and V. Henkel (2010). "Effects of pharmacotherapy and psychotherapy in depressed primary-care patients: A randomized, controlled trial including a patients' choice arm." <i>International Journal of Neuropsychopharmacology</i> 13: 31-44.	3
Huffman, J. C., C. A. Mastromauro, G. Sowden, G. L. Fricchione, B. C. Healy and J. L. Januzzi (2011). "Impact of a depression care management program for hospitalized cardiac patients." <i>Circ Cardiovasc Qual Outcomes</i> 4: 198-205.	3
Hunkeler, E. M., W. Katon, L. Tang, J. W. Williams, Jr, K. Kroenke, E. H. Lin, L. H. Harpole, P. Arean, S. Levine, L. M. Grypma, W. A. Hargreaves and J. Unutzer (2006). "Long term outcomes from the IMPACT randomised trial for depressed elderly patients in primary care." <i>BMJ</i> 332: 259-263.	3
Hunkeler, E. M., J. F. Meresman, W. A. Hargreaves, B. Fireman, W. H. Berman, A. J. Kirsch, J. Groebe, S. W. Hurt, P. Braden, M. Getzell, P. A. Feigenbaum, T. Peng and M. Salzer (2000). "Efficacy of nurse telehealth care and peer support in augmenting treatment of depression in primary care." <i>Arch Fam Med</i> 9: 700-708.	3
Jenkins, D. and A. MacDonald (1994). "Should general practitioners refer more of their elderly depressed patients to psychiatric services?" <i>Int J Geriatr Psychiatry</i> 9: 461-465.	4
Joubert, J., C. Reid, L. Joubert, D. Barton, D. Ruth, D. Jackson, J. O. Sullivan and S. M. Davis (2006). "Risk factor management and depression post-stroke: The value of an integrated model of care." <i>Journal of Clinical Neuroscience</i> 13: 84-90.	3
Katon, W., J. Russo, E. Frank, J. Barrett, J. W. Williams, Jr, T. Oxman, M. Sullivan and J. Cornell (2002). "Predictors of nonresponse to treatment in primary care patients with dysthymia." <i>Gen Hosp Psychiatry</i> 24: 20-27.	7
Katon, W., J. Unutzer, M. Y. Fan, J. W. Williams, Jr., M. Schoenbaum, E. H. Lin and E. M. Hunkeler (2006). "Cost-effectiveness and net benefit of enhanced treatment of depression for older adults with diabetes and depression." <i>Diabetes Care</i> 29(2): 265-270.	6
Katon, W. J., K. Von, E. H. B. Lin, G. Simon, E. Ludman, J. Russo, P. Ciechanowski, E. Walker and T. Bush (2004). "The pathways study: A randomized trial of collaborative care in patients with diabetes and depression." <i>Arch Gen Psychiatry</i> 61: 1042-1049.	3
Kinder, L. S., W. J. Katon, E. Ludman, J. Russo, G. Simon, E. H. Lin, P. Ciechanowski, K. Von and B. Young (2006). "Improving depression care in patients with diabetes and multiple complications." <i>J Gen Intern Med</i> 21: 1036-1041.	3
Klug, G., G. Hermann, B. Fuchs-Nieder, M. Panzer, A. Haider-Stipacek, H. G. Zapotoczky and S. Priebe (2011). "Clinically and economically effectiveness of home treatment for elderly patients with depression: Randomised controlled trial." <i>Journal of Mental Health Policy and Economics</i> 14: S15.	4

Kocsis, J. H., A. C. Leon, J. C. Markowitz, R. Manber, B. Arnow, D. N. Klein and M. E. Thase (2009). "Patient preference as a moderator of outcome for chronic forms of major depressive disorder treated with nefazodone, cognitive behavioral analysis system of psychotherapy, or their combination." <i>Journal of Clinical Psychiatry</i> 70: 354-361.	3
Kominski, G., R. Andersen, R. Bastani, R. Gould, C. Hackman, D. Huang, L. Jarvik, A. Maxwell, J. Moye, E. Olsen, R. Rohrbaugh, J. Rosansky, S. Taylor and S. Van (2001). "UPBEAT: the impact of a psychogeriatric intervention in VA medical centers. Unified Psychogeriatric Biopsychosocial Evaluation and Treatment." <i>Med Care</i> 39: 500-512.	3
Krahn, D. D., S. J. Bartels, E. Coakley, D. W. Oslin, H. Chen, J. McIntyre, H. Chung, J. Maxwell, J. Ware and S. E. Levkoff (2006). "PRISM-E: comparison of integrated care and enhanced specialty referral models in depression outcomes." <i>Psychiatr Serv</i> 57: 946-953.	4
Kravitz, R. L., P. Franks, M. D. Feldman, D. J. Tancredi, C. A. Slee, R. M. Epstein, P. R. Duberstein, R. A. Bell, M. Jackson-Triche, D. A. Paterniti, C. Cipri, A. M. Iosif, S. Olson, S. Kelly-Reif, A. Hudnut, S. Dvorak, C. Turner and A. Jerant (2013). "Patient engagement programs for recognition and initial treatment of depression in primary care: a randomized trial." <i>Jama</i> 310(17): 1818-1828.	3
Kroenke, K., M. J. Bair, T. M. Damush, J. Wu, S. Hoke, J. Sutherland and W. Tu (2009). "Optimized antidepressant therapy and pain self-management in primary care patients with depression and musculoskeletal pain: a randomized controlled trial." <i>JAMA</i> 301: 2099-2110.	3
Kroenke, K., J. Wu, M. J. Bair, E. E. Krebs, T. M. Damush and W. Tu (2011). "Reciprocal relationship between pain and depression: a 12-month longitudinal analysis in primary care." <i>J Pain</i> 12: 964-973.	3
Leontjevas, R., D. L. Gerritsen, M. Smalbrugge, S. Teerenstra, M. J. Vernooij-Dassen and R. T. Koopmans (2013). "A structural multidisciplinary approach to depression management in nursing-home residents: a multicentre, stepped-wedge cluster-randomised trial." <i>Lancet</i> 381(9885): 2255-2264.	3
Leontjevas, R., S. Teerenstra, M. Smalbrugge, M. J. Vernooij-Dassen, E. T. Bohlmeijer, D. L. Gerritsen and R. T. Koopmans (2013). "More insight into the concept of apathy: a multidisciplinary depression management program has different effects on depressive symptoms and apathy in nursing homes." <i>Int Psychogeriatr</i> 25(12): 1941-1952.	9
Lesperance, F., N. Frasure-Smith, D. Koszycki, M. A. Laliberte, Z. van, T. L, B. Baker, J. R. Swenson, K. Ghatali, B. L. Abramson, P. Dorian, M. C. Guertin and C. Investigators (2007). "Effects of citalopram and interpersonal psychotherapy on depression in patients with coronary artery disease: the Canadian Cardiac Randomized Evaluation of Antidepressant and Psychotherapy Efficacy (CREATE) trial." <i>JAMA</i> 297: 367-379.	3
Lin, E. H., K. Von, P. Ciechanowski, D. Peterson, E. J. Ludman, C. M. Rutter, M. Oliver, B. A. Young, J. Gensichen, M. McGregor, D. K. McCulloch, E. H. Wagner and W. J. Katon (2012). "Treatment adjustment and medication adherence for complex patients with diabetes, heart disease, and depression: a randomized controlled trial." <i>Ann Fam Med</i> 10: 6-14.	3

Lin, E. H. B., W. Katon, K. Von, L. Tang, Williams, Jr., W. J. K. Kroenke, E. Hunkeler, L. Harpole, M. Hegel, P. Arean, M. Hoffing, P. Della, C. Langston and J. Unutzer (2003). "Effect of Improving Depression Care on Pain and Functional Outcomes among Older Adults with Arthritis: A Randomized Controlled Trial." <i>Journal of the American Medical Association</i> 290: 2428-2434.	3
Lin, E. H. B., K. Von, E. J. Ludman, C. Rutter, T. M. Bush, G. E. Simon, J. Unutzer, E. Walker and W. J. Katon (2003). "Enhancing adherence to prevent depression relapse in primary care." <i>Gen Hosp Psychiatry</i> 25: 303-310.	3
Linden, M., A. Westram, L. G. Schmidt and C. Haag (2008). "Impact of the WHO depression guideline on patient care by psychiatrists: a randomized controlled trial." <i>Eur Psychiatry</i> 23: 403-408.	3
Liu, S. J., C. J. Lin, Y. M. Chen and X. Y. Huang (2007). "The effects of reminiscence group therapy on self-esteem, depression, loneliness and life satisfaction of elderly people living alone." <i>Mid-Taiwan Journal of Medicine</i> 12: 133-142.	11
Llewellyn-Jones, R. H., K. A. Baikie, H. Smithers, J. Cohen, J. Snowdon and C. C. Tennant (1999). "Multifaceted shared care intervention for late life depression in residential care: randomised controlled trial." <i>BMJ</i> 319: 676-682.	7
Lund, A., M. Michelet, L. Sandvik, T. Wyller and U. Sveen (2012). "A lifestyle intervention as supplement to a physical activity programme in rehabilitation after stroke: a randomized controlled trial." <i>Clin Rehabil</i> 26: 502-512.	3
Lundervold, D. and L. M. Lewin (1990). "Older adults' acceptability of pharmacotherapy and behavior therapy for depression: Initial results." <i>Journal of Applied Gerontology</i> : 211-215.	6
Lynch, T. R., J. S. Cheavens, K. C. Cukrowicz, S. R. Thorp, L. Bronner and J. Beyer (2007). "Treatment of older adults with co-morbid personality disorder and depression: a dialectical behavior therapy approach." <i>Int J Geriatr Psychiatry</i> 22: 131-143.	3
Malt, U. F., O. H. Robak, H. P. Madsbu, O. Bakke and M. Loeb (1999). "The Norwegian naturalistic treatment study of depression in general practice (NORDEP)-I: randomised double blind study." <i>BMJ</i> 318: 1180-1184.	3
Manber, R., A. J. Rush, M. E. Thase, B. Amow, D. Klein, M. H. Trivedi, S. G. Korenstein, J. C. Markowitz, D. L. Dunner, M. Munsaka, F. E. Borian, Martin and B. Keller (2003). "The effects of psychotherapy, nefazodone, and their combination on subjective assessment of disturbed sleep in chronic depression." <i>Sleep</i> 26: 130-136.	3
Manes, F., R. Jorge, M. Morcuende, T. Yamada, S. Paradiso and R. G. Robinson (2001). "A controlled study of repetitive transcranial magnetic stimulation as a treatment of depression in the elderly." <i>Int Psychogeriatr</i> 13(2): 225-231.	3
Martire, L. M., R. Schulz, C. F. Reynolds, J. Q. Morse, M. A. Butters and G. A. Hinrichsen (2008). "Impact of close family members on older adults' early response to depression treatment." <i>Psychol Aging</i> : 447-452.	7
Martire, L. M., R. Schulz, C. F. Reynolds, rd, J. F. Karp, A. G. Gildengers and E. M. Whyte (2010). "Treatment of late-life depression alleviates caregiver burden." <i>J Am Geriatr Soc</i> 58: 23-29.	7
Mavandadi, S., T. R. Ten Have, I. R. Katz, U. N. Durai, D. D. Krahn, M. D. Llorente, J. E. Kirchner, E. J. Olsen, W. W. Van Stone, S. L. Cooley and D. W. Oslin (2007). "Effect of depression treatment on depressive symptoms in older adulthood: the moderating role of pain." <i>J Am Geriatr Soc</i> 55(2): 202-211.	3

McCusker, J., M. Cole, M. Yaffe, P. Cappeliez, M. Dawes, M. Sewitch, A. Ciampi and E. Latimer (2008). "Project direct: Pilot study for a collaborative intervention for depressed seniors." Canadian Journal of Community Mental Health 27: 201-218.	3
McSweeney, K., A. Jeffreys, J. Griffith, C. Plakiotis, R. Kharsas and D. W. O'Connor (2012). "Specialist mental health consultation for depression in Australian aged care residents with dementia: a cluster randomized trial." Int J Geriatr Psychiatry 27: 1163-1171.	4
Mitchell, P. H., L. Teri, R. Veith, A. Buzaitis, D. Tirschwell, K. Becker, M. Fruin, R. Kohen and K. C. Cain (2008). "Living well with stroke: design and methods for a randomized controlled trial of a psychosocial behavioral intervention for poststroke depression." J Stroke Cerebrovasc Dis 17: 109-115.	8
Mohr, D. C., T. Carmody, L. Erickson, L. Jin and J. Leader (2011). "Telephone-administered cognitive behavioral therapy for veterans served by community-based outpatient clinics." J Consult Clin Psychol 79: 261-265.	3
Moscovici, L. and M. Kotler (2009). "A multistage chronobiologic intervention for the treatment of depression: a pilot study." J Affect Disord 116: 201-207.	3
Nadal Rosa, M., A. Losada Salvador, B. Cuixart Carlos, P. Selma Jaume, S. González Sebastián, M. Peláez Irene, M. Fornasini and S. Sallas Jaume (2008). "Ensayo clínico aleatorizado sobre la eficacia de la valoración geriátrica integral seguida de intervención en atención primaria." Rev Esp Geriatr Gerontol 43: 5-12.	3
Naismith, S. L., K. Diamond, P. E. Carter, L. M. Norrie, M. A. Redoblado-Hodge, S. J. G. Lewis and I. B. Hickie (2011). "Enhancing memory in late-life depression: The effects of a combined psychoeducation and cognitive training program." American Journal of Geriatric Psychiatry 19: 240-248.	3
Natarajan, M. K. R. and J. R. Geddes (2008). "Ongoing pharmacotherapy maintains quality of life benefits in older adults with major depression." Evidence Based Mental Health 11: 43-43.	9
Nickel, M. K., C. Nickel, C. Lahmann, F. O. Mitterlehner, K. Tritt, P. K. Leiberich, W. K. Rother and T. H. Loew (2005). "Recovering the ability to function socially in elderly depressed patients: a prospective, controlled trial." Arch Gerontol Geriatr 41: 41-49.	4
Oslin, D. W., R. Thompson, M. J. Kallan, T. TenHave, F. C. Blow, R. Bastani, R. L. Gould, A. E. Maxwell, J. Rosansky, S. Van and L. Jarvik (2004). "Treatment effects from UPBEAT: a randomized trial of care management for behavioral health problems in hospitalized elderly patients." J Geriatr Psychiatry Neurol 17: 99-106.	3
Oxman, T. E. (2003). "Collaborative care may improve depression management in older adults." Evidence Based Mental Health 6(3): 86-86.	9
Parker, J. C., K. L. Smarr, J. R. Slaughter, S. K. Johnston, M. L. Priesmeyer, K. D. Hanson, G. E. Johnson, J. E. Hewett, J. E. Hewett, W. S. Irvin, G. R. Komatireddy and S. E. Walker (2003). "Management of Depression in Rheumatoid Arthritis: A Combined Pharmacologic and Cognitive-Behavioral Approach." Arthritis Care and Research 49: 766-777.	3
Piette, J. D., C. Richardson, J. Himle, S. Duffy, T. Torres, M. Vogel, K. Barber and M. Valenstein (2011). "A randomized trial of telephonic counseling plus walking for depressed diabetes patients." Med Care 49: 641-648.	3
Raj, A. (2004). "PEARLS home based treatment significantly improves depression, dysthymia, and health related quality of life in older people." Evidence Based Mental Health 7: 110-110.	9

Raue, P. J., H. C. Schulberg, M. Heo, S. Klimstra and M. L. Bruce (2009). "Patients' depression treatment preferences and initiation, adherence, and outcome: a randomized primary care study." <i>Psychiatr Serv</i> 60: 337-343.	3
Reynolds, C. F., rd, M. A. Dew, L. M. Martire, M. D. Miller, J. M. Cyranowski, E. Lenze, E. M. Whyte, B. H. Mulsant, B. G. Pollock, J. F. Karp, A. Gildengers, K. Szanto, A. Y. Dombrovski, C. Andreeescu, M. A. Butters, J. Q. Morse, P. R. Houck, S. Bensasi, S. Mazumdar, J. A. Stack and E. Frank (2010). "Treating depression to remission in older adults: a controlled evaluation of combined escitalopram with interpersonal psychotherapy versus escitalopram with depression care management." <i>Int J Geriatr Psychiatry</i> 25: 1134-1141.	7
Reynolds, C. F., rd, E. Frank, J. M. Perel, M. D. Miller, C. Cornes, A. H. Rifai, B. G. Pollock, S. Mazumdar, C. J. George, P. R. Houck and et al. (1994). "Treatment of consecutive episodes of major depression in the elderly." <i>Am J Psychiatry</i> 151: 1740-1743.	7
Reynolds, I., F. C. E. Frank, P. R. Houck, S. Mazumdar, M. A. Dew, C. Cornes, D. J. Buysse, A. Begley and D. J. Kupfer (1997). "Which elderly patients with remitted depression remain well with continued interpersonal psychotherapy after discontinuation of antidepressant medication?" <i>American Journal of Psychiatry</i> 154: 958-962.	7
Rollman, B. L., B. H. Belnap, M. S. LeMenager, S. Mazumdar, P. R. Houck, P. J. Counihan, W. N. Kapoor, H. C. Schulberg, C. F. Reynolds and rd (2009). "Telephone-delivered collaborative care for treating post-CABG depression: a randomized controlled trial." <i>JAMA</i> 302: 2095-2103.	3
Rothblum, E. D., A. J. Sholomskas, C. Berry and B. A. Prusoff (1982). "Issues in clinical trials with the depressed elderly." <i>J Am Geriatr Soc</i> 30: 694-699.	6
Schaik, D. J. F., E. Dozeman, H. W. J. Marwijk, M. L. Stek, F. Smit, A. T. F. Beekman and H. E. Horst (2014) "Preventing depression in homes for older adults: Are effects sustained over 2 years?" <i>International journal of geriatric psychiatry</i> 29, 191-197 DOI: 10.1002/gps.3989.	3
Schapira, M., L. Camera, C. Finkelsztein, D. Matusevich, M. Smietniansky, J. Esteban, H. Patino and A. Acuna (2010). "[A multidisciplinary program for the treatment and follow up of depression in ambulatory elderly]." <i>Vertex</i> 21: 284-290.	11
Schneider, L. S., E. R. Fredrickson, J. A. Severson and R. B. Sloane (1986) "3H-Imipramine binding in depressed elderly: relationship to family history and clinical response." <i>Psychiatry Res</i> , 257-266.	6
Shah, A., K. Odutoye and T. De (2001). "Depression in acutely medically ill elderly inpatients: a pilot study of early identification and intervention by formal psychogeriatric consultation." <i>J Affect Disord</i> 62(3): 233-240.	4
Silveira, H., A. C. Deslandes, H. Moraes, R. Mouta, P. Ribeiro, R. Piedade and J. Laks (2010). "Effects of exercise on electroencephalographic mean frequency in depressed elderly subjects." <i>Neuropsychobiology</i> : 141-147.	7
Sirey, J. A., M. L. Bruce and G. S. Alexopoulos (2005). "The Treatment Initiation Program: an intervention to improve depression outcomes in older adults." <i>Am J Psychiatry</i> 162: 184-186.	3
Sirey, J. A., M. L. Bruce and H. C. Kales (2010). "Improving antidepressant adherence and depression outcomes in primary care: the treatment initiation and participation (TIP) program." <i>Am J Geriatr Psychiatry</i> 18: 554-562.	4
Slusher, L. W. (2007). "A care management intervention improved depression after stroke." <i>Evidence Based Nursing</i> 10(4): 117-117.	9

Smit, A., B. G. Tiemens, J. Ormel, H. Kluiter, J. A. Jenner, M. van de, O. van, W. D. P. T and H. J. Conradi (2005). "Enhanced treatment for depression in primary care: First year results on compliance, self-efficacy, the use of antidepressants and contacts with the primary care physician." Primary Care & Community Psychiatry 10: 39-49.	3
Stoudemire, A., C. D. Hill and R. Morris (1993). "Long-term affective cognitive outcome in depressed older adults." American Journal of Psychiatry: 896-900.	3
Szanto, K., B. H. Mulsant, P. Houck, M. A. Dew, C. F. Reynolds and rd (2003). "Occurrence and course of suicidality during short-term treatment of late-life depression." Arch Gen Psychiatry 60: 610-617.	9
Unutzer, J., W. Katon, C. M. Callahan, J. W. Williams, Jr., E. Hunkeler, L. Harpole, M. Hoffing, R. D. Della Penna, P. H. Noel, E. H. Lin, P. A. Arean, M. T. Hegel, L. Tang, T. R. Belin, S. Oishi, C. Langston and I. I. I. M.-P. A. t. C. Treatment (2002). "Collaborative care management of late-life depression in the primary care setting: a randomized controlled trial." JAMA 288(22): 2836-2845.	3
Unutzer, J., L. Tang, S. Oishi, W. Katon, J. W. Williams, Jr, E. Hunkeler, H. Hendrie, E. H. Lin, S. Levine, L. Grypma, D. C. Steffens, J. Fields, C. Langston and I. Investigators (2006). "Reducing suicidal ideation in depressed older primary care patients." J Am Geriatr Soc 54: 1550-1556.	3
Unützer, J., W. Katon, C. M. Callahan, J. W. Williams, Jr., E. Hunkeler, L. Harpole, M. Hoffing, R. D. Della Penna, P. H. Noel, E. H. B. Lin, L. Tang and S. Oishi (2003). "Depression treatment in a sample of 1,801 depressed older adults in primary care." J Am Geriatr Soc 51(4): 505-514.	7
van der Aa, H. P., G. H. van Rens, H. C. Comijs, J. E. Bosmans, T. H. Margrain and R. M. van Nispen (2013). "Stepped-care to prevent depression and anxiety in visually impaired older adults--design of a randomised controlled trial." BMC Psychiatry 13: 209.	9
Van, H. L., J. Dekker, J. Peen, A. van and R. A. Schoevers (2008). "Identifying patients at risk of complete nonresponse in the outpatient treatment of depression." Psychother Psychosom 77: 358-364.	3
Van Leeuwen Williams, E., J. Unutzer, S. Lee and P. H. Noel (2009). "Collaborative depression care for the old-old: findings from the IMPACT trial." Am J Geriatr Psychiatry 17(12): 1040-1049.	7
van, M., W. H. H. Ader, H. de and A. Beekman (2008). "Primary care management of major depression in patients aged > or =55 years: outcome of a randomised clinical trial." Br J Gen Pract 58: 680-686, I.	3
Van Orden, K. A., A. E. Begley, Y. Conwell and I. C. F. Reynolds (2014). "Does reducing social disconnectedness represent a mechanism whereby depression treatment reduces suicide risk?" American Journal of Geriatric Psychiatry 22(3): S134-S135.	3
Vannoy, S. D., P. Duberstein, K. Cukrowicz, E. Lin, M. Y. Fan and J. Unutzer (2007). "The relationship between suicide ideation and late-life depression." Am J Geriatr Psychiatry 15: 1024-1033.	7
Van't Veer-Tazelaar, P., F. Smit, H. van Hout, P. van Oppen, H. van der Horst, A. Beekman and H. van Marwijk (2010). "Cost-effectiveness of a stepped care intervention to prevent depression and anxiety in late life: randomised trial." Br J Psychiatry 196(4): 319-325.	6
Warner, B. (2010). "Reducing suicidal ideation and depression in older primary care patients: The oldest old and pain." American Journal of Psychiatry 167: 102-103.	9

Verkaik, R., A. L. Francke, M. van, P. M. M. Spreeuwenberg, M. W. Ribbe and J. M. Bensing (2011). "The introduction of a nursing guideline on depression at psychogeriatric nursing home wards: Effects on Certified Nurse Assistants." <i>Int J Nurs Stud</i> 48: 710-719.	4
Williams, J. W. and Jr (2008). "Managed depression care reduces mortality in older adults with major depression." <i>Evidence Based Mental Health</i> 11: 16-16.	9
Williams, J. W., Jr, W. Katon, E. H. Lin, P. H. Noel, J. Worchel, J. Cornell, L. Harpole, B. A. Fultz, E. Hunkeler, V. S. Mika, J. Unutzer and I. Investigators (2004). "The effectiveness of depression care management on diabetes-related outcomes in older patients." <i>Ann Intern Med</i> 140: 1015-1024.	7
Wolfson, L., M. Miller, P. Houck, L. Ehrenpreis, J. A. Stack, E. Frank, C. Cornes, S. Mazumdar, D. J. Kupfer and C. F. Reynolds (1997). "Foci of interpersonal psychotherapy (IPT) in depressed elders: Clinical and outcome correlates in a combined IPT/nortriptyline protocol." <i>Psychotherapy Research</i> : 45-56.	9
Von, K., W. J. Katon, E. H. B. Lin, P. Ciechanowski, D. Peterson, E. J. Ludman, B. Young and C. M. Rutter (2011). "Functional outcomes of multi-condition collaborative care and successful ageing: Results of randomised trial." <i>BMJ (Online)</i> 343: 1083.	3
Xia, Q., J. Wang and K. Zheng (2006). "Rehabilitative therapy for the elderly chronic somatic disease complicated with depression." <i>Chinese Journal of Clinical Rehabilitation</i> 10: 166-168.	4
Xu, S., K. Rost, F. Dong and L. M. Dickinson (2011). "Stakeholder benefit from depression disease management: differences by rurality?" <i>J Behav Health Serv Res</i> 38: 114-121.	6
Zeng, W., H. Ma, Q. Liang, Y. Dong, H. Ye and Y. Zhang (2001). "[The influence of antidepressive therapy on short-term prognosis in elderly patients with unstable angina and depression]." <i>Zhonghua Nei Ke Za Zhi</i> 40: 809-810.	11
Zhang, H. W., C. Y. Wang, H. N. Xu, X. Zhao, Q. Dai, J. Li, X. Du, Z. Song, G. Han, G. Liu, P. Li and H. Lin (2006). "[Clinical study on effect of fluoxetine combined with Chinese medicine or tibetan drugs in treating senile depression in plateau district]." <i>Zhongguo Zhong Xi Yi Jie He Za Zhi</i> 26(3): 202-204.	11
Zou, X. B., Z. X. Lin, J. D. Lin, D. Lu and G. M. Chen (2005). "Interventional efficacy of citalopram combined with shining and psychological morning exercise in the attack of depression in elderly people." <i>Chinese Journal of Clinical Rehabilitation</i> 9: 24-25.	11

Ljusterapi

Avery, D. and G. Winokur (1976). "Mortality in depressed patients treated with electroconvulsive therapy and antidepressants." <i>Arch Gen Psychiatry</i> 33: 1029-1037.	3
Blumberger, D. M., B. H. Mulsant, P. B. Fitzgerald, T. K. Rajji, A. V. Ravindran, L. T. Young, A. J. Levinson and Z. J. Daskalakis (2012). "A randomized double-blind sham-controlled comparison of unilateral and bilateral repetitive transcranial magnetic stimulation for treatment-resistant major depression." <i>World J Biol Psychiatry</i> 13: 423-435.	3
Brunoni, A. R., R. Ferrucci, M. Bortolomasi, M. Vergari, L. Tadini, P. S. Boggio, M. Giacopuzzi, S. Barbieri and A. Priori (2011). "Transcranial direct current stimulation (tDCS) in unipolar vs. bipolar depressive disorder." <i>Prog Neuropsychopharmacol Biol Psychiatry</i> 35: 96-101.	3
Cronholm, B. and J. O. Ottosson (1996). "Experimental studies of the therapeutic action of electroconvulsive therapy in endogenous depression. The role of the electrical stimulation and of the seizure studied by variation of stimulus intensity and modification by lidocaine of seizure discharge." <i>Convuls Ther</i> 12: 172-194.	3
Devanand, D. P., S. Lisanby, E. S. Lo, L. Fitzsimons, T. B. Cooper, U. Halbreich and H. A. Sackeim (1998). "Effects of electroconvulsive therapy on plasma vasopressin and oxytocin." <i>Biol Psychiatry</i> 44: 610-616.	3
Dowling, G. A., C. L. Graf, E. M. Hubbard and J. S. Luxenberg (2007). "Light treatment for neuropsychiatric behaviors in Alzheimer's disease." <i>West J Nurs Res</i> 29: 961-975.	3
Fitzgerald, P. B., K. Hoy, R. Gunewardene, C. Slack, S. Ibrahim, M. Bailey and Z. J. Daskalakis (2011). "A randomized trial of unilateral and bilateral prefrontal cortex transcranial magnetic stimulation in treatment-resistant major depression." <i>Psychol Med</i> 41: 1187-1196.	3
Flint, A. J. and S. L. Rifat (1998). "The treatment of psychotic depression in later life: a comparison of pharmacotherapy and ECT." <i>Int J Geriatr Psychiatry</i> 13: 23-28.	3
Folkerts, H. W., N. Michael, R. Tolle, K. Schonauer, S. Mucke and H. Schulze-Monking (1997). "Electroconvulsive therapy vs. paroxetine in treatment-resistant depression -- a randomized study." <i>Acta Psychiatr Scand</i> 96: 334-342.	3
Genhart, M. J., K. A. Kelly, R. D. Coursey, M. Datiles and N. E. Rosenthal (1993). "Effects of bright light on mood in normal elderly women." <i>Psychiatry Res</i> 47: 87-97.	3
Hansen, P. E., B. Ravnkilde, P. Videbech, K. Clemmensen, R. Sturlason, M. Reiner, E. Parner, R. Rosenberg and P. Vestergaard (2011). "Low-frequency repetitive transcranial magnetic stimulation inferior to electroconvulsive therapy in treating depression." <i>J ECT</i> 27: 26-32.	3
Hickman, S. E., A. L. Barrick, C. S. Williams, S. Zimmerman, B. R. Connell, J. S. Preisser, C. M. Mitchell and P. D. Sloane (2007). "The effect of ambient bright light therapy on depressive symptoms in persons with dementia." <i>J Am Geriatr Soc</i> 55(11): 1817-1824.	3
Husain, M. M., S. M. McClintonck, A. J. Rush, R. G. Knapp, M. Fink, T. A. Rummans, K. Rasmussen, C. Claassen, G. Petrides, M. M. Biggs, M. Mueller, S. Sampson, S. H. Bailine, S. H. Lisanby and C. H. Kellner (2008). "The efficacy of acute electroconvulsive therapy in atypical depression." <i>J Clin Psychiatry</i> 69: 406-411.	3

Husain, M. M., A. J. Rush, M. Fink, R. Knapp, G. Petrides, T. Rummans, M. M. Biggs, K. O'Connor, K. Rasmussen, M. Little, W. Zhao, H. J. Bernstein, G. Smith, M. Mueller, S. M. McClintock, S. H. Bailine and C. H. Kellner (2004). "Speed of response and remission in major depressive disorder with acute electroconvulsive therapy (ECT): a Consortium for Research in ECT (CORE) report." <i>J Clin Psychiatry</i> 65: 485-491.	3
Janicak, P. G., J. P. O'Reardon, S. M. Sampson, M. M. Husain, S. H. Lisanby, J. T. Rado, K. L. Heart and M. A. Demitrack (2008). "Transcranial magnetic stimulation in the treatment of major depressive disorder: a comprehensive summary of safety experience from acute exposure, extended exposure, and during reintroduction treatment." <i>J Clin Psychiatry</i> 69: 222-232.	3
Janicak, P. G., R. P. Sharma, T. H. Israni, S. M. Dowd, E. Altman and J. M. Davis (1991). "Effects of unilateral-nondominant vs. bilateral ECT on memory and depression: a preliminary report." <i>Psychopharmacol Bull</i> 27: 353-357.	3
Jorge, R. E., D. J. Moser, L. Acion and R. G. Robinson (2008). "Treatment of vascular depression using repetitive transcranial magnetic stimulation." <i>Arch Gen Psychiatry</i> 65: 268-276.	3
Kellner, C. H., M. Fink, R. Knapp, G. Petrides, M. Husain, T. Rummans, M. Mueller, H. Bernstein, K. Rasmussen, K. O'Connor, G. Smith, A. J. Rush, M. Biggs, S. McClintock, S. Bailine and C. Malur (2005). "Relief of expressed suicidal intent by ECT: a consortium for research in ECT study." <i>American Journal of Psychiatry</i> 162: 977-982.	3
Kellner, C. H., R. Knapp, M. M. Husain, K. Rasmussen, S. Sampson, M. Cullum, S. M. McClintock, K. G. Tobias, C. Martino, M. Mueller, S. H. Bailine, M. Fink and G. Petrides (2010). "Bifrontal, bitemporal and right unilateral electrode placement in ECT: randomised trial." <i>Br J Psychiatry</i> 196(3): 226-234.	3
Kellner, C. H., Monroe, Jr., R. R. J. Pritchett, M. P. Jarrell, H. J. Bernstein and C. M. Burns (1992). "Weekly ECT in geriatric depression." <i>Convuls Ther</i> 8: 245-252.	3
Kripke, D. F. (2011). "Bright light treatment reduces symptoms in older adults with non-seasonal major depression." <i>Evidence Based Mental Health</i> 14(3): 75-75.	3
Krystal, A. D., T. Holsinger, R. D. Weiner and C. E. Coffey (2000). "Prediction of the utility of a switch from unilateral to bilateral ECT in the elderly using treatment 2 ictal EEG indices." <i>J ECT</i> 16: 327-337.	3
Lieverse, R., M. Nielen, B. Uitdehaag, E. Van Someren, J. Smit and W. Hoogendoijk (2009). "Double blind randomised clinical trial of bright light therapy in elderly subjects with nonseasonal major depressive disorder." <i>European Psychiatry</i> 24: S320.	3
Lieverse, R., M. M. Nielen, D. J. Veltman, B. M. Uitdehaag, E. J. van Someren, J. H. Smit and W. J. Hoogendoijk (2008). "Bright light in elderly subjects with nonseasonal major depressive disorder: a double blind randomised clinical trial using early morning bright blue light comparing dim red light treatment." <i>Trials</i> 9: 48.	3
Loving, R. T., D. F. Kripke, J. A. Elliott, N. C. Knickerbocker and M. A. Grandner (2005). "Bright light treatment of depression for older adults [ISRCTN55452501]." <i>BMC Psychiatry</i> 5: 41.	3
Loving, R. T., D. F. Kripke, N. C. Knickerbocker and M. A. Grandner (2005). "Bright green light treatment of depression for older adults [ISRCTN69400161]." <i>BMC Psychiatry</i> 5: 42.	3

Lyness, J. M. (2004). "Treatment of depressive conditions in later life: real-world light for dark (or dim) tunnels." <i>JAMA: Journal of the American Medical Association</i> 291: 1626-1628.	9
Martiny, K. (2011). "Bright light treatment is effective in treating older patients with non-seasonal major depression." <i>Evidence Based Nursing</i> 14(4): 117-118.	3
McCall, W. V., C. C. Colenda and B. A. Farah (1996). "Ictal EEG regularity declines during a course of RUL ECT." <i>Convuls Ther</i> 12: 213-216.	3
McCall, W. V., A. Dunn, P. B. Rosenquist and D. Hughes (2002). "Markedly suprathreshold right unilateral ECT versus minimally suprathreshold bilateral ECT: antidepressant and memory effects." <i>J ECT</i> 18: 126-129.	3
McCall, W. V., B. A. Farah, D. Reboussin and C. C. Colenda (1995). "Comparison of the efficacy of titrated, moderate-dose and fixed, high- dose right unilateral ECT in elderly patients." <i>American Journal of Geriatric Psychiatry</i> 3(4): 317-324.	3
Moffit, M. T. and S. Ancoli-Israel (1993) "Bright light treatment of late-life depression." <i>Society of Light Treatment and Biological Rhythms Abstracts</i> , 41.	3
Mosimann, U. P., W. Schmitt, B. D. Greenberg, M. Kosel, R. M. Muri, M. Berkhoff, C. W. Hess, H. U. Fisch and T. E. Schlaepfer (2004). "Repetitive transcranial magnetic stimulation: a putative add-on treatment for major depression in elderly patients." <i>Psychiatry Res</i> 126: 123-133.	3
Most, E. I. S., P. Scheltens, S. Van and J. W. E (2012). "Prevention of depression in elderly with memory-problems by activation of the biological clock with light." <i>Journal of Sleep Research</i> 21: 34-35.	3
O'Connor Daniel, W., B. Gardner, I. Presnell, D. Singh, M. Tsanglis and E. White (2010). "The effectiveness of continuation-maintenance ECT in reducing depressed older patients' hospital re-admissions." <i>J Affect Disord</i> 120: 62-66.	7
O'Leary, D., D. Gill, S. Gregory and C. Shawcross (1994). "The effectiveness of real versus simulated electroconvulsive therapy in depressed elderly patients." <i>Int J Geriatr Psychiatry</i> : 567-571.	3
O'Reardon, J. P., H. B. Solvason, P. G. Janicak, S. Sampson, K. E. Isenberg, Z. Nahas, W. M. McDonald, D. Avery, P. B. Fitzgerald, C. Loo, M. A. Demitrack, M. S. George and H. A. Sackeim (2007). "Efficacy and safety of transcranial magnetic stimulation in the acute treatment of major depression: a multisite randomized controlled trial." <i>Biol Psychiatry</i> 62: 1208-1216.	3
Petrides, G., M. Fink, M. M. Husain, R. G. Knapp, A. J. Rush, M. Mueller, T. A. Rummans, K. M. O'Connor, K. G. Rasmussen, Jr., H. J. Bernstein, M. Biggs, S. H. Bailine and C. H. Kellner (2001). "ECT remission rates in psychotic versus nonpsychotic depressed patients: a report from CORE." <i>J ECT</i> 17(4): 244-253.	3
Quante, A., A. Luborzewski, E. L. Brakemeier, A. Merkl, H. Danker-Hopfe and M. Bajbouj (2011). "Effects of 3 different stimulus intensities of ultrabrief stimuli in right unilateral electroconvulsive therapy in major depression: a randomized, double-blind pilot study." <i>J Psychiatr Res</i> 45: 174-178.	3

Riemersma-van der Lek, R. F., D. F. Swaab, J. Twisk, E. M. Hol, W. J. Hoogendoijk and E. J. Van Someren (2008). "Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities: a randomized controlled trial." <i>JAMA</i> 299(22): 2642-2655.	3
Royer, M., N. H. Ballantine, P. J. Eslinger, K. Houser, R. Mistrick, R. Behr and K. Rakos (2012). "Light therapy for seniors in long term care." <i>J Am Med Dir Assoc</i> 13: 100-102.	3
Sackeim, H. A., J. Prudic, D. P. Devanand, M. S. Nobler, S. H. Lisanby, S. Peyser, L. Fitzsimons, B. J. Moody and J. Clark (2000). "A prospective, randomized, double-blind comparison of bilateral and right unilateral electroconvulsive therapy at different stimulus intensities." <i>Arch Gen Psychiatry</i> 57: 425-434.	3
Sackeim, H. A., J. Prudic, M. S. Nobler, L. Fitzsimons, S. H. Lisanby, N. Payne, R. M. Berman, E. L. Brakemeier, T. Perera and D. P. Devanand (2008). "Effects of pulse width and electrode placement on the efficacy and cognitive effects of electroconvulsive therapy." <i>Brain Stimul</i> 1: 71-83.	7
Sackeim, H. A., A. J. Rush, M. S. George, L. B. Marangell, M. M. Husain, Z. Nahas, C. R. Johnson, S. Seidman, C. Giller, S. Haines, Simpson, Jr., K. R and R. R. Goodman (2001). "Vagus nerve stimulation (VNS(trademark)) for treatment-resistant depression: Efficacy, side effects, and predictors of outcome." <i>Neuropsychopharmacology</i> 25: 713-728.	3
Segman, R. H., B. Shapira, M. Gorfine and B. Lerer (1995). "Onset and time course of antidepressant action: psychopharmacological implications of a controlled trial of electroconvulsive therapy." <i>Psychopharmacology (Berl)</i> 119: 440-448.	3
Serra, M., C. Gasto, V. Navarro, X. Torres, J. Blanch and G. Masana (2006). "[Maintenance electroconvulsive therapy in elderly psychotic unipolar depression]." <i>Med Clin (Barc)</i> 126: 491-492.	11
Shapira, B., N. Tubi and B. Lerer (2000). "Balancing speed of response to ECT in major depression and adverse cognitive effects: role of treatment schedule." <i>J ECT</i> 16(2): 97-109.	3
Sherman, F. T. (2009). "Life-saving treatment for depression in elderly: always think of electroconvulsive therapy (ECT)." <i>Geriatrics</i> 64: 8.	9
Stek, M. L., F. B. van der Wurff, B. M. Uitdehaag, A. T. Beekman and W. J. Hoogendoijk (2007). "ECT in the treatment of depressed elderly: lessons from a terminated clinical trial." <i>Int J Geriatr Psychiatry</i> 22(10): 1052-1054.	3
Stoppe, A., M. Louza, M. Rosa, G. Gil and S. Rigonatti (2006). "Fixed high-dose electroconvulsive therapy in the elderly with depression: a double-blind, randomized comparison of efficacy and tolerability between unilateral and bilateral electrode placement." <i>J ECT</i> 22: 92-99.	3
Stoudemire, A., G. Knos, M. Gladson, H. Markwalter, Y. F. Sung, R. Morris and R. Cooper (1990). "Labetalol in the control of cardiovascular responses to electroconvulsive therapy in high-risk depressed medical patients." <i>J Clin Psychiatry</i> 51: 508-512.	3
Sumaya, I. C., B. M. Rienzi, J. F. Deegan, 2nd and D. E. Moss (2001). "Bright light treatment decreases depression in institutionalized older adults: a placebo-controlled crossover study." <i>J Gerontol A Biol Sci Med Sci</i> 56(6): M356-360.	3

Tew, J. D., Jr, B. H. Mulsant, R. F. Haskett, D. Dolata, L. Hixson and J. J. Mann (2002). "A randomized comparison of high-charge right unilateral electroconvulsive therapy and bilateral electroconvulsive therapy in older depressed patients who failed to respond to 5 to 8 moderate-charge right unilateral treatments." <i>J Clin Psychiatry</i> 63: 1102-1105.	3
Tew, J. D., Jr., B. H. Mulsant, R. F. Haskett, J. Prudic, M. E. Thase, R. R. Crowe, D. Dolata, A. E. Begley, C. F. Reynolds, 3rd and H. A. Sackeim (1999). "Acute efficacy of ECT in the treatment of major depression in the old-old." <i>Am J Psychiatry</i> 156(12): 1865-1870.	7
Tsai, Y. F., T. K. Wong, Y. Y. Juang and H. H. Tsai (2004). "The effects of light therapy on depressed elders." <i>Int J Geriatr Psychiatry</i> 19: 545-548.	3
Zis, A. P., L. N. Yatham, R. W. Lam, C. M. Clark, M. Srisurapanont and K. McGarvey (1996). "Effect of stimulus intensity on prolactin and cortisol release induced by unilateral electroconvulsive therapy." <i>Neuropsychopharmacology</i> 15: 263-270.	3

Hälsoekonomi

Araya, R., T. Flynn, G. Rojas, R. Fritsch and G. Simon (2006). "Cost-effectiveness of a primary care treatment program for depression in low-income women in Santiago, Chile." <i>Am J Psychiatry</i> 163(8): 1379-1387.	3
Aziz, M., A. M. Mehringer, E. Mozurkewich and G. N. Razik (2005). "Cost-utility of 2 maintenance treatments for older adults with depression who responded to a course of electroconvulsive therapy: results from a decision analytic model." <i>Can J Psychiatry</i> 50(7): 389-397.	6
Borghi, J. and J. F. Guest (2000). "Economic impact of using mirtazapine compared to amitriptyline and fluoxetine in the treatment of moderate and severe depression in the UK." <i>Eur Psychiatry</i> 15(6): 378-387.	3
Bosmans JE, Dozeman E, van Marwijk HWJ, van Schaik DJF, Stek ML, Beekman ATF, et al. Cost-effectiveness of a stepped care programme to prevent depression and anxiety in residents in homes for the older people: A randomised controlled trial. <i>International Journal of Geriatric Psychiatry</i> 2014;29:182-190.	3
Bosmans, J. E., B. Schreuders, H. W. van Marwijk, J. H. Smit, P. van Oppen and M. W. van Tulder (2012). "Cost-effectiveness of problem-solving treatment in comparison with usual care for primary care patients with mental health problems: a randomized trial." <i>BMC Fam Pract</i> 13: 98.	3
Bosmans, J. E., D. J. van Schaik, M. W. Heymans, H. W. van Marwijk, H. P. van Hout and M. C. de Bruijne (2007). "Cost-effectiveness of interpersonal psychotherapy for elderly primary care patients with major depression." <i>Int J Technol Assess Health Care</i> 23(4): 480-487.	3
Bosmans, J., M. de Bruijne, H. van Hout, H. van Marwijk, A. Beekman, L. Bouter, W. Stalman and M. van Tulder (2006). "Cost-effectiveness of a disease management program for major depression in elderly primary care patients." <i>J Gen Intern Med</i> 21(10): 1020-1026.	3
Boyer, P., J. M. Danion, J. C. Bisserbe, J. M. Hotton and S. Troy (1998). "Clinical and economic comparison of sertraline and fluoxetine in the treatment of depression. A 6-month double-blind study in a primary-care setting in France." <i>Pharmacoeconomics</i> 13(1 Pt 2): 157-169.	3
Brettschneider C, Konig HH. Cost-utility analyses of cognitive-behavioral therapy for major depressive disorder - A systematic review. <i>Value in Health</i> , 2013; 16 (7): A549.	9
Brown, M. C. J., A. A. Nimmerrichter and J. F. Guest (1999). "Cost-effectiveness of mirtazapine compared to amitriptyline and fluoxetine in the treatment of moderate and severe depression in Austria." <i>European Psychiatry</i> 14(4): 230-244.	3
Browne, G., M. Steiner, J. Roberts, A. Gafni, C. Byrne, E. Dunn, B. Bell, M. Mills, L. Chalklin, D. Wallik and J. Kraemer (2002). "Sertraline and/or interpersonal psychotherapy for patients with dysthymic disorder in primary care: 6-month comparison with longitudinal 2-year follow-up of effectiveness and costs." <i>J Affect Disord</i> 68(2-3): 317-330.	3
Chisholm, D., K. Sanderson, J. L. Ayuso-Mateos and S. Saxena (2004). "Reducing the global burden of depression: population-level analysis of intervention cost-effectiveness in 14 world regions." <i>Br J Psychiatry</i> 184: 393-403.	3
Dezetter A, Briffault X, Ben Lakhdar C, Kovess-Masfety V. Costs and benefits of improving access to psychotherapies for common mental disorders. <i>J Ment Health Policy Econ</i> 2013;16:161-77.	3

Fantino, B., N. Moore, H. Verdoux and J.-P. Auray (2007). "Cost-effectiveness of escitalopram vs. citalopram in major depressive disorder." <i>Int Clin Psychopharmacol</i> 22(2): 107-115.	3
Gensichen, J., J. J. Petersen, M. Von Korff, D. Heider, S. Baron, J. Konig, A. Freytag, C. Krauth, F. M. Gerlach and H. H. Konig (2013). "Cost-effectiveness of depression case management in small practices." <i>British Journal of Psychiatry</i> 202(6): 441-446.	3
Gerhards, S. A., L. E. de Graaf, L. E. Jacobs, J. L. Severens, M. J. Huibers, A. Arntz, H. Riper, G. Widdershoven, J. F. Metsemakers and S. M. Evers (2010). "Economic evaluation of online computerised cognitive-behavioural therapy without support for depression in primary care: randomised trial." <i>Br J Psychiatry</i> 196(4): 310-318.	3
Gusi, N., M. C. Reyes, J. L. Gonzalez-Guerrero, E. Herrera and J. M. Garcia (2008). "Cost-utility of a walking programme for moderately depressed, obese, or overweight elderly women in primary care: a randomised controlled trial." <i>BMC Public Health</i> 8: 231.	3
Haji EO, Mann K, Dragicevic A, Muller MJ, Boland K, Rao ML, et al. Potential cost-effectiveness of therapeutic drug monitoring for depressed patients treated with citalopram. <i>Therapeutic Drug Monitoring</i> 2013;35:396-401.	3
Holland J, Bhogle M. Sertraline and mirtazapine as geriatric antidepressants. <i>Psychiatr Danub</i> 2013;25 Suppl 2:S286-90.	6
Hollinghurst S, Carroll FE, Abel A, Campbell J, Garland A, Jerrom B, et al. Cost-effectiveness of cognitive-behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: economic evaluation of the CoBalT Trial. <i>The British Journal of Psychiatry</i> 2014;204:69-76.	3
Hollinghurst, S., T. J. Peters, S. Kaur, N. Wiles, G. Lewis and D. Kessler (2010). "Cost-effectiveness of therapist-delivered online cognitive-behavioural therapy for depression: randomised controlled trial." <i>Br J Psychiatry</i> 197(4): 297-304.	3
Joling KJ, Bosmans JE, van Marwijk HW, van der Horst HE, Scheltens P, Vroomen JL, et al. The cost-effectiveness of a family meetings intervention to prevent depression and anxiety in family caregivers of patients with dementia: a randomized trial. <i>Trials</i> 2013;14:305.	3
Jonkers, C. C., F. Lamers, S. M. Evers, H. Bosma and J. T. Van Eijk (2010). "Cost-utility estimates in depression: does the valuation method matter?" <i>J Ment Health Policy Econ</i> 13(4): 189-197.	4
Jonkers, C. C., F. Lamers, S. M. Evers, H. Bosma, J. F. Metsemakers and J. T. Van Eijk (2009). "Economic evaluation of a minimal psychological intervention in chronically ill elderly patients with minor or mild to moderate depression: a randomized trial (the DELTA-study)." <i>Int J Technol Assess Health Care</i> 25(4): 497-504.	3
Katon, W., J. Unutzer, M. Y. Fan, J. W. Williams, Jr., M. Schoenbaum, E. H. Lin and E. M. Hunkeler (2006). "Cost-effectiveness and net benefit of enhanced treatment of depression for older adults with diabetes and depression." <i>Diabetes Care</i> 29(2): 265-270.	3
Kendrick, T., R. Peveler, L. Longworth, D. Baldwin, M. Moore, J. Chatwin, A. Thornett, J. Goddard, M. Campbell, H. Smith, M. Buxton and C. Thompson (2006). "Cost-effectiveness and cost-utility of tricyclic antidepressants, selective serotonin reuptake inhibitors and lofepramine: Randomised controlled trial." <i>British Journal of Psychiatry</i> 188(APR.): 337-345.	3
Klug, G., G. Hermann, B. Fuchs-Nieder, M. Panzer, A. Haider-Stipacek, H. G. Zapotoczky and S. Priebe (2010). "Effectiveness of home treatment for elderly people with depression: randomised controlled trial." <i>Br J Psychiatry</i> 197(6): 463-467.	4

Klug, G., G. Hermann, B. Fuchs-Nieder, M. Panzer, A. Haider-Stipacek, H. G. Zapotoczky and S. Priebe (2011). "Clinically and economically effectiveness of home treatment for elderly patients with depression: Randomised controlled trial." <i>Journal of Mental Health Policy and Economics</i> 14: S15.	3
Knapp M, King D, Romeo R, Schehl B, Barber J, Griffin M, et al. Cost effectiveness of a manual based coping strategy programme in promoting the mental health of family carers of people with dementia (the START (STrAtegies for RelaTives) study): a pragmatic randomised controlled trial. <i>Bmj</i> 2013;347:f6342.	3
Kono, A., Y. Kanaya, T. Fujita, C. Tsumura, T. Kondo, K. Kushiyama and L. Z. Rubenstein (2012) "Effects of a preventive home visit program in ambulatory frail older people: a randomized controlled trial (Provisional abstract)." <i>Journals of Gerontology Series A - Biological Sciences and Medical Sciences</i> , 302-309.	3
Ladapo, J. A., J. A. Shaffer, Y. Fang, S. Ye and K. W. Davidson (2012). "Cost-effectiveness of enhanced depression care after acute coronary syndrome: results from the Coronary Psychosocial Evaluation Studies randomized controlled trial." <i>Arch Intern Med</i> 172(21): 1682-1684.	9
Lapierre, Y., J. Bentkover, S. Schainbaum and S. Manners (1995). "Direct cost of depression: analysis of treatment costs of paroxetine versus Imipramine in Canada." <i>Can J Psychiatry</i> 40(7): 370-377.	3
Liu, C.-F., S. C. Hedrick, E. F. Chaney, P. Heagerty, B. Felker, N. Hasenberg, S. Fihn and W. Katon (2003). "Cost-effectiveness of collaborative care for depression in a primary care veteran population." <i>Psychiatric Services</i> 54(5): 698-704.	3
Lokkerbol J, Adema D, Cuijpers P, Reynolds ICF, Schulz R, Weehuizen R, et al. Improving the cost-effectiveness of a healthcare system for depressive disorders by implementing telemedicine: A health economic modeling study. <i>American Journal of Geriatric Psychiatry</i> 2014;22:253-262.	3
McCrone, P., M. Knapp, J. Proudfoot, C. Ryden, K. Cavanagh, D. A. Shapiro, S. Ilson, J. A. Gray, D. Goldberg, A. Mann, I. Marks, B. Everitt and A. Tylee (2004). "Cost-effectiveness of computerised cognitive-behavioural therapy for anxiety and depression in primary care: randomised controlled trial." <i>Br J Psychiatry</i> 185: 55-62.	3
McCusker, J., P. Jacobs, N. Dendukuri, E. Latimer, P. Tousignant and J. Verdon (2003) "Cost-effectiveness of a brief two-stage emergency department intervention for high-risk elders: results of a quasi-randomized controlled trial (Structured abstract)." <i>Annals of Emergency Medicine</i> , 45-56.	3
McDonald, W. M., V. L. Phillips, G. S. Figiel, F. A. Marsteller, C. D. Simpson and M. C. Bailey (1998). "Cost-effective maintenance treatment of resistant geriatric depression." <i>Psychiatric Annals</i> 28(1): 47-52.	3
Meyers, B. S. (2007). "Optimizing the use of data generated by geriatric depression treatment studies during a time of diminishing resources." <i>Am J Geriatr Psychiatry</i> 15(7): 545-552.	9
Olgiati P, Bajo E, Bigelli M, Montgomery S, Serretti A. Challenging sequential approach to treatment resistant depression: Cost-utility analysis based on the Sequenced Treatment Alternatives to Relieve Depression (STAR-D) trial. <i>European Neuropsychopharmacology</i> 2013;23:1739-1746.	3

Oligati, P., E. Bajo, M. Bigelli, D. De Ronchi and A. Serretti (2012). "Should pharmacogenetics be incorporated in major depression treatment? Economic evaluation in high- and middle-income European countries." <i>Prog Neuropsychopharmacol Biol Psychiatry</i> 36(1): 147-154.	3
Parker, S. G., P. Oliver, M. Pennington, J. Bond, C. Jagger, P. M. Enderby, R. Curless, T. Chater, A. Vanoli, K. Fryer, C. Cooper, S. Julious, C. Donaldson, C. Dyer, T. Wynn, A. John and D. Ross (2009) "Rehabilitation of older patients: day hospital compared with rehabilitation at home - a randomised controlled trial (Provisional abstract)." <i>Health Technol Assess</i> , i-xi.	4
Pyne, J. M., J. C. Fortney, S. P. Tripathi, M. L. Maciejewski, M. J. Edlund and D. K. Williams (2010). "Cost-effectiveness analysis of a rural telemedicine collaborative care intervention for depression." <i>Arch Gen Psychiatry</i> 67(8): 812-821.	3
Reynolds, C. F., 3rd, P. Cuijpers, V. Patel, A. Cohen, A. Dias, N. Chowdhary, O. I. Okereke, M. A. Dew, S. J. Anderson, S. Mazumdar, F. Lotrich and S. M. Albert (2012). "Early intervention to reduce the global health and economic burden of major depression in older adults." <i>Annu Rev Public Health</i> 33: 123-135.	4
Richards, A., M. Barkham, J. Cahill, D. Richards, C. Williams and P. Heywood (2003). "PHASE: a randomised, controlled trial of supervised self-help cognitive behavioural therapy in primary care." <i>Br J Gen Pract</i> 53(495): 764-770.	3
Rijken, L. H. V., A. Van Straten, A. Maiwenn, F. Rutten and M. Donker (2006). "Cost-utility of brief psychological treatment for depression and anxiety." <i>British Journal of Psychiatry</i> 188(APR.): 323-329.	3
Romeo, R., A. Patel, M. Knapp and C. Thomas (2004). "The cost-effectiveness of mirtazapine versus paroxetine in treating people with depression in primary care." <i>Int Clin Psychopharmacol</i> 19(3): 125-134.	3
Romeo, R., M. Knapp, S. Banerjee, J. Morris, R. Baldwin, N. Tarrier, N. Pendleton, M. Horan and A. Burns (2011). "Treatment and prevention of depression after surgery for hip fracture in older people: cost-effectiveness analysis." <i>J Affect Disord</i> 128(3): 211-219.	3
Rost, K., J. Fortney, M. Zhang, J. Smith and G. R. Smith, Jr. (1999). "Treatment of depression in rural Arkansas: policy implications for improving care." <i>J Rural Health</i> 15(3): 308-315.	3
Saylan M, Treur MJ, Postema R, Dilbaz N, Savas H, Heeg BM, et al. Cost-effectiveness analysis of aripiprazole augmentation treatment of patients with major depressive disorder compared to olanzapine and quetiapine augmentation in Turkey: A microsimulation approach. <i>Value in Health Regional Issues</i> 2013;2:171-180.	3
Schapira, M., M. Smietniansky, H. Patino, G. Castejon, M. B. Outumuro, J. Esteban, M. Montero Odasso, D. Matusevich and L. Camera (2013). "Medical services use by ambulatory elderly with depression and anxiety. A collaborative care management program." <i>J Am Geriatr Soc</i> 61: S59.	9
Serrano-Blanco, A., D. Suárez, A. Pinto-Meza, M. T. Peñarrubia and J. M. Haro (2009). "Fluoxetine and imipramine: Are there differences in cost-utility for depression in primary care?" <i>J Eval Clin Pract</i> 15(1): 195-203.	3
Serrano-Blanco, A., E. Gabarron, I. Garcia-Bayo, M. Soler-Vila, E. Caramés, M. T. Peñarrubia-Maria, A. Pinto-Meza and J. M. Haro (2006). "Effectiveness and cost-effectiveness of antidepressant treatment in primary health care: A six-month randomised study comparing fluoxetine to imipramine." <i>J Affect Disord</i> 91(2-3): 153-163.	3

Shah, A., B. Wuntakal, J. Fehler and P. Sullivan (2001). "Is a dedicated specialist social worker working exclusively with psychogeriatric inpatients and an associated dedicated domiciliary care package cost-effective?" <i>Int Psychogeriatr</i> 13(3): 337-346.	3
Shapira, B., N. Tubi, H. Drexler, D. Lidsky, A. Calev and B. Lerer (1998). "Cost and benefit in the choice of ECT schedule. Twice versus three times weekly ECT." <i>Br J Psychiatry</i> 172: 44-48.	6
Shen C, Shah N, Findley PA, Sambamoorthi U. Depression treatment and short-term healthcare expenditures among elderly Medicare beneficiaries with chronic physical conditions. <i>J Negat Results Biomed</i> 2013;12	5
Shih, Y. C., N. B. Bekele and Y. Xu (2007). "Use of Bayesian net benefit regression model to examine the impact of generic drug entry on the cost effectiveness of selective serotonin reuptake inhibitors in elderly depressed patients." <i>Pharmacoeconomics</i> 25(10): 843-862.	6
Sicras-Mainar, A., R. Navarro-Artieda, M. Blanca-Tamayo, V. Gimeno-de la Fuente and J. Salvatella-Pasant (2010). "Comparison of escitalopram vs. citalopram and venlafaxine in the treatment of major depression in Spain: Clinical and economic consequences." <i>Curr Med Res Opin</i> 26(12): 2757-2764.	3
Simon, G. E., M. VonKorff, J. H. Heiligenstein, D. A. Revicki, L. Grothaus, W. Katon and E. H. Wagner (1996). "Initial antidepressant choice in primary care: Effectiveness and cost of fluoxetine vs tricyclic antidepressants." <i>JAMA</i> 275(24): 1897-1902.	3
Smit, F., G. Willemse, M. Koopmanschap, S. Onrust, P. Cuijpers and A. Beekman (2006). "Cost-effectiveness of preventing depression in primary care patients: randomised trial." <i>Br J Psychiatry</i> 188: 330-336.	3
Smith, A., L. Graham and S. Senthinathan (2007). "Mindfulness-based cognitive therapy for recurring depression in older people: a qualitative study." <i>Aging Ment Health</i> 11(3): 346-357.	6
Sobocki, P., B. Jonsson, J. Angst and C. Rehnberg (2006). "Cost of depression in Europe." <i>J Ment Health Policy Econ</i> 9(2): 87-98.	4
Sobocki, P., M. Ekman, H. Agren, B. Runeson and B. Jonsson (2006). "The mission is remission: Health economic consequences of achieving full remission with antidepressant treatment for depression." <i>Int J Clin Pract</i> 60(7): 791-798.	6
Stant, A. D., E. M. TenVergert, H. Kluiter, H. J. Conradi, A. Smit and J. Ormel (2009). "Cost-effectiveness of a psychoeducational relapse prevention program for depression in primary care." <i>J Ment Health Policy Econ</i> 12(4): 195-204.	3
Tournier, M., R. Crott, Y. Gaudron and H. Verdoux (2013). "Economic impact of antidepressant treatment duration in naturalistic conditions." <i>Acta Psychiatr Scand</i> 127(5): 365-372.	3
Unutzer, J., W. J. Katon, M. Y. Fan, M. C. Schoenbaum, E. H. Lin, R. D. Della Penna and D. Powers (2008). "Long-term cost effects of collaborative care for late-life depression." <i>Am J Manag Care</i> 14(2): 95-100.	3
Unützer, J., D. Powers, W. Katon and C. Langston (2005). "From establishing an evidence-based practice to implementation in real-world settings: IMPACT as a case study." <i>Psychiatr Clin North Am</i> 28(4): 1079-1092.	6
Wade, A. G., D. Saragoussi, N. Despiegel, C. Francois, F. Guelfucci and M. Toumi (2010). "Healthcare expenditure in severely depressed patients treated with escitalopram, generic SSRIs or venlafaxine in the UK." <i>Curr Med Res Opin</i> 26(5): 1161-1170.	3
van den Berg, M., F. Smit, T. Vos and P. H. van Baal (2011). "Cost-effectiveness of opportunistic screening and minimal contact psychotherapy to prevent depression in primary care patients." <i>PLoS One</i> 6(8): e22884.	3

van der Weele, G. M., M. W. de Waal, W. B. van den Hout, R. C. van der Mast, A. J. de Craen, W. J. Assendelft and J. Gussekloo (2011). "Yield and costs of direct and stepped screening for depressive symptoms in subjects aged 75 years and over in general practice." <i>Int J Geriatr Psychiatry</i> 26(3): 229-238.	4
Vasiliadis HM, Latimer E, Dionne PA, Preville M. The costs associated with antidepressant use in depression and anxiety in community living older adults. <i>European Psychiatry</i> 2013;28.	6
Wiley-Exley, E., M. E. Domino, J. Maxwell and S. E. Levkoff (2009). "Cost-effectiveness of integrated care for elderly depressed patients in the PRISM-E study." <i>J Ment Health Policy Econ</i> 12(4): 205-213.	6
Von Korff, M., W. Katon, T. Bush, E. H. Lin, G. E. Simon, K. Saunders, E. Ludman, E. Walker and J. Unutzer (1998). "Treatment costs, cost offset, and cost-effectiveness of collaborative management of depression." <i>Psychosom Med</i> 60(2): 143-149.	3
Wu, E., P. Greenberg, E. Yang, A. Yu, R. Ben-Hamadi and M. H. Erder (2008). "Comparison of treatment persistence, hospital utilization and costs among major depressive disorder geriatric patients treated with escitalopram versus other SSRI/SNRI antidepressants." <i>Curr Med Res Opin</i> 24(10): 2805-2813.	6
Wunner C, Reichhart C, Strauss B, Söllner W. Effectiveness of a psychosomatic day hospital treatment for the elderly: A naturalistic longitudinal study with waiting time before treatment as control condition. <i>Journal of Psychosomatic Research</i> 2014;76:121-126.	6
Zimmer, B. (1999). "Direct and indirect costs of venlafaxine treatment of depression in the elderly with comorbid medical disorders." <i>Annals of Long-Term Care</i> 7(11): 405-409.	6