

## Bilaga 4. Exkluderade studier

---

### Studier exkluderade från AHRQ:s rapport

#### **Inkluderade patienter med insulinbehandling**

Kibriya MG, Ali L, Banik NG, Khan AK. Home monitoring of blood glucose (HMBG) in Type-2 diabetes mellitus in a developing country. *Diabetes Res Clin Pract* 1999;46:253-7.

#### **Icke-randomiserad interventionsstudie**

Rutten G, van Eijk J, de Nobel E, Beek M, van der Velden H. Feasibility and effects of a diabetes type II protocol with blood glucose self-monitoring in general practice. *Fam Pract* 1990;7:273-8.

#### **Kohortstudie utan kontrollgrupp**

Banister NA, Jastrow ST, Hodges V, Loop R, Gillham MB. Diabetes self-management training program in a community clinic improves patient outcomes at modest cost. *J Am Diet Assoc* 2004;104:807-10.

Franciosi M, Pellegrini F, De Berardis G, Belfiglio M, Di Nardo B, Greenfield S, et al. Self-monitoring of blood glucose in non-insulin-treated diabetic patients:

a longitudinal evaluation of its impact on metabolic control. *Diabet Med* 2005; 22:900-6.

Halimi S, Charpentier G, Grimaldi A, Grenier JL, Baut F, Germain B, Magnette J. Effect on compliance, acceptability of blood glucose self-monitoring and HbA(1c) of a self-monitoring system developed according to patient's wishes. The ACCORD study. *Diabetes Metab* 2001;27:681-7.

Miles P, Everett J, Murphy J, Kerr D. Comparison of blood or urine testing by patients with newly diagnosed non-insulin dependent diabetes: patient survey after randomised crossover trial. [erratum appears in *BMJ* 1998 Jan 17;316:195]. *BMJ* 1997;315:348-9.

Ozmen B, Boyvada S. The relationship between self-monitoring of blood glucose control and glycosylated haemoglobin in patients with type 2 diabetes with and without diabetic retinopathy. *J Diabet Complications* 2003;17:128-34.

### **Fel intervention, kontrollintervention eller effektmått**

Afandi B, Ahmad S, Saadi H, Elkhumaidi S, Karkoukli MA, Kelly B, et al. Audit of a diabetes clinic at Tawam hospital, United Arab Emirates, 2004–2005. *Ann N Y Acad Sci* 2006;1084:319-24.

Benhamou PY, Melki V, Boizel R, Perreal F, Quesada JL, Bessieres-Lacombe S, et al.

One-year efficacy and safety of Web-based follow-up using cellular phone in type 1 diabetic patients under insulin pump therapy: the PumpNet study. *Diabetes Metab* 2007;33:220-6.

Brown SA, Garcia AA, Kouzekanani K, Hanis CL. Culturally competent diabetes

- self-management education for Mexican Americans: The Starr County border health initiative. *Diabetes Care* 2002; 25:259-68.
- Clarke P, Coleman MA, Holt RI. Alternative site self blood glucose testing is preferred by women with gestational diabetes. *Diabetes Technol Ther* 2005;7:604-8.
- Cox DJ, Kovatchev B, Koev D, Koeva L, Dachev S, Tcharaktchiev D, et al. Hypoglycemia anticipation, awareness and treatment training (HAATT) reduces occurrence of severe hypoglycemia among adults with type 1 diabetes mellitus. *Int J Behav Med* 2004;11:212-8.
- Deiss D, Hartmann R, Schmidt J, Kordonouri O. Results of a randomised controlled cross-over trial on the effect of continuous subcutaneous glucose monitoring (CGMS) on glycaemic control in children and adolescents with type 1 diabetes. *Exp Clin Endocrinol Diabetes* 2006;114:63-7.
- Elnour AA, El Mugammar IT, Jaber T, Revel T, McElnay JC. Pharmaceutical care of patients with gestational diabetes mellitus. *J Eval Clin Pract* 2008;14: 131-40.
- Farmer A, Gibson OJ, Tarassenko L, Neil A. A systematic review of telemedicine interventions to support blood glucose self-monitoring in diabetes. *Diabet Med* 2005; 22:1372-8.
- von Ferber L, Koster I, Hauner H. [Costs of the treatment of hyperglycemia in patients with diabetes mellitus. The impact of age, type of therapy and complications: results of the German CoDiM study]. *Med Klin (Munich)* 2006;101:384-93.
- Gagliardino JJ, Lapertosa S, Villagra M, Caporale JE, Oliver P, Gonzalez C, et al. PRODIACOR: a patient-centered treatment program for type 2 diabetes and associated cardiovascular risk factors in the city of Corrientes, Argentina: study design and baseline data. *Contemp Clin Trials* 2007;28:548-56.
- Garg S, Zisser H, Schwartz S, Bailey T, Kaplan R, Ellis S, Jovanovic L. Improvement in glycemic excursions with a transcutaneous, real-time continuous glucose sensor: a randomized controlled trial. *Diabetes Care* 2006;29:44-50.
- Hee-Sung K. Impact of Web-based nurse's education on glycosylated haemoglobin in type 2 diabetic patients. *J Clin Nurs* 2007; 16:1361-6.
- Heneghan C, Perera R, Ward AA, Fitzmaurice D, Meats E, Glasziou P. Assessing differential attrition in clinical trials: self-monitoring of oral anticoagulation and type II diabetes. *BMC Med Res Methodol* 2007;7:18.
- Hill-Briggs F, Yeh HC, Gary TL, Batts-Turner M, D'Zurilla T, Brancati FL. Diabetes problem-solving scale development in an adult, African American sample. *Diabetes Educ* 2007;33:291-9.
- Hirsch IB, Abelson J, Bode BW, Fischer JS, Kaufman FR, Mastrototaro J, et al. Sensor-augmented insulin pump therapy: results of the first randomized treat-to-target study. *Diabetes Technol Ther* 2008; 10:377-83.
- Karter AJ, Stevens MR, Brown AF, Duru OK, Gregg EW, Gary TL, et al. Educational disparities in health behaviors among patients with diabetes: the Translating Re-

- search Into Action for Diabetes (TRIAD) Study. *BMC Public Health* 2007;7:308.
- King AB, Wolfe GS, Armstrong DU. Evaluation of a patient education booklet (SimpleStart) effect on postprandial glucose control in type 2 diabetes. *Diabetes Technol Ther* 2007;9:241-5.
- Krass I, Armour CL, Mitchell B, Brilliant M, Dienaar R, Hughes J, et al. The Pharmacy Diabetes Care Program: assessment of a community pharmacy diabetes service model in Australia. *Diabet Med* 2007;24:677-83.
- Kuwa, K. [Noninvasive blood glucose monitoring: new technology using metabolic heat conformation method]. *Rinsho Byori* 2006;54:519-25.
- Mastura I, Mimi O, Piterman L, Teng CL, Wijesinha S. Self-monitoring of blood glucose among diabetes patients attending government health clinics. *Med J Malaysia* 2007;62:147-51.
- Miller ST, Schlundt DG, Larson C, Reid R, Pichert JW, Hargreaves M, et al. Exploring ethnic disparities in diabetes, diabetes care, and lifestyle behaviors: the Nashville REACH 2010 community baseline survey. *SOURCE* 2004;14 (3 Suppl 1):S38-45. Date of Publication: Jun 2004.
- Mitchell CG, Bowker SL, Majumdar SR, Toth EL, Johnson JA. Lack of correlation between patient-reported outcomes and glycemic control in type 2 diabetes not managed by insulin. *Canadian Journal of Diabetes* 2004;28:362-8.
- Muller U, Hammerlein A, Casper A, Schulz M. Community pharmacy-based intervention to improve self-monitoring of blood glucose in type 2 diabetic patients. *Pharmacy Practice* 2006;4:195-203.
- Palmer AJ, Dinneen S, Gavin JR, 3rd, Gray A, Herman WH, Karter AJ. Cost-utility analysis in a UK setting of self-monitoring of blood glucose in patients with type 2 diabetes. *Curr Med Res Opin* 2006;22:861-72.
- Perriello G, Pampanelli S, Di Pietro C, Brunetti P. Comparison of glycaemic control over 1 year with pioglitazone or gliclazide in patients with Type 2 diabetes. *Diabet Med* 2006;23:246-52.
- Raskin P, Klaff L, McGill J, South SA, Hollander P, Khutoryansky N, Hale PM. Efficacy and safety of combination therapy: Repaglinide plus metformin versus nateglinide plus metformin. *Diabetes Care* 2003;26:2063-8.
- Rosal MC, Olendzki B, Reed GW, Gumieniak O, Scavron J, Ockene I. Diabetes self-management among low-income Spanish-speaking patients: a pilot study. *Ann Behav Med* 2005;29:225-35.
- Saloranta C, Guitard C, Pecher E, De Pablos-Velasco P, Lahti K, Brunel P, Groop L. Nateglinide improves early insulin secretion and controls postprandial glucose excursions in a prediabetic population. *Diabetes Care* 2002;25:2141-6.
- Scherbaum WA, Ohmann C, Abholz HH, Dragano N, Lankisch M. Effect of the frequency of self-monitoring blood glucose in patients with type 2 diabetes treated with oral antidiabetic drugs-a multi-centre, randomized controlled trial. *PLoS ONE* 2008;3:e3087.

Wen L, Parchman ML, Linn WD, Lee S. Association between self-monitoring of blood glucose and glycemic control in patients with type 2 diabetes mellitus. *Am J Health Syst Pharm* 2004;61:2401-5.

Woo J, Sea MM, Tong P, Ko GT, Lee Z, Chan J, Chow FC. Effectiveness of a lifestyle modification programme in

weight maintenance in obese subjects after cessation of treatment with Orlistat. *J Eval Clin Pract* 2008;13:853-9.

Yoo HJ, An HG, Park SY, Ryu OH, Kim HY, Seo JA, et al. Use of a real time continuous glucose monitoring system as a motivational device for poorly controlled type 2 diabetes. *Diabetes Res Clin Pract* 2008;82:73-9. Epub 2008 Aug 12.

## Insulinbehandlade patienter

Albisser AM, Wright CE, Sakkal S. Averting iatrogenic hypoglycemia through glucose prediction in clinical practice: progress towards a new procedure in diabetes. *Diabetes Res Clin Pract* 2007;76:207-14.

Ashwell SG, Gebbie J, Home PD. Twice-daily compared with once-daily insulin glargine in people with Type 1 diabetes using meal-time insulin aspart. *Diabet Med* 2006;23:879-86.

Ashwell SG, Gebbie J, Home PD. Optimal timing of injection of once-daily insulin glargine in people with Type 1 diabetes using insulin lispro at meal-times. *Diabet Med* 2006;23:46-52.

Bergenstal RM, Johnson M, Powers MA, Wynne A, Vlainic A, Hollander P, Rendell M. Adjust to target in type 2 diabetes: comparison of a simple algorithm with carbohydrate counting for adjustment of mealtime insulin glulisine. *Diabetes Care* 2008;31:1305-10.

Bretzel RG, Nuber U, Landgraf W, Owens DR, Bradley C, Linn T. Once-daily basal insulin glargine versus thrice-daily prandial insulin lispro in people with type 2 diabetes on oral hypoglycaemic agents (APOLLO): an open randomised controlled trial. *Lancet* 2008;371:1073-84.

Brodows RG, Qu Y, Johns D, Kim D, Holcombe JH. Quantifying the effect of exenatide and insulin glargine on postprandial glucose excursions in patients with type 2 diabetes. *Curr Med Res Opin* 2008;24:1395-7.

Forti A, Garcia EG, Yu MB, Jimenez MC, Brodows RG, Oliveira JH. Efficacy and safety of exenatide administered before the two largest daily meals of Latin American patients with type 2 diabetes. *Curr Med Res Opin* 2008;24:2437-47.

Chen JW, Lauritzen T, Bojesen A, Christiansen JS. Multiple mealtime administration of biphasic insulin aspart 30 versus traditional basal-bolus human insulin treatment in patients with type 1 diabetes. *Diabetes Obes Metab* 2006;8:682-9.

Chen HS, Wu TE, Jap TS, Lin SH, Hsiao LC, Lin HD. Improvement of glycaemia control in subjects with type 2 diabetes by self-monitoring of blood glucose: comparison of two management programs adjusting bedtime insulin dosage. *Diabetes Obes Metab* 2008;10:34-40.

Haak T, Tiengo A, Draeger E, Suntum M, Waldhausl W. Lower within-subject variability of fasting blood glucose and reduced

weight gain with insulin detemir compared to NPH insulin in patients with type 2 diabetes. *Diabetes Obes Metab* 2005;7:56-64.

Kennedy L, Herman WH, Strange P, Harris A. Impact of active versus usual algorithmic titration of basal insulin and point-of-care versus laboratory measurement of HbA1c on glycemic control in patients with type 2 diabetes: the Glycemic Optimization with Algorithms and Labs at Point of Care (GOAL A1C) trial. *Diabetes Care* 2006;29:1-8.

Lagarde WH, Barrows FP, Davenport ML, Kang M, Guess HA, Calikoglu AS. Continuous subcutaneous glucose monitoring in children with type 1 diabetes mellitus: a single-blind, randomized, controlled trial. *Pediatr Diabetes* 2006;7:159-64.

Lush CW, Darsow T, Zhang B, Lorenzi G, Frias JP. Pramlintide as an adjunct to basal insulin: Effects on glycemic control and weight in patients with type 2 diabetes mellitus. *Insulin* 2007;2:166-72.

Murata GH, Duckworth WC, Shah JH, Wendel CS, Mohler MJ, Hoffman RM. Hypoglycemia in stable, insulin-treated veterans with type 2 diabetes: a prospect-

ive study of 1662 episodes. *J Diabetes Complications* 2005;19:10-7.

Oria-Pino A, Montero-Perez FJ, Luna-Morales S, del Campo-Vazquez P, Sanchez-Guijo P. [Effectiveness and efficacy of self-measurement of capillary blood glucose in patients with type 2 diabetes mellitus]. *Med Clin (Barc)* 2006;126:728-35.

Sheppard P, Bending JJ, Huber JW. Pre- and post-prandial capillary glucose self-monitoring achieves better glycaemic control than pre-prandial only monitoring. A study in insulin treated diabetic patients. *Practical Diabetes International* 2005;22:15-22.

Tirgoviste CI, Strachinariu R, Farcasiu E, Milicevic Z, Teodorescu G. Humalog Mix 25 in patients with type 2 diabetes which do not achieve acceptable glycemic control with oral agents: results from a phase III, randomized, parallel study. *Rom J Intern Med* 2003;41:153-62.

Yates K, Hasnat Milton A, Dear K, Ambler G. Continuous glucose monitoring-guided insulin adjustment in children and adolescents on near-physiological insulin regimens: a randomized controlled trial. *Diabetes Care* 2006;29:1512-7.

## Studier som inkluderat även typ 1-diabetes

Abdelgadir M, Elbagir M, Eltom M, Berne C. The influence of glucose self-monitoring on glycaemic control in patients with diabetes mellitus in Sudan. *Diabetes Res Clin Pract* 2006;74:90-4.

Cobry E, Chase HP, Burdick P, McFann K, Yetzer H, Scrimgeour L. Use of CoZ-monitor in youth with type 1 diabetes. *Pediatr Diabetes* 2008;9:148-51.

Golicki DT, Golicka D, Groele L, Pankowska E. Continuous Glucose Monitoring System in children with type 1 diabetes mellitus: a systematic review and meta-analysis. *Diabetologia* 2008;51:233-40.

Kovatchev BP, Crean J, McCall A. Pramlintide reduces the risks associated with glucose variability in type 1

diabetes. *Diabetes Technol Ther* 2008; 10:391-6.

Lecomte P, Romon I, Fosse S, Simon D, Fagot-Campagna A. Self-monitoring of blood glucose in people with type 1 and type 2 diabetes living in France: the Entred study 2001. *Diabetes Metab* 2008;34:219-26.

Radman M, Jurisic D, Ljusic D, Jerkovic R, Kovacic N, Hozo IS. Assessing glycemia

in type 1 diabetic patients using a microdialysis system for continuous glucose monitoring. *Ann Saudi Med* 2007;27: 166-70.

Tamborlane WV, Ruedy KJ, Wysocki T, O'Grady M, Kollman C, Block J, et al. JDRF randomized clinical trial to assess the efficacy of real-time continuous glucose monitoring in the management of type 1 diabetes: Research design and methods. *Diabetes Technol Ther* 2008;10:310-21.

## **Inga primärdata, översikt/beskrivning av kommande studie, "editorial", kommentar, dubbelpublikation**

Home glucose monitoring makes little difference in type 2 diabetes. *J Fam Pract* 2005;54:936.

Bouter LM. [From the Cochrane Library and *Diabetes Care*: self-monitoring of blood glucose probably an effective way to improve glycaemic control in patients with type 2 diabetes not taking insulin]. *Ned Tijdschr Geneesk* 2006;150:1826-9.

Davis WA, Bruce DG, Davis TME. Is self-monitoring of blood glucose appropriate for all type 2 diabetic patients? The fremantle diabetes study: Response to Kolb et al. [5]. *Diabetes Care* 2007;30:184-5.

Farmer A, Waldron-Lynch F, Dinneen S. Self-monitoring of blood glucose did not improve glycaemic control in patients with type 2 diabetes not treated with insulin: Commentary. *Evid Based Med* 2008;13:7.

Gallo A, Tiengo A. Self monitoring blood glucose in type 2 diabetic patients [L'autocontrollo domiciliare della glicemia nel soggetti con diabete mellito tipo 2]. *Giornale Italiano di Diabetologia e Metabolismo* 2005;25:213-20.

Gerich JE, Odawara M, Terauchi Y. The rationale for paired pre- and postprandial self-monitoring of blood glucose: The role of glycemic variability in micro- and macrovascular risk. *Curr Med Res Opin* 2007;23:1791-8.

Gulliford M. Self monitoring of blood glucose in type 2 diabetes. *BMJ* 2008; 336:1139-40.

Heneghan C, Perera R, Ward AA, Fitzmaurice D, Meats E, Glasziou P. Assessing differential attrition in clinical trials: Self-monitoring of oral anticoagulation and type II diabetes. *BMC Med Res Methodol* 2007;7:18.

Karter AJ. Role of self-monitoring of blood glucose in glycemic control. *Endocr Pract* 2006;12:110-7.

Krentz AJ, Bailey CJ. Oral antidiabetic agents: current role in type 2 diabetes mellitus. *Drugs* 2005;65:385-411.

McAndrew L, Schneider SH, Burns E, Leventhal H. Does patient blood glucose monitoring improve diabetes control? A

systematic review of the literature. *Diabetes Educ* 2007;33:991-1011; discussion 1012-3.

McGeoch G, Derry S, Moore RA. Self-monitoring of blood glucose in type-2 diabetes: what is the evidence?. *Diabetes Metab Res Rev* 2007;23:423-40.

Murff HJ. Routine self-monitoring of blood glucose does not improve glycemic control in noninsulin-treated diabetic patients: Commentary. *J Clin Outcomes Manag* 2007;14:537-8.

Poolsup N, Suksomboon N, Jiamsathit W. Systematic review of the benefits of self-monitoring of blood glucose on glycemic control in type 2 diabetes patients. *Diabetes Technol Ther* 2008;10:S51-S66.

Sarol JN, Jr, Nicodemus NA, Jr, Tan KM, Grava MB. Self-monitoring of blood glucose as part of a multi-component therapy among non-insulin requiring type 2 diabetes patients: a meta-analysis (1966–2004). *Curr Med Res Opin* 2005;21:173-84.

Siebolds M, Gaedeke O, Schwedes U. Self-monitoring of blood glucose – psychological aspects relevant to changes in HbA1c in type 2 diabetic patients treated with diet or diet plus oral antidiabetic medication. *Patient Educ Couns* 2006;62:104-10.

Slack S. Near patient testing for glycated haemoglobin in people with type 2 diabetes mellitus managed in primary care: acceptability and satisfaction; Impact of self-mon-

itoring of blood glucose in the management of patients with non-insulin-treated diabetes: open parallel group randomized trial. *Ann Clin Biochem* 2008;45:114-5.

Towfigh A, Romanova M, Weinreb JE, Munjas B, Suttorp MJ, Zhou A, Shekelle PG. Self-monitoring of blood glucose levels in patients with type 2 diabetes mellitus not taking insulin: a meta-analysis. *Am J Manag Care* 2008;14:468-75.

Welschen LM, Bloemendal E, Nijpels G, Dekker JM, Heine RJ, Stalman WA, Bouter LM. Self-monitoring of blood glucose in patients with type 2 diabetes who are not using insulin. *Cochrane Database Syst Rev* 2005:CD005060.

Welschen LM, Bloemendal E, Nijpels G, Dekker JM, Heine RJ, Stalman WA, Jansen JP. Self-monitoring of glucose in type 2 diabetes mellitus: a Bayesian meta-analysis of direct and indirect comparisons. *Curr Med Res Opin* 2006;22:671-81.

Welschen LMC, Bloemendal E, Nijpels G, Dekker JM, Heine RJ, Stalman WAB, Bouter LM. From the Cochrane Library and Diabetes Care: Self-monitoring of blood glucose probably an effective way to improve glycaemic control in patients with type 2 diabetes not taking insulin [Zelfcontrole van bloedglucosewaarden bij diabetes mellitus type 2 zonder gebruik van insuline waarschijnlijk effectieve methode om glykemische instelling te verbeteren]. *Ned Tijdschr Geneesk* 2006;150:1826-9.

## Ej RCT

Karter AJ, Parker MM, Moffet HH, Spence MM, Chan J, Ettner SL, Selby JV. Longitudinal study of new and

prevalent use of self-monitoring of blood glucose. *Diabetes Care* 2006; 29:1757-63.