

Studier med låg kvalitet

Beteende och livsstil	orsak
Aslan E, Komurcu N, Beji NK, Yalcin O. Bladder training and Kegel exercises for women with urinary complaints living in a rest home. Gerontology 2008;54:224-31.	Stort bortfall, randomisering och blindning framgår inte.
Colling J, Ouslander J, Hadley BJ, Eisch J, Campbell E. The effects of patterned urge-response toileting (PURT) on urinary incontinence among nursing home residents. J Am Geriatr Soc. [Clinical Trial Randomized Controlled Trial Research Support, U.S. Gov't, P.H.S.]. 1992 Feb;40(2):135-41.	Signifika baslinjeskillnader, stort bortfall. Det framgår inte om datainsamlare/utvärderare var blindade.
Colling J, Owen TR, McCreedy M, Newman D. The effects of a continence program on frail community-dwelling elderly persons. Urologic Nursing. 2003;23:117-22.	Det framgår inte om datainsamlare/utvärderare var blindade. Stort bortfall, baslinjeskillander.
Creason NS, Grybowski JA, Burgener S, Whippo C, Yeo S, Richardson B. Prompted voiding therapy for urinary incontinence in aged female nursing home residents. Journal of Advanced Nursing. 1989;14:120-6.	Det framgår inte om datainsamlare/utvärderare var blindade. Dåligt beskrivna resultat.
Filocamo MT, Li M, Del P, Cecconi F, Villari D, Marzocco M, et al. Pharmacologic Treatment in Postprostatectomy Stress Urinary Incontinence. European Urology 2007;51:1559-1564.	Randomisering och bortfall framgår inte.
Guibert R, Rodda M, Barda D. A randomised controlled trial of an IT technology medical device on urinary incontinence in a residential aged care facility 16th National Conference on Incontinence, Gold Coast, Queensland, 2007. Australian & New Zealand Continence Journal. 2007;13:119-20.	Randomisering och bortfall framgår inte. Liten studie, abstract från konferens?
Jirovec MM, Templin T. Predicting success using individualized scheduled toileting for memory-impaired elders at home. Res Nurs Health 2001;24:1-8.	Stort bortfall. Det framgår inte om datainsamlare/utvärderare var blindade.
Kincade JE, Dougherty MC, Carlson JR, Hunter GS, Busby-Whitehead J. Randomized clinical trial of efficacy of self-monitoring techniques to treat urinary incontinence in women. Neurourology and Urodynamics. 2007;26:507-11.	Dåligt beskriven för vår population. Bortfall framgår inte. Det framgår inte om

Studier med låg kvalitet

	datainsamlare/utvärderare var blindade.
Krichbaum K, Pearson V, Savik K, Mueller C. Improving resident outcomes with GAPN organization level interventions gerontological advanced practice nurses. <i>Western Journal of Nursing Research.</i> 2005;27:322-37.	Det framgår inte om datainsamlare/utvärderare var blindade. Stort bortfall.
Ryden Muriel B, Snyder M, Gross Cynthia R, Savik K, Pearson V, Krichbaum K, et al. Value-added outcomes: The use of advanced practice nurses in long-term care facilities. <i>The Gerontologist.</i> 2000;40:654-62.	Det framgår inte om datainsamlare/utvärderare var blindade, stort bortfall.
Simmons SF, Ouslander JG. Resident and family satisfaction with incontinence and mobility care: sensitivity to intervention effects? <i>Gerontologist</i> 2005;45:318-26.	Stort bortfall randomiseering och blindning framgår inte.
Schnelle JF, Traughber B, Sowell VA, Newman DR, Petrilli CO, Ory M. Prompted voiding treatment of urinary incontinence in nursing home patients. A behavior management approach for nursing home staff. <i>J Am Geriatr Soc.</i> 1989;37:1051-7.	Bortfall oklart, randomiseering oklar, blinding oklar.
Tobin GW, Brocklehurst JC. The management of urinary incontinence in local authority residential homes for the elderly. <i>Age Ageing</i> 1986;15:292-8.	Problematiska baslinjeskillnader. Det framgår inte om datainsamlare/utvärderade vad blindade.
Wenger NS, Roth CP, Hall WJ, Ganz DA, Snow V, Byrkit J, et al. Practice redesign to improve care for falls and urinary incontinence: primary care intervention for older patients. <i>Arch Intern Med.</i> 2010;170:1765-72.	Randomisering framgår inte.

Kirurgi	orsak
Dalpiaz O, Primus G, Schips L. SPARC sling system for treatment of female stress urinary incontinence in the elderly. <i>Eur Urol</i> 2006;50:826-30; discussion 830.	Liten population, uppgift om bortfall saknades
Herschorn S, Radomski SB. Collagen injections for genuine stress urinary incontinence: patient selection and durability. <i>Int Urogynecol J Pelvic Floor Dysfunct</i> 1997;8:18-24.	Liten population, uppgift om bortfall saknades
Lo TS, Huang HJ, Chang CL, Wong SY, Horng SG, Liang CC. Use of intravenous anesthesia for tension-free vaginal tape therapy in elderly	Liten population, uppgift om bortfall

Studier med låg kvalitet

women with genuine stress incontinence. Urology 2002;59:349-53.	saknades
Touloupidis S, Papatsoris AG, Thanopoulos C, Charalambous S, Giannakopoulos S, Rombis V. Tension-free vaginal tape for the treatment of stress urinary incontinence in geriatric patients. Gerontology 2007;53:125-7.	Liten population, uppgift om bortfall saknades

Läkemedel	orsak
Isik AT, Celik T, Bozoglu E, Doruk H. Trospium and cognition in patients with late onset Alzheimer disease. J Nutr Health Aging. 2009;13:672-6.	Stort bortfall, oklart om blindning
Minassian VA, Ross S, Sumabat O, Lovatsis D, Pascali D, Al-Badr A, et al. Randomized trial of oxybutynin extended versus immediate release for women aged 65 and older with overactive bladder: lessons learned from conducting a trial. J Obstet Gynaecol Can. 2007;29:726-32.	Stort bortfall , ingen blindning, avbruten studie pga dålig power.