

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

Alm A, Hornmark AM, Fall PA, Linder L, Bergstrand B, Ehrnebo M, et al. Care of pressure sores: a controlled study of the use of a hydrocolloid dressing compared with wet saline gauze compresses. <i>Acta Derm Venereol Suppl (Stockh)</i> . 1989;149:1-10.
Alvarez OM, Fernandez-Obregon A, Rogers RS, Bergamo L, Masso J, Black M. A prospective, randomized, comparative study of collagenase and papain-urea for pressure ulcer debridement. <i>Wounds: A Compendium of Clinical Research & Practice</i> 2002;14:293-301.
Alvarez OM, Fernandez-Obregon A, Rogers RS, Bergamo L, Masso J, Black M. A prospective, randomized, comparative study of collagenase and papain-urea for pressure ulcer debridement. <i>Wounds: A Compendium of Clinical Research & Practice</i> 2003;15:293-301.
Amione P, Ricci E, Topo F, Izzo L, Pirovano R, Rega V, et al. Comparison of Allevyn Adhesive and Biatain Adhesive in the management of pressure ulcers [corrected] [published erratum appears in <i>J WOUND CARE</i> 2005 Sep;14(8):insert 1p]. <i>Journal of wound care</i> . 2005;14:365-70.
Anders J, Heinemann A, Leffmann C, Leutenegger M, Profener F, von R-K. Decubitus ulcers: pathophysiology and primary prevention. <i>Dtsch Arztebl Int</i> 2010;107:371-81; quiz 382.
Baier RR, Gifford DR, Lyder CH, Schall MW, Funston-Dillon DL, Lewis JM, et al. Quality improvement for pressure ulcer care in the nursing home setting: the Northeast Pressure Ulcer Project. <i>J Am Med Dir Assoc</i> 2003;4:291-301.
Bale S, Banks V, Haglestein S, Harding KG. A comparison of two amorphous hydrogels in the debridement of pressure sores. <i>Journal of wound care</i> . 1998;7:65-8.
Ballard K. Pressure-relief mattresses and patient comfort Debut MR (SSI Hill-Rom) mattress Nimbus (HNE Huntleigh) mattress. <i>Professional Nurse</i> . 1997;13:27-8.
Banks S, Bridel J. A descriptive evaluation of pressure-reducing cushions. <i>British Journal of Nursing</i> 1995;4:736, 738, 740 passim.
Banks V, Bale S, Harding K, Harding EF. Evaluation of a new polyurethane foam dressing. <i>Journal of wound care</i> . 1997;6:266-9.
Banks V, Bale SE, Harding KG. Comparing two dressings for exuding pressure sores in community patients. <i>J Wound Care</i> 1994;3:175-8.
Barrois B, Carles M, Rumeau M, Tell L, Toussaint JF, Bonnefoy M, et al. Efficacy and tolerability of hyaluronan (ialuset) in the treatment of pressure ulcers: a multicentre, non-randomised, pilot study. <i>Drugs R D</i> 2007;8:267-73.
Beeckman D, Clays E, Van Hecke A, Vanderwee K, Schoonhoven L, Verhaeghe S. A multi-faceted tailored strategy to implement an electronic clinical decision support system for pressure ulcer prevention in nursing homes: A two-armed randomized controlled trial. <i>Int J Nurs Stud</i> 2013;50:475-486.
Beele H, Meuleneire F, Nahuys M, Percival SL. A prospective randomised open label study to evaluate the potential of a new silver alginate/carboxymethylcellulose antimicrobial wound dressing

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

to promote wound healing. <i>Int Wound J.</i> 2010;7:262-70.
Bennett RG, Bellantoni MF, Ouslander JG. Air-fluidized bed treatment of nursing home patients with pressure sores. <i>J Am Geriatr Soc.</i> 1989;37:235-42.
Bergquist S, Frantz R. Braden Scale: validity in community-based older adults receiving home health care. <i>Applied Nursing Research</i> 2001;14:36-43.
Bergstrom N, Horn SD, Smout RJ, Bender SA, Ferguson ML, Taler G, et al. The National Pressure Ulcer Long-Term Care Study: outcomes of pressure ulcer treatments in long-term care. <i>J Am Geriatr Soc</i> 2005;53:1721-9.
Bergstrom N, Smout R, Horn S, Spector W, Hartz A, Limcangco MR. Stage 2 pressure ulcer healing in nursing homes. <i>J Am Geriatr Soc.</i> 2008;56:1252-8.
Biglari B, Vd L, P H, Simon A, Aytac S, Gerner HJ, et al. Use of Medihoney as a non-surgical therapy for chronic pressure ulcers in patients with spinal cord injury. <i>Spinal Cord</i> 2012;50:165-169.
Bito S, Mizuhara A, Oonishi S, Takeuchi K, Suzuki M, Akiyama K, et al. Randomised controlled trial evaluating the efficacy of wrap therapy for wound healing acceleration in patients with NPUAP stage II and III pressure ulcer. <i>BMJ Open.</i> 2012;2:e000371.
Boykin A, Winland-Brown J. Pressure sores: nursing management compare hydrocolloid occlusive dressing with povidone-iodine therapy. <i>Using nursing research National League for Nursing.</i> 1989;11.
Breslow RA. The effect of dietary protein intake on pressure sore healing and nutritional status in nursing home subjects. <i>University Of Maryland College Park.</i> 1990;129.
Brod M, McHenry E, Plasse TF, Fedorczyk D, Trout JR. A randomized comparison of poly-hema and hydrocolloid dressings for treatment of pressure sores. <i>Arch Dermatol.</i> 1990;126:969-70.
Brown-Etris M, Milne C, Orsted H, Gates JL, Netsch D, Punchello M, et al. A prospective, randomized, multisite clinical evaluation of a transparent absorbent acrylic dressing and a hydrocolloid dressing in the management of Stage II and shallow Stage III pressure ulcers. <i>Adv Skin Wound Care.</i> 2008;21:169-74.
Callaghan S, Trapp M. Evaluating two dressings for the prevention of nasal bridge pressure sores. <i>Professional Nurse</i> 1998;13:361-4.
Cassino R, Ippolito AM, Cuffaro C, Corsi A, Ricci E. A controlled, randomised study on the efficacy of two overlays in the treatment of decubitus ulcers. <i>Minerva Chir</i> 2013;68:105-16.
Cassino R, Ricci E. Effectiveness of topical application of amino acids to chronic wounds: a prospective observational study. <i>Journal of wound care.</i> 2010;19:29-34.
Ceolim MF, Araujo IEM, Diogo MJE, Bajay HM, Rosa KM. Comparing hydrocolloid to dry gauze dressings in the treatment of sacral pressure sores. <i>Nursing: A New Era for Action, International</i>

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

Council of Nurses ICN 22nd Quadrennial Congress. 2001.
Cereda E, Gini A, Pedrolli C, Vanotti A. Disease-specific, versus standard, nutritional support for the treatment of pressure ulcers in institutionalized older adults: a randomized controlled trial. <i>J Am Geriatr Soc</i> 2009;57:1395-402.
Chang KW, Alsagoff S, Ong KT, Sim PH. Pressure ulcers--randomised controlled trial comparing hydrocolloid and saline gauze dressings. <i>Med J Malaysia</i> 1998;53:428-31.
Chuangsuwanich A, Charmsanti O, Lohsiriwat V, Kangwanpoom C, Thong-In N. The efficacy of silver mesh dressing compared with silver sulfadiazine cream for the treatment of pressure ulcers. <i>J Med Assoc Thai</i> 2011;94:559-65.
Colwell JC, Foreman MD, Trotter JP. A comparison of the efficacy and cost-effectiveness of two methods of managing pressure ulcers. <i>Decubitus: The Journal of Skin Ulcers</i> 1993;6:28-36.
Cullum N, Petherick E. Pressure ulcers. <i>Clin Evid (Online)</i> 2008;2008.
Davis P, Wood L, Wood Z, Eaton A, Wilkins J. Clinical experience with a glucose oxidase-containing dressing on recalcitrant wounds. <i>Journal of wound care</i> . 2009;18:116-21.
Dehlin O, Elmståhl S, Gottrup F. Monochromatic phototherapy: effective treatment for grade II chronic pressure ulcers in elderly patients. <i>Aging Clinical & Experimental Research</i> . 2007;19:478-83.
Desneves KJ, Todorovic BE, Cassar A, Crowe TC. Treatment with supplementary arginine, vitamin C and zinc in patients with pressure ulcers: a randomised controlled trial. <i>Clin Nutr</i> . 2005;24:979-87.
Durovic A, Maric D, Brdareski Z, Jevtic M, Durdevic S. The effects of polarized light therapy in pressure ulcer healing. <i>Vojnosanit Pregl</i> . 2008;65:906-12.
Edsberg LE, Brogan MS, Jaynes CD, Fries K. Topical hyperbaric oxygen and electrical stimulation: exploring potential synergy. <i>Ostomy Wound Management</i> 2002;48:42-4.
Felzani G, Spoletini I, Convento A, Di L, Rossi P, Miceli M, et al. Effect of lysine hyaluronate on the healing of decubitus ulcers in rehabilitation patients. <i>Adv Ther</i> . 2011;28:439-45.
Flynn D, Williams S. Barrier creams for skin breakdown. <i>Nursing & Residential Care</i> . 2011;13:553-4.
Franek A, Kostur R, Polak A, Taradaj J, Szlachta Z, Blaszczyk E, et al. Using high-voltage electrical stimulation in the treatment of recalcitrant pressure ulcers: results of a randomized, controlled clinical study. <i>Ostomy Wound Management</i> 2012;58:30-44.
Frykberg RG, Driver VR, Carman D, Lucero B, Borris-Hale C, Fylling CP, et al. Chronic wounds treated with a physiologically relevant concentration of platelet-rich plasma gel: a prospective case series. <i>Ostomy Wound Management</i> . 2010;56:36-8.
Gentzkow GD, Alon G, Taler GA, Eltorai IM, Montroy RE. Healing of refractory stage III and IV pressure ulcers by a new electrical stimulation device. <i>Wounds: A Compendium of Clinical Research</i>

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

& Practice 1993;5:160-72.
Gray M, Jones DP. The effect of different formulations of equivalent active ingredients on the performance of two topical wound treatment products. <i>Ostomy Wound Management</i> 2004;50:34-6.
Hamilton L, Cleverly C. A skin care resource nurse program: is it effective? <i>Perspectives: The Journal of the Gerontological Nursing Association</i> 1994;18:10-14.
Harrison-Balestra C, Eaglstein WH, Falabela AF, Kirsner RS. Recombinant human platelet-derived growth factor for refractory nondiabetic ulcers: A retrospective series. <i>Dermatologic Surgery</i> 2002;28:755-760.
Health Quality O. Management of chronic pressure ulcers: an evidence-based analysis. <i>Ont Health Technol Assess Ser</i> 2009;9:1-203.
Health Quality O. Negative pressure wound therapy: an evidence-based analysis. <i>Ont Health Technol Assess Ser</i> 2006;6:1-38.
Ho CH, Powell HL, Collins JF, Bauman WA, Spungen AM. Poor nutrition is a relative contraindication to negative pressure wound therapy for pressure ulcers: preliminary observations in patients with spinal cord injury. <i>Adv Skin Wound Care</i> 2010;23:508-16.
Hommel A, Ulander K, Thorngren KG. Improvements in pain relief, handling time and pressure ulcers through internal audits of hip fracture patients. <i>Scand J Caring Sci</i> 2003;17:78-83.
Honde C, Derks C, Tudor D. Local treatment of pressure sores in the elderly: amino acid copolymer membrane versus hydrocolloid dressing. <i>J Am Geriatr Soc.</i> 1994;42:1180-3.
Houghton PE, Campbell KE, Fraser CH, Harris C, Keast DH, Potter PJ, et al. Electrical stimulation therapy increases rate of healing of pressure ulcers in community-dwelling people with spinal cord injury. <i>Arch Phys Med Rehabil</i> 2010;91:669-78.
Hurlow J, Bowler PG. Potential implications of biofilm in chronic wounds: a case series. <i>J Wound Care</i> 2012;21:109-119.
Iizaka S, Sanada H, Matsui Y, Furue M, Tachibana T, Nakayama T, et al. Predictive validity of weekly monitoring of wound status using DESIGN-R score change for pressure ulcer healing: A multicenter prospective cohort study. <i>Wound Repair & Regeneration</i> 2012;20:473-481.
Iizaka S, Sanada H, Minematsu T, Oba M, Nakagami G, Koyanagi H, et al. Do nutritional markers in wound fluid reflect pressure ulcer status? <i>Wound Repair & Regeneration</i> 2010;18:31-7.
Iizaka S, Sanada H, Nakagami G, Sekine R, Koyanagi H, Konya C, et al. Estimation of protein loss from wound fluid in older patients with severe pressure ulcers. <i>Nutrition.</i> 2010;26:890-5.
Iordanou P, Baltopoulos G, Giannakopoulou M, Bellou P, Ktenas E. Effect of polarized light in the healing process of pressure ulcers [corrected] [published erratum appears in <i>INT J NURS PRACT</i> 2002 Apr;8(2):116]. <i>Int J Nurs Pract.</i> 2002;8:49-55.

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

Isenberg R. Pulsed radio frequency energy therapy in elderly patients with Stage III and Stage IV pressure ulcers in long-term care settings. <i>Ostomy Wound Management</i> . 2008;54:72-3.
Jones KR, Fennie K. Factors Influencing Pressure Ulcer Healing in Adults Over 50: An Exploratory Study. <i>J Am Med Dir Assoc</i> . 2007;8:378-87.
Kilpadi DV, Stechmiller JK, Childress B, Cowan L, Comerio M, Kieswetter K, et al. Composition of wound fluid from pressure ulcers treated with negative pressure wound therapy using V A C therapy in home health or extended care patients: a pilot study. <i>Wounds: A Compendium of Clinical Research & Practice</i> 2006;18:119-28.
Kloth LC, Berman JE, Dumit-Minkel S, Sutton CH, Papanek PE, Wurzel J. Effects of a normothermic dressing on pressure ulcer healing. <i>Adv Skin Wound Care</i> 2000;13:69-74.
Kloth LC, Berman JE, Nett M, Papanek PE, Dumit-Minkel S. A randomized controlled clinical trial to evaluate the effects of noncontact normothermic wound therapy on chronic full-thickness pressure ulcers. <i>Adv Skin Wound Care</i> . 2002;15:270-6.
Knops SP, Van L, E M, Spanjersberg WR, Patka P, Schipper IB. Randomised clinical trial comparing pressure characteristics of pelvic circumferential compression devices in healthy volunteers. <i>Injury</i> 2011;42:1020-6.
Konya C, Sanada H, Sugama J, Okuwa M, Kitagawa A. Does the use of a cleanser on skin surrounding pressure ulcers in older people promote healing? <i>Journal of wound care</i> . 2005;14:169-71.
Kurring PA, Roberts CD, Quinlan D. Evaluation of a hydrocellular dressing in the management of exuding wounds in the community. <i>British Journal of Nursing</i> . 1994;3:1049-50, 52.
Labanaris AP, Polykandriotis E, Horch RE. The effect of vacuum-assisted closure on lymph vessels in chronic wounds. <i>Journal of Plastic, Reconstructive and Aesthetic Surgery</i> . 2009;62:1068-75.
Lee SK, Posthauer ME, Dorner B, Redovian V, Maloney MJ. Pressure ulcer healing with a concentrated, fortified, collagen protein hydrolysate supplement: a randomized controlled trial. <i>Adv Skin Wound Care</i> . 2006;19:92-6.
Lee Y, Cesario T, McCauley V, Flionis L, Pax A, Thrupp L. Low-level colonization and infection with ciprofloxacin-resistant gram-negative bacilli in a skilled nursing facility. <i>American Journal of Infection Control</i> . 1998;26:552-7.
Lepisto M, Lauri S, Eriksson E, Vahlberg T. Pressure ulcer patients in long-term care. A follow-up study. <i>Reviews in Clinical Gerontology</i> . 2004;14:91-103.
Lindahl E, Gilje F, Norberg A, Söderberg A. Nurses' ethical reflections on caring for people with malodorous exuding ulcers. <i>Nursing Ethics</i> 2010;17:777-90.
Maeshige N, Fujiwara H, Honda H, Yoshikawa Y, Terashi H, Usami M, et al. Evaluation of the combined use of ultrasound irradiation and wound dressing on pressure ulcers. <i>J Wound Care</i>

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

2010;19:63-8.
Malbrain M, Hendriks B, Wijnands P, Denie D, Jans A, Vanpellicom J, et al. A pilot randomised controlled trial comparing reactive air and active alternating pressure mattresses in the prevention and treatment of pressure ulcers among medical ICU patients. <i>J Tissue Viability</i> 2010;19:7-15.
Mao CL, Rivet AJ, Sidora T, Pasko MT. Update on pressure ulcer management and deep tissue injury. <i>Ann Pharmacother</i> 2010;44:325-32.
Marchand AC, Lidowski H. Reassessment of the use of genuine sheepskin for pressure ulcer prevention and treatment. <i>Decubitus: The Journal of Skin Ulcers</i> . 1993;6:44-7.
Martin SJ, Corrado OJ, Kay EA. Enzymatic debridement for necrotic wounds. <i>Journal of wound care</i> . 1996;5:310-1.
McMullen D. Clinical experience with a calcium alginate dressing. <i>Dermatology Nursing</i> . 1991;3:216-9.
Meaume S, Perez J, Rethore V, Sebbane G, Dompmartin A, Bressieux JM, et al. Management of chronic wounds with an innovative absorbent wound dressing. <i>J Wound Care</i> 2012;21:315-6, 318, 320-2.
Meaume S, Vallet D, Morere MN, Teot L. Evaluation of a silver-releasing hydroalginate dressing in chronic wounds with signs of local infection. <i>Journal of wound care</i> . 2005;14:411-9.
Meaume S, Van De L, Heyman H, Romanelli M, Ciangherotti A, Charpin S. A study to compare a new self-adherent soft silicone dressing with a self-adherent polymer dressing in stage II pressure ulcers. <i>Ostomy Wound Manage</i> 2003;49:44-51.
Miyachi Y, Imamura S. Improved healing of pressure sores with sugar and povidone-iodine. <i>Current Therapeutic Research - Clinical and Experimental</i> . 1991;49:81-7.
Mizokami F, Murasawa Y, Furuta K, Isogai Z. Iodoform gauze removes necrotic tissue from pressure ulcer wounds by fibrinolytic activity. <i>Biological and Pharmaceutical Bulletin</i> 2012;35:1048-1053.
Mody GN, Nirmal IA, Duraisamy S, Perakath B. A blinded, prospective, randomized controlled trial of topical negative pressure wound closure in India. <i>Ostomy Wound Management</i> 2008;54:36-46.
Moore ZE, Cowman S. Wound cleansing for pressure ulcers. <i>Cochrane Database of Systematic Reviews</i> 2013.
Moore ZE, Cowman S. Wound cleansing for pressure ulcers. <i>Cochrane Database of Systematic Reviews</i> . 2005;4:4.
Motta G, Dunham L, Dye T, Mentz J, O'Connell-Gifford E, Smith E. Clinical efficacy and cost-effectiveness of a new synthetic polymer sheet wound dressing. <i>Ostomy Wound Manage</i> . 1999;45:41, 4-6, 8.
Motta GJ, Milne CT, Corbett LQ. Impact of antimicrobial gauze on bacterial colonies in wounds that require packing. <i>Ostomy Wound Manage</i> 2004;50:48-62.
Motta GJ. The effectiveness of Dermagran topical therapy for treating chronic wounds in nursing

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

facility residents. <i>Ostomy Wound Management</i> . 1991;36:35-8.
Mulder GD, Altman M, Seeley JE, Tintle T. Prospective randomized study of the efficacy of hydrogel, hydrocolloid, and saline solution -- moistened dressings on the management of pressure ulcers. <i>Wound Repair & Regeneration</i> . 1993;1:213-8.
Münter KC, Beele H, Russell L, Crespi A, Gröchenig E, Basse P, et al. Effect of a sustained silver-releasing dressing on ulcers with delayed healing: the CONTOP study. <i>J Wound Care</i> 2006;15:199-206.
Mustoe TA, Cutler NR, Allman RM, Goode PS, Deuel TF, Prause JA, et al. A phase II study to evaluate recombinant platelet-derived growth factor-BB in the treatment of stage 3 and 4 pressure ulcers. <i>Arch Surg</i> . 1994;129:213-9.
Nagase T, Iizaka S, Kato H, Nakagami G, Kaitani T, Machida M, et al. Undermining incision and healing of deep pressure ulcers: A prospective cohort study of pressure ulcers by the Japanese National Hospital Organization. <i>Wound Repair & Regeneration</i> 2013;21:512-519.
Nakagami G, Sanada H, Iizaka S, Kadono T, Higashino T, Koyanagi H, et al. Predicting delayed pressure ulcer healing using thermography: a prospective cohort study. <i>Journal of wound care</i> . 2010;19:465-6, 8, 70 passim.
Nixon J, Nelson EA, Cranny G, Iglesias CP, Hawkins K, Cullum NA, et al. Pressure relieving support surfaces: a randomised evaluation. <i>Health Technology Assessment</i> . 2006;10:1-180.
Ochs RF, Horn SD, van R, Pietsch C, Smout RJ. Comparison of air-fluidized therapy with other support surfaces used to treat pressure ulcers in nursing home residents. <i>Ostomy Wound Manage</i> . 2005;51:38-68.
Ohura T, Nakajo T, Moriguchi T, Oka H, Tachi M, Ohura, Jr., et al. Clinical efficacy of basic fibroblast growth factor on pressure ulcers: Case-control pairing study using a new evaluation method. <i>Wound Repair and Regeneration</i> . 2011;19:542-51.
Paris A, Horvath R, Basset P, Thiery S, Couturier P, Franco A, et al. Nitrous oxide-oxygen mixture during care of bedsores and painful ulcers in the elderly: a randomized, crossover, open-label pilot study. <i>J Pain Symptom Manage</i> . 2008;35:171-6.
Parish LC, Dryjski M, Cadden S, Versiva XCPUSG. Prospective clinical study of a new adhesive gelling foam dressing in pressure ulcers. <i>Int Wound J</i> . 2008;5:60-7.
Parnell LKS, Ciufi B, Gokoo CF. Preliminary use of a hydrogel containing enzymes in the treatment of Stage II and Stage III pressure ulcers. <i>Ostomy Wound Management</i> 2005;51:50-2.
Payne WG, Posnett J, Alvarez O, Brown-Etris M, Jameson G, Wolcott R, et al. A prospective, randomized clinical trial to assess the cost-effectiveness of a modern foam dressing versus a traditional saline gauze dressing in the treatment of stage II pressure ulcers. <i>Ostomy Wound Management</i> . 2009;55:50-5.

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

Payne WG, Wright TE, Ochs D, Mannari RJ, Robson MC. An exploratory study of dermal replacement therapy in the treatment of stage III pressure ulcers. <i>Journal of Applied Research</i> . 2004;4:12-23.
Price P, Bale S, Crook H, Harding KG. The effect of a radiant heat dressing on pressure ulcers. <i>J Wound Care</i> 2000;9:201-205.
Pullen R, Popp R, Fusgen I. Multicentre evaluation of the topical treatment of decubital ulcers in geriatric patients. <i>European Journal of Geriatrics</i> . 2000;2:27-30.
Pullen R, Popp R, Volkens P, Fusgen I. Prospective randomized double-blind study of the wound-debriding effects of collagenase and fibrinolysin/deoxyribonuclease in pressure ulcers. <i>Age Ageing</i> . 2002;31:126-30.
Raffoul W, Far MS, Cayeux MC, Berger MM. Nutritional status and food intake in nine patients with chronic low-limb ulcers and pressure ulcers: importance of oral supplements. <i>Nutrition</i> 2006;22:82-8.
Rees RS, Robson MC, Smiell JM, Perry BH. Becaplermin gel in the treatment of pressure ulcers: a phase II randomized, double-blind, placebo-controlled study. <i>Wound Repair Regen</i> 1999;7:141-7.
Robicsek A, Beaumont JL, Thomson RB, Jr, Govindarajan G, Peterson LR. Topical therapy for methicillin-resistant <i>Staphylococcus aureus</i> colonization: impact on infection risk. <i>Infection Control & Hospital Epidemiology</i> 2009;30:623-32.
Robson MC, Hill DP, Smith PD, Wang X, Meyer-Siegler K, Ko F, et al. Sequential cytokine therapy for pressure ulcers: Clinical and mechanistic response. <i>Ann Surg</i> 2000;231:600-611.
Robson MC, Phillips LG, Lawrence WT, Bishop JB, Youngerman JS, Hayward PG, et al. The safety and effect of topically applied recombinant basic fibroblast growth factor on the healing of chronic pressure sores. <i>Ann Surg</i> 1992;216:401-6; discussion 406.
Romanelli M, Dimi V, Bertone MS, Mazzatenta C, Martini P. Efficacy and tolerability of "Fitostimoline(registered trademark) antibiotico" soaked gauzes in the topical treatment of cutaneous sores, ulcers and burns, complicated with bacterial contamination. An open-label, controlled, randomised, multicentre, parallel group. <i>Gazzetta Medica Italiana Archivio per le Scienze Mediche</i> 2008;167:251-260.
Romanelli M, Dini V, Barbanera S, Bertone MS. Evaluation of the efficacy and tolerability of a solution containing propyl betaine and polihexanide for wound irrigation. <i>Skin Pharmacol Physiol</i> 2010;23 Suppl:41-4.
Rosenthal MJ, Felton RM, Nastasi AE, Naliboff BD, Harker J, Navach JH. Healing of advanced pressure ulcers by a generic total contact seat: 2 randomized comparisons with low air loss bed treatments. <i>Arch Phys Med Rehabil</i> 2003;84:1733-42.
Rudensky B, Lipschits M, Isaacsohn M, Sonnenblick M. Infected pressure sores: Comparison of methods for bacterial identification. <i>Southern Medical Journal</i> 1992;85:901-903.

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

Saji M, Taguchi S, Uchiyama K, Osono E, Hayama N, Ohkuni H. Efficacy of gentian violet in the eradication of methicillin-resistant <i>Staphylococcus aureus</i> from skin lesions. <i>Journal of Hospital Infection</i> . 1995;31:225-8.
Salzberg CA, Cooper-Vastola SA, Perez F, Viehbeck MG, Byrne DW. The effects of non-thermal pulsed electromagnetic energy on wound healing of pressure ulcers in spinal cord-injured patients: a randomized, double-blind study. <i>Ostomy Wound Management</i> . 1995;41:42-4.
Santangelo A, Testai M, Ossino MC, Barbagallo P, Crisafulli C, Muscara G, et al. Management and treatment of decubital ulcers of an elderly population in the assisted sanitary residence of Futura-Viagrande (Catania, Sicily, Italy). <i>Arch Gerontol Geriatr</i> . 2009;48:332-4.
Sanz-Adell V, Casper-Pedraza, Montse, Gendre-Jornet, M. Isabel Alegria-Alvaro, Patricia. Acupuncture treatment in ulcers. <i>Agora de Enfermeria</i> 2011;15:27-30.
Saydak SJ. A pilot test of two methods for the treatment of pressure ulcers. <i>Journal of Enterostomal Therapy</i> . 1990;17:139-42.
Schessel ES, Ger R. The management of pressure sores by constant-tension approximation. <i>British Journal of Plastic Surgery</i> . 2001;54:439-46.
Scott EM, Leaper DJ, Clark M, Kelly PJ. Effects of warming therapy on pressure ulcers--a randomized trial. <i>AORN J</i> . 2001;73:921-7, 9.
Seaman S, Herbster S, Muglia J, Murray M, Rick C. Simplifying modern wound management for nonprofessional caregivers. <i>Ostomy Wound Manage</i> 2000;46:18-27.
Selkowitz DM, Cameron MH, Mainzer A, Wolfe R. Efficacy of pulsed low-intensity ultrasound in wound healing: a single-case design. <i>Ostomy Wound Manage</i> . 2002;48:40-4, 6.
Serena T, Lee SK, Lam K, Attar P, Meneses P, Ennis W. The impact of noncontact, nonthermal, low-frequency ultrasound on bacterial counts in experimental and chronic wounds. <i>Ostomy Wound Manage</i> . 2009;55:22-30.
Shahin ESM, Dassen T, Halfens RJG. Incidence, prevention and treatment of pressure ulcers in intensive care patients: a longitudinal study. <i>Int J Nurs Stud</i> 2009;46:413-21.
Shaw CG. Seat cushion comparison for nursing home wheelchair users. <i>Assistive Technology</i> 1993;5:92-105.
Sherman RA, Wyle F, Vulpe M. Maggot therapy for treating pressure ulcers in spinal cord injury patients. <i>J Spinal Cord Med</i> . 1995;18:71-4.
Sherman RA. Maggot versus conservative debridement therapy for the treatment of pressure ulcers. <i>Wound Repair & Regeneration</i> . 2002;10:208-14.
Sibbald RG, Coutts P, Woo KY. Reduction of bacterial burden and pain in chronic wounds using a new polyhexamethylene biguanide antimicrobial foam dressing-clinical trial results. <i>Adv Skin</i>

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

Wound Care. 2011;24:78-84.
Singer P. Nutritional care to prevent and heal pressure ulcers. Israel Medical Association Journal. 2002;4:713-6.
Small N, Mulder M, Mackenzie MJ, Nel M. A comparative analysis of pressure sore treatment modalities in community settings. Curationis. 2002;25:74-82.
Sopata M, Luczak J, Ciupinska M. Effect of bacteriological status on pressure ulcer healing in patients with advanced cancer. Journal of wound care. 2002;11:107-10.
St C. Descriptive study of the use of a specialty bed in the United Kingdom. Decubitus: The Journal of Skin Ulcers. 1992;5:28-30.
Takahashi J, Yokota O, Fujisawa Y, Sasaki K, Ishizu H, Aoki T, et al. An evaluation of polyvinylidene film dressing for treatment of pressure ulcers in older people. Journal of wound care. 2006;15:449-50, 52.
Taly AB, Sivaraman N, K P, Murali T, John A. Efficacy of multiwavelength light therapy in the treatment of pressure ulcers in subjects with disorders of the spinal cord: A randomized double-blind controlled trial. Arch Phys Med Rehabil 2004;85:1657-61.
ter R, Kessels AG, Knipschild P. A randomized clinical trial of ultrasound in the treatment of pressure ulcers. Phys Ther. 1996;76:1301-11.
The effect of stochastic electrical noise on hard-to-heal wounds. Journal of wound care. 2011;20:96-103.
Thomas AC, Wysocki AB. The healing wound: a comparison of three clinically useful methods of measurement. Decubitus: The Journal of Skin Ulcers. 1990;3:18-20.
Thomas DR, Diebold MR, Eggemeyer LM. A controlled, randomized, comparative study of a radiant heat bandage on the healing of stage 3-4 pressure ulcers: a pilot study. J Am Med Dir Assoc. 2005;6:46-9.
Thomas DR, Goode PS, LaMaster K, Tennyson T. Acemannan hydrogel dressing versus saline dressing for pressure ulcers. A randomized, controlled trial. Adv Wound Care. 1998;11:273-6.
Thomas DR. Prevention and management of pressure ulcers. Reviews in Clinical Gerontology 2001;11:115-130.
Trial C, Darbas H, Lavigne J, Sotto A, Simoneau G, Tillet Y, et al. Assessment of the antimicrobial effectiveness of a new silver alginate wound dressing: a RCT [corrected] [published erratum appears in J WOUND CARE 2010 Mar;19(3):109]. J Wound Care 2010;19:20-6.
Ullah MO. A study to detect the efficacy of Micro-current Electrical Therapy on decubitus wound. Journal of Medical Sciences 2007;7:1320-1324.
Ultrasound for pressure ulcers. Nurses' Drug Alert 1995;19.

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

Valente SA, Greenough I, W B, DeMarco SL, Andersen RE. More expensive surfaces are not always better. <i>Kuwait Medical Journal</i> . 2012;44:40-5.
van A, R D, Sobotka L, Meijer EP, Heyman H, Groen HW, et al. Specific nutritional support accelerates pressure ulcer healing and reduces wound care intensity in non-malnourished patients. <i>Nutrition</i> . 2010;26:867-72.
van L, Hovius S, Neyens J, Halfens R, Schols J. Pressure relief, cold foam or static air? A single center, prospective, controlled randomized clinical trial in a Dutch nursing home. <i>J Tissue Viability</i> 2011;20:30-4.
van N, C A, Vandenbroucke-Grauls C, van T, F H, Joore HCA, et al. Feasibility and effects of the semirecumbent position to prevent ventilator-associated pneumonia: a randomized study. <i>Crit Care Med</i> . 2006;34:396-402.
Van R. Pressure sores and pressure-decreasing mattresses: controlled clinical trial <i>The Lancet. Ostomy Wound Management</i> 1994;40:12.
Verdú S, Nolasco B, Lopez C, Torra i B. Study on the use of a range of ALLEVYN hydro-cellular dressings in the treatment of bed sores and leg ulcers by primary health care professionals [Spanish]. <i>Revista Rol de Enfermería</i> . 2006;29:43-9.
Wanner MB, Schwarzl F, Strub B, Zaech GA, Pierer G. Vacuum-assisted wound closure for cheaper and more comfortable healing of pressure sores: a prospective study. <i>Scand J Plast Reconstr Surg Hand Surg</i> 2003;37:28-33.
Ward L, Fenton K, Maher L. The high impact actions for nursing and midwifery 4: your skin matters fifth in a series. <i>Nursing Times</i> 2010;106:14-5.
Wild T, Stremitzer S, Budzanowski A, Hoelzenbein T, Ludwig C, Ohrenberger G. Definition of efficiency in vacuum therapy--a randomised controlled trial comparing with V.A.C. Therapy. <i>Int Wound J</i> . 2008;5:641-7.
Winter A, Hewitt H. Testing a hydrocolloid. <i>Nursing Times</i> 1990;86:59-60.
Witkowski W, Jawien A, Witkiewicz W, Zon B. Initial multi-centre observations upon the effect of a new topical negative pressure device upon patient and clinician experience and the treatment of wounds. <i>Int Wound J</i> 2009;6:167-74.
Worsley M. Comparing efficacies treatment of leg ulcers and pressure sores. <i>Nursing Standard</i> . 1991;5:5-6.
Wu T, Wang S-T, Lin P-C, Liu C-L, Chao Yann-Fen C. Effects of Using a High-Density Foam Pad Versus a Viscoelastic Polymer Pad on the Incidence of Pressure Ulcer Development During Spinal Surgery. <i>Biological Research for Nursing</i> 2011;13:419-24.
Xakellis GC, Chrischilles EA. Hydrocolloid versus saline-gauze dressings in treating pressure ulcers:

Exklusionslista Trycksår/Exclusion list Pressure ulcer

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

a cost-effectiveness analysis. Arch Phys Med Rehabil. 1992;73:463-9.
Yamamoto-Mitani N, Katakura N, Fujita J, Shinohara Y, Sonoda Y, Hayashi K. Evaluating pressure ulcer care by home healthcare nurses. International Journal of Older People Nursing 2011;6:201-215.
Yapucu G, Eser I. Effectiveness of a honey dressing for healing pressure ulcers. J Wound Ostomy Continence Nurs 2007;34:184-90.
Yastrub DJ. Relationship between type of treatment and degree of wound healing among institutionalized geriatric patients with stage II pressure ulcers reprinted with permission from the author and Nurse Practitioner Associates for Continuing Education. Care Management Journals. 2004;5:213-8.
Yastrub DJ. Relationship between type of treatment and degree of wound healing among institutionalized geriatric patients with stage II pressure ulcers. Clinical Excellence for Nurse Practitioners. 2005;9:89-94.
Young J, Nikoletti S, McCaul K, Twigg D, Morey P. Risk factors associated with pressure ulcer development at a major western Australian teaching hospital from 1998 to 2000: secondary data analysis. Journal of Wound, Ostomy & Continence Nursing. 2002;29:234-41.
Zeron HM, Krotzsch G, F E, Munoz RE. Pressure ulcers: a pilot study for treatment with collagen polyvinylpyrrolidone. Int J Dermatol. 2007;46:314-7.