

## Exkluderade studier/Excluded studies

### Organisation – (bensår/leg ulcer, fotsår/foot ulcer, trycksår/pressure ulcer)

Nedanstående studier har efter fulltextgranskning inte uppfyllt inklusionskriterierna och ligger således inte till grund för de evidensbaserade resultaten. En och samma studie kan ha förekommit i flera interventioner men redovisas endast en gång.

The following studies have after full text review, not fulfilled the inclusion criteria and do therefore not form the basis for the evidence-based results. A single study might have occurred in several interventions but is only reported once.

<b>Bensår/Leg ulcer</b>
Adam DJ, Naik J, Hartshorne T, Bello M, London NJ. The diagnosis and management of 689 chronic leg ulcers in a single-visit assessment clinic. <i>Eur J Vasc Endovasc Surg</i> 2003;25:462-8.
Apelqvist J, Ragnarson-Tennvall G, Persson U, Larsson J. Diabetic foot ulcers in a multidisciplinary setting. An economic analysis of primary healing and healing with amputation. <i>J Intern Med</i> . 1994;235:463-71.
Bale S. The contribution of the wound care nurse in developing a diabetic foot clinic. <i>British Journal of Clinical Governance</i> . 2002;7:22-6.
Bjellerup M, Akesson H. The role of vascular surgery in chronic leg ulcers: Report from a specialized ulcer clinic. <i>Acta Derm Venereol</i> 2002;82:266-9.
Campbell LV, Graham AR, Kidd RM, Molloy HF, O'Rourke SR, Colagiuri S. The lower limb in people with diabetes. Position statement of the Australian Diabetes Society. <i>Med J Aust</i> . 2000;173:369-72.
Castineira F. The Limerick leg-ulcer project: Early results. <i>Irish Journal of Medical Science</i> 1999;168:17-20.
Chen M-Y, Huang W-C, Peng Y-S, Guo J-S, Chen C-P, Jong M-C, et al. Effectiveness of a health promotion programme for farmers and fishermen with type-2 diabetes in Taiwan. <i>Journal of Advanced Nursing</i> . 2011;67:2060-7.
Ciardullo AV, Daghigh MM, Brunetti M, Bevini M, Daya G, Feltri G, et al. Audit of a shared-care program for persons with diabetes: baseline and 3 annual follow-ups. <i>Acta Diabetol</i> . 2004;41:9-13.
Collins K, Morrell J, Peters J, Walters S, Brereton L. Research Variations in venous leg ulcer care in the community. <i>Br J Community Nurs</i> 1998;3:6-8.
Corbett CF. A randomized pilot study of improving foot care in home health patients with diabetes. <i>Diabetes Educator</i> . 2003;29:273-82.
Edwards H, Courtney M, Finlayson K, Lewis C, Lindsay E, Dumble J. Improved healing rates for chronic venous leg ulcers: pilot study results from a randomized controlled trial of a community nursing intervention. <i>Int J Nurs Pract</i> 2005;11:169-76.
Edwards H, Courtney M, Finlayson K, Shuter P, Lindsay E. A randomised controlled trial of a

community nursing intervention: improved quality of life and healing for clients with chronic leg ulcers. <i>Journal of Clinical Nursing</i> 2009;18:1541-9.
Ellison DA, Hayes L, Lane C, Tracey A, McCollum CN. Evaluating the cost and efficacy of leg ulcer care provided in two large UK health authorities. <i>J Wound Care</i> 2002;11:47-51.
Faglia E, Favales F, Aldeghi A, Calia P, Quarantiello A, Barbano P, et al. Change in major amputation rate in a center dedicated to diabetic foot care during the 1980s: Prognostic determinants for major amputation. <i>J Diabetes Complications</i> 1998;12:96-102.
FitzJohn M. Innovations in practice Treating diabetic foot ulceration in the 21st century. <i>British Journal of Community Health Nursing</i> . 1996;1:22-4.
Fletton JA, Perkins J, Jaap AJ, Brash PD, Tooke JE. Is community chiropodial/podiatric care appropriately targeted at the 'at-risk' diabetic foot? <i>Foot</i> . 1995;5:176-9.
Ghuri ASK, Nyamekye I, Grabs AJ, Farndon JR, Whyman MR, Poskitt KR. Influence of a specialised leg ulcer service and venous surgery on the outcome of venous leg ulcers. <i>European Journal of Vascular and Endovascular Surgery</i> 1998;16:238-244.
Gibbons GW, Marcaccio, Jr., E J, Burgess AM, Pomposelli, Jr., F B, et al. Improved quality of diabetic foot care, 1984 vs 1990: Reduced length of stay and costs, insufficient reimbursement. <i>Archives of Surgery</i> . 1993;128:576-81.
Girod I, Valensi P, Laforet C, Moreau-Defarges T, Guillon P, Baron F. An economic evaluation of the cost of diabetic foot ulcers: Results of a retrospective study on 239 patients. <i>Diabetes and Metabolism</i> . 2003;29:269-77.
Harrison Margaret B, VanDenKerkhof E, Hopman Wilma M, Graham Ian D, Lorimer K, Carley M. Evidence-informed leg ulcer care: a cohort study comparing outcomes of individuals choosing nurse-led clinic or home care. <i>Ostomy Wound Management</i> 2011;57:38-45.
Harrison MB, Graham ID, Lorimer K, Vandenberg E, Buchanan M, Wells PS, et al. Nurse clinic versus home delivery of evidence-based community leg ulcer care: a randomized health services trial. <i>BMC Health Serv Res</i> 2008;8:243.
Heinen MM, Persoon A, van de K, Otero M, van A. Ulcer-related problems and health care needs in patients with venous leg ulceration: a descriptive, cross-sectional study. <i>International Journal of Nursing Studies</i> 2007;44:1296-303.
Lavery LA, Armstrong DG, Wunderlich RP, Tredwell J, Boulton AJ. Predictive value of foot pressure assessment as part of a population-based diabetes disease management program. <i>Diabetes Care</i> . 2003;26:1069-73.
Lavery LA, Wunderlich RP, Tredwell JL. Disease management for the diabetic foot: Effectiveness of a diabetic foot prevention program to reduce amputations and hospitalizations. <i>Diabetes Res Clin Pract</i> . 2005;70:31-7.
Nather A, Bee CS, Lin WK, Valerie CXB, Liang S, Tambyah PA, et al. Value of team approach combined with clinical pathway for diabetic foot problems: A clinical evaluation. <i>Diabetic Foot and Ankle</i> 2010;1.

Nelzen O. Fifty percent reduction in venous ulcer prevalence is achievable - Swedish experience. <i>J Vasc Surg</i> 2010;52:39S-44S.
Pataky Z, Herrmann FR, Regat D, Vuagnat H. The at-risk foot concerns not only patients with diabetes mellitus. <i>Gerontology</i> . 2008;54:349-53.
Prompers L, Huijberts M, Apelqvist J, Jude E, Piaggese A, Bakker K, et al. Delivery of care to diabetic patients with foot ulcers in daily practice: Results of the Eurodiale Study, a prospective cohort study. <i>Diabetic Medicine</i> 2008;25:700-707.
Prompers L, Huijberts M, Apelqvist J, Jude E, Piaggese A, Bakker K, et al. Optimal organization of health care in diabetic foot disease: introduction to the Eurodiale study. <i>International Journal of Lower Extremity Wounds</i> . 2007;6:11-7.
Purwins S, Herberger K, Debus ES, Rustenbach SJ, Pelzer P, Rabe E, et al. Cost-of-illness of chronic leg ulcers in Germany. <i>Int Wound J</i> 2010;7:97-102.
Real JT, Valls M, Ascaso P, Basanta ML, Viguer A, Ascaso JF, et al. Risk factors associated to hospitalization in diabetic patients with foot ulcers. <i>Medicina Clinica</i> . 2001;117:641-4.
Rezende KF, Ferraz MB, Malerbi DA, Melo NH, Nunes MP, Pedrosa HC, et al. Direct costs and outcomes for inpatients with diabetes mellitus and foot ulcers in a developing country: The experience of the public health system of Brazil. <i>Diabetes and Metabolic Syndrome: Clinical Research and Reviews</i> . 2009;3:228-32.
Ronnemaa T, Hamalainen H, Toikka T, Liukkonen I. Evaluation of the impact of podiatrist care in the primary prevention of foot problems diabetic subjects. <i>Diabetes Care</i> 1997;20:1833-1837.
Ruckley CV. Towards a model of care for chronic leg ulcer: Lessons learned from studies in Scotland. <i>Phlebology</i> 2001;16:17-19.
Scriven JM, Hartshorne T, Bell PRF, Naylor AR, London NJM. Single-visit venous ulcer assessment clinic: The first year. <i>British Journal of Surgery</i> 1997;84:334-336.
Sturmberg JP, Overend D. General practice based diabetes clinics An integration model. <i>Australian Family Physician</i> . 1999;28:240-5.
Öien RF. Registering Ulcer Treatment through a national quality register: RUT -- a winning concept for both patients and the health care sector. <i>EWMA Journal</i> 2009;9:41-4.

<b>Fotsår/foot ulcer</b>
Apelqvist J, Ragnarson-Tennvall G, Persson U, Larsson J. Diabetic foot ulcers in a multidisciplinary setting. An economic analysis of primary healing and healing with amputation. <i>J Intern Med</i> 1994;235:463-71.
Bale S. The contribution of the wound care nurse in developing a diabetic foot clinic. <i>British Journal of Clinical Governance</i> 2002;7:22-26.
Campbell LV, Graham AR, Kidd RM, Molloy HF, O'Rourke SR, Colagiuri S. The lower limb in people with diabetes. Position statement of the Australian Diabetes Society. <i>Med J Aust</i> 2000;173:369-72.
Chen M-Y, Huang W-C, Peng Y-S, Guo J-S, Chen C-P, Jong M-C, et al. Effectiveness of a health promotion programme for farmers and fishermen with type-2 diabetes in Taiwan. <i>Journal of Advanced Nursing</i> 2011;67:2060-7.
Ciardullo AV, Daghio MM, Brunetti M, Bevini M, Daya G, Feltri G, et al. Audit of a shared-care program for persons with diabetes: baseline and 3 annual follow-ups. <i>Acta Diabetol</i> 2004;41:9-13.
Corbett CF. A randomized pilot study of improving foot care in home health patients with diabetes. <i>Diabetes Educator</i> 2003;29:273-82.
FitzJohn M. Innovations in practice Treating diabetic foot ulceration in the 21st century. <i>British Journal of Community Health Nursing</i> 1996;1:22-4.
Fletton JA, Perkins J, Jaap AJ, Brash PD, Tooke JE. Is community chiropodial/podiatric care appropriately targeted at the 'at-risk' diabetic foot? <i>Foot</i> 1995;5:176-179.
Gibbons GW, Marcaccio, Jr., E J, Burgess AM, Pomposelli, Jr., F B, et al. Improved quality of diabetic foot care, 1984 vs 1990: Reduced length of stay and costs, insufficient reimbursement. <i>Archives of Surgery</i> 1993;128:576-581.
Lavery LA, Armstrong DG, Wunderlich RP, Tredwell J, Boulton AJ. Predictive value of foot pressure assessment as part of a population-based diabetes disease management program. <i>Diabetes Care</i> 2003;26:1069-73.
Lavery LA, Wunderlich RP, Tredwell JL. Disease management for the diabetic foot: Effectiveness of a diabetic foot prevention program to reduce amputations and hospitalizations. <i>Diabetes Res Clin Pract</i> 2005;70:31-37.
Pataky Z, Herrmann FR, Regat D, Vuagnat H. The at-risk foot concerns not only patients with diabetes mellitus. <i>Gerontology</i> 2008;54:349-353.
Prompers L, Huijberts M, Apelqvist J, Jude E, Piaggese A, Bakker K, et al. Optimal organization of health care in diabetic foot disease: introduction to the Eurodiale study. <i>International Journal of Lower Extremity Wounds</i> 2007;6:11-7.
Real JT, Valls M, Ascaso P, Basanta ML, Viguer A, Ascaso JF, et al. Risk factors associated to hospitalization in diabetic patients with foot ulcers. <i>Medicina Clinica</i> 2001;117:641-644.

<p>Rezende KF, Ferraz MB, Malerbi DA, Melo NH, Nunes MP, Pedrosa HC, et al. Direct costs and outcomes for inpatients with diabetes mellitus and foot ulcers in a developing country: The experience of the public health system of Brazil. <i>Diabetes and Metabolic Syndrome: Clinical Research and Reviews</i> 2009;3:228-232.</p>
<p>Sturmberg JP, Overend D. General practice based diabetes clinics An integration model. <i>Australian Family Physician</i> 1999;28:240-5.</p>
<p><b>Trycksår/Pressure ulcer</b></p>
<p>Arnold A, McEntagart M, Younger DS. Psychosocial issues that face patients with Charcot-Marie-Tooth disease: the role of genetic counseling. <i>J Genet Couns.</i> 2005;14:307-18.</p>
<p>Bjellerup M, Akesson H. The role of vascular surgery in chronic leg ulcers: Report from a specialized ulcer clinic. <i>Acta Derm Venereol.</i> 2002;82:266-9.</p>
<p>Brink P, Smith TF, Linkewich B. Factors associated with pressure ulcers in palliative home care. <i>J Palliat Med.</i> 2006;9:1369-75.</p>
<p>Brooks J, Ersser SJ, Lloyd A, Ryan TJ. Nurse-led education sets out to improve patient concordance and prevent recurrence of leg ulcers. <i>Journal of wound care.</i> 2004;13:111-6.</p>
<p>Burnside J, Wells W, Smith M, Game F, Jeffcoate W, Radford K. Education for the prevention of foot ulcers: views of those with neuropathy. <i>Diabetic Foot Journal.</i> 2007;10.</p>
<p>Burt KM. The relationship between nurse caring and selected outcomes of care in hospitalized older adults. <i>Catholic University of America.</i> 2007;190.</p>
<p>Cai S, Mukamel DB, Temkin-Greener H. Pressure ulcer prevalence among black and white nursing home residents in new york state: Evidence of racial disparity? <i>Medical Care.</i> 2010;48:233-9.</p>
<p>Chang C, Jackson SS, Bullman TA, Cobbs EL. Impact of a home-based primary care program in an urban Veterans Affairs medical center. <i>J Am Med Dir Assoc.</i> 2009;10:133-7.</p>
<p>Clemensen J, Larsen SB, Ejksjaer N. Telemedical treatment at home of diabetic foot ulcers. <i>J Telemed Telecare.</i> 2005;11 Suppl 2:S14-6.</p>
<p>Coggrave MJ, Rose LS. A specialist seating assessment clinic: changing pressure relief practice. <i>Spinal Cord.</i> 2003;41:692-5.</p>
<p>Edwards LM. Compression therapy Why patients do not comply with compression bandaging. <i>British Journal of Nursing.</i> 12:Tissue Viability Supplement S5-6, S8, S10 passim.</p>
<p>Ferrell BA, Josephson K, Norvid P, Alcorn H. Pressure ulcers among patients admitted to home care. <i>J Am Geriatr Soc.</i> 2000;48:1042-7.</p>
<p>Frantz RA, Gardner S, Harvey P, Specht J. Adoption of research-based practice for treatment of pressure ulcers in long-term care. <i>Decubitus: The Journal of Skin Ulcers.</i> 1992;5:44-5.</p>
<p>Garber SL, Rintala DH, Ann H, Rodriguez GP, Friedman J. A structured educational model to improve pressure ulcer prevention knowledge in veterans with spinal cord dysfunction. <i>J Rehabil</i></p>

Res Dev. 2002;39:575-87.
Gengenbacher M, Stahelin HB, Scholer A, Seiler WO. Low biochemical nutritional parameters in acutely ill hospitalized elderly patients with and without stage III to IV pressure ulcers. <i>Aging - Clinical and Experimental Research</i> . 2002;14:420-3.
Gray DG, Cooper PJ, Campbell M. A study of the performance of a pressure reducing foam mattress after three years of use. <i>J Tissue Viability</i> . 1998;8:9-13.
Green SM, Winterberg H, Franks PJ, Moffatt CJ, Eberhardie C, McLaren S. Nutritional intake in community patients with pressure ulcers. <i>Journal of wound care</i> . 1999;8:325-30.
Gunningberg L, Stotts NA, Idvall E. Hospital-acquired pressure ulcers in two Swedish County Councils: Cross-sectional data as the foundation for future quality improvement. <i>Int Wound J</i> . 2011;8:465-73.
Haleem S, Heinert G, Parker MJ. Pressure sores and hip fractures. <i>Injury</i> . 2008;39:219-23.
Hayashi T, Narita Y, Okugawa N, Hamaguchi E, Shibahara M, Kuzuhara S. Pressure ulcers in ALS patients on admission at a university hospital in Japan. <i>Amyotrophic Lateral Sclerosis</i> . 2007;8:310-3.
Hengstermann S, Fischer A, Steinhagen-Thiessen E, Schulz RJ. Nutrition status and pressure ulcer: What we need for nutrition screening. <i>Journal of Parenteral and Enteral Nutrition</i> . 2007;31:288-94.
Hommel A, Bjorkelund KB, Thorngren KG, Ulander K. Nutritional status among patients with hip fracture in relation to pressure ulcers. <i>Clinical Nutrition</i> . 2007;26:589-96.
Iizaka S, Okuwa M, Sugama J, Sanada H. The impact of malnutrition and nutrition-related factors on the development and severity of pressure ulcers in older patients receiving home care. <i>Clinical Nutrition</i> . 2010;29:47-53.
Indredavik B, Rohweder G, Naalsund E, Lydersen S. Medical complications in a comprehensive stroke unit and an early supported discharge service. <i>Stroke</i> . 39:414-20.
Isaia G, Bo M, Tibaldi V, Scarafiotti C, Ruatta C, Michelis G, et al. The experience of a geriatric hospital at home service for acutely ill elderly patients with pressure ulcers: an observational study. <i>Wounds: A Compendium of Clinical Research &amp; Practice</i> . 2010;22:121-6.
Iwata M, Kuzuya M, Kitagawa Y, Suzuki Y, Iguchi A. Underappreciated predictors for postdischarge mortality in acute hospitalized oldest-old patients. <i>Gerontology</i> . 2006;52:92-8.
Keelaghan E, Margolis D, Zhan M, Baumgarten M. Prevalence of pressure ulcers on hospital admission among nursing home residents transferred to the hospital. <i>Wound Repair and Regeneration</i> . 2008;16:331-6.
Kletzenbauer SP, Blakemore ME. Fast track hip trial: an evaluation of a change in X-ray referral. <i>Journal of Clinical Effectiveness</i> . 1996;1:55-6.
Krichbaum K, Pearson V, Savik K, Mueller C. Improving resident outcomes with GAPN organization level interventions gerontological advanced practice nurses. <i>Western Journal of</i>

Nursing Research. 2005;27:322-37.
Kwong Enid WY, Lau Ada TY, Lee Rainbow LP, Kwan Rick YC. A pressure ulcer prevention programme specially designed for nursing homes: Does it work? J Clin Nurs. 2011;20:2777-86.
Landi F, Onder G, Russo A, Bernabei R. Pressure ulcer and mortality in frail elderly people living in community. Arch Gerontol Geriatr. 2007;44 Suppl 1:217-23.
Lang NM, Kraegel JM, Rantz MJ, Krejci JW. Quality of health care for older people in America: a review of nursing studies. American Nurses' Association. 1990;131.
Langer G, Schloemer G, Knerr A, Kuss O, Behrens J. Nutritional interventions for preventing and treating pressure ulcers. Cochrane Database Syst Rev. 2003:CD003216.
Lilly CM, Cody S, Zhao H, Landry K, Baker SP, McIlwaine J, et al. Hospital mortality, length of stay, and preventable complications among critically ill patients before and after tele-ICU reengineering of critical care processes. JAMA: Journal of the American Medical Association. 2011;305:2175-83.
Marek KD, Popejoy L, Petroski G, Mehr D, Rantz M, Lin W. Clinical outcomes of aging in place. Nursing Research. 2005;54:202-11.
Maugham L, Cox R, Amsters D, Battistutta D. Reducing inpatient hospital usage for management of pressure sores after spinal cord lesions. International Journal of Rehabilitation Research. 2004;27:311-5.
Maylor M. Senior nurses' control expectations and the development of pressure ulcers. Nurs Stand. 2001;15:33-7.
McNaughton V, Brazil K. Wound and skin team: impact on pressure ulcer prevalence in chronic care. J Gerontol Nurs. 1995;21:45-9.
Miyazaki Margareth Y, Caliri Maria Helena L, dos S, Claudia B. Knowledge on pressure ulcer prevention among nursing professionals. Revista Latino-Americana de Enfermagem. 2010;18:1203-11.
Morrow JB, Goetz LL, Marcee AK, Snell LE, Kowalske KJ, Roaten SP. Providing access to specialty care for severely disabled institutional residents: an academic health center telemedicine model. Journal of Health Care for the Poor & Underserved. 2009;20:415-22.
Newton M. Integrated care pathway: the prevention and management of pressure ulcers. J Tissue Viability. 2003;13:126-9.
Nixon J, Cranny G, Bond S. Pathology, diagnosis, and classification of pressure ulcers: comparing clinical and imaging techniques. Wound Repair Regen. 2005;13:365-72.
O'Hare J. Maintaining standards for pressure ulcer management in care homes. Nurs Stand. 2008;22:62-4, 6, 8.
Ohura T, Nakajo T, Okada S, Omura K, Adachi K. Evaluation of effects of nutrition intervention on

healing of pressure ulcers and nutritional states (randomized controlled trial). <i>Wound Repair and Regeneration</i> 2011;19:330-336.
Paquay L, Verstraete S, Wouters R, Buntinx F, Vanderwee K, Defloor T, et al. Implementation of a guideline for pressure ulcer prevention in home care: pretest-post-test study. <i>J Clin Nurs</i> . 2010;19:1803-11.
Pham B, Teague L, Mahoney J, Goodman L, Paulden M, Poss J, et al. Early prevention of pressure ulcers among elderly patients admitted through emergency departments: a cost-effectiveness analysis. <i>Ann Emerg Med</i> . 2011;58:468-78 e3.
Rafter L, Reynolds T. Using hue, saturation and intensity compared with three other pressure ulcer grading systems. <i>Wounds UK</i> . 2008;4:25-38.
Rodrigues I, Mégie MF. Prevalence of chronic wounds in Quebec home care: an exploratory study. <i>Ostomy Wound Management</i> . 2006;52:46-8.
Sae-Sia W, Wipke-Tevis D. Pressure ulcer prevention and treatment practices in inpatient rehabilitation facilities. <i>Rehabilitation Nursing</i> . 2002;27:192-8.
Schoonhoven L, Haalboom JRE, Bousema MT, Algra A, Grobbee DE, Grypdonck MH, et al. Prospective cohort study of routine use of risk assessment scales for prediction of pressure ulcers. <i>British Medical Journal</i> . 2002;325:797-800.
Strauss MJ, Gong J, Gary BD, Kalsbeek WD, Spear S. The cost of home air-fluidized therapy for pressure sores. A randomized controlled trial. <i>J Fam Pract</i> . 1991;33:52-9.
Thomas DR, Goode PS, Tarquine PH, Allman RM. Hospital-acquired pressure ulcers and risk of death. <i>J Am Geriatr Soc</i> . 1996;44:1435-40.
Thomas DR. Are all pressure ulcers avoidable? <i>J Am Med Dir Assoc</i> . 2001;2:297-301.
van L, M A, Post MWM, Bloemen JHA, van A, F WA. Care needs of persons with long-term spinal cord injury living at home in the Netherlands. <i>Spinal Cord</i> . 2010;48:423-8.