

Bilaga 3. Förteckning/referenslista över exkluderade studier och studier med låga bevisvärden

3.1 Multimodal rehabilitering

Studier som granskats men som inte uppfyllde inklusionskriterierna

United Kingdom back pain exercise and manipulation (UK BEAM) randomised trial: effectiveness of physical treatments for back pain in primary care. *BMJ* 2004;329:1377.

United Kingdom back pain exercise and manipulation (UK BEAM) randomised trial: cost effectiveness of physical treatments for back pain in primary care. *BMJ* 2004;329:1381.

Almeida TF, Roizenblatt S, Benedito-Silva AA, Tufik S. The effect of combined therapy (ultrasound and interferential current) on pain and sleep in fibromyalgia. *Pain* 2003;104:665-72.

Abasolo L, Carmona L, Hernandez-Garcia C, Lajas C, Loza E, Blanco M, et al. Musculoskeletal work disability for clinicians: time course and effectiveness of a specialized intervention program by diagnosis. *Arthritis Rheum* 2007;57:335-42.

Altan L, Bingol U, Aykac M, Koc Z, Yurtkuran M. Investigation of the effects of pool-based exercise on fibromyalgia syndrome. *Rheumatol Int* 2004;24:272-7.

Andersen LL, Kjaer M, Sogaard K, Hansen L, Kryger AI, Sjogaard G. Effect of two

contrasting types of physical exercise on chronic neck muscle pain. *Arthritis Rheum* 2008;59:84-91.

Anema JR, Steenstra IA, Bongers PM, de Vet HC, Knol DL, Loisel P, et al. Multidisciplinary rehabilitation for subacute low back pain: graded activity or workplace intervention or both? A randomized controlled trial. *Spine* 2007;32:291-8; discussion 299-300.

Ansari NN, Ebadi S, Talebian S, Naghdi S, Mazaheri H, Olyaei G, et al. A randomized, single blind placebo controlled clinical trial on the effect of continuous ultrasound on low back pain. *Electromyogr Clin Neurophysiol* 2006;46:329-36.

Asenlof P, Denison E, Lindberg P. Individually tailored treatment targeting activity, motor behavior, and cognition reduces pain-related disability: a randomized controlled trial in patients with musculoskeletal pain. *J Pain* 2005;6:588-603.

Aure OF, Nilsen JH, Vasseljen O. Manual therapy and exercise therapy in patients with chronic low back pain: a randomized, controlled trial with 1-year follow-up. *Spine* 2003;28:525-31; discussion 531-2.

Babu AS, Mathew E, Danda D, Prakash H. Management of patients with fibromyalgia using biofeedback: a randomized control trial. *Indian J Med Sci* 2007;61:455-61.

- Bastiaenen CH, de Bie RA, Vlaeyen JW, Goossens ME, Leffers P, Wolters PM, et al. Long-term effectiveness and costs of a brief self-management intervention in women with pregnancy-related low back pain after delivery. *BMC Pregnancy Childbirth* 2008;8:19.
- Bekkering GE, van Tulder MW, Hendriks EJ, Koopmanschap MA, Knol DL, Bouter LM, et al. Implementation of clinical guidelines on physical therapy for patients with low back pain: randomized trial comparing patient outcomes after a standard and active implementation strategy. *Phys Ther* 2005;85:544-55.
- Bergman GJ, Winters JC, Groenier KH, Pool JJ, Meyboom-de Jong B, Postema K, et al. Manipulative therapy in addition to usual medical care for patients with shoulder dysfunction and pain: a randomized, controlled trial. *Ann Intern Med* 2004;141:432-9.
- Bernaards CM, Ariens GA, Hildebrandt VH. The (cost-)effectiveness of a lifestyle physical activity intervention in addition to a work style intervention on the recovery from neck and upper limb symptoms in computer workers. *BMC Musculoskeletal Disord* 2006;7:80.
- Blangsted AK, Sogaard K, Hansen EA, Hannerz H, Sjogaard G. One-year randomized controlled trial with different physical-activity programs to reduce musculoskeletal symptoms in the neck and shoulders among office workers. *Scand J Work Environ Health* 2008; 34:55-65.
- Browder DA, Childs JD, Cleland JA, Fritz JM. Effectiveness of an extension-oriented treatment approach in a subgroup of subjects with low back pain: a randomized clinical trial. *Phys Ther* 2007;87:1608-18; discussion 1577-9.
- Brox JI, Reikeras O, Nygaard O, Sorensen R, Indahl A, Holm I, et al. Lumbar instrumented fusion compared with cognitive intervention and exercises in patients with chronic back pain after previous surgery for disc herniation: a prospective randomized controlled study. *Pain* 2006;122:145-55.
- Buchner M, Zuhlten-Hinguranage A, Schiltenswolf M, Neubauer E. Therapy outcome after multidisciplinary treatment for chronic neck and chronic low back pain: a prospective clinical study in 365 patients. *Scand J Rheumatol* 2006;35: 363-7.
- Cairns MC, Foster NE, Wright C. Randomized controlled trial of specific spinal stabilization exercises and conventional physiotherapy for recurrent low back pain. *Spine* 2006;31:E670-81.
- Cambron JA, Gudavalli MR, Hedeker D, McGregor M, Jedlicka J, Keenum M, et al. One-year follow-up of a randomized clinical trial comparing flexion distraction with an exercise program for chronic low-back pain. *J Altern Complement Med* 2006;12:659-68.
- Carr JL, Klaber Moffett JA, Howarth E, Richmond SJ, Torgerson DJ, Jackson DA, et al. A randomized trial comparing a group exercise programme for back pain patients with individual physiotherapy in a severely deprived area. *Disabil Rehabil* 2005;27:929-37.
- Castel A, Perez M, Sala J, Padrol A, Rull M. Effect of hypnotic suggestion on fibromyalgic pain: comparison between hypnosis and relaxation. *Eur J Pain* 2007; 11:463-8.

- Celestini M, Marchese A, Serenelli A, Graziani G. A randomized controlled trial on the efficacy of physical exercise in patients braced for instability of the lumbar spine. *Eura Medicophys* 2005; 41:223-31.
- Chatzitheodorou D, Kabitsis C, Malliou P, Mougios V. A pilot study of the effects of high-intensity aerobic exercise versus passive interventions on pain, disability, psychological strain, and serum cortisol concentrations in people with chronic low back pain. *Phys Ther* 2007;87:304-12.
- Chatzitheodorou D, Mavromoustakos S, Milioti S. The effect of exercise on adrenocortical responsiveness of patients with chronic low back pain, controlled for psychological strain. *Clin Rehabil* 2008;22:319-28.
- Childs JD, Fritz JM, Flynn TW, Irrgang JJ, Johnson KK, Majkowski GR, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med* 2004;141:920-8.
- Chiu TT, Hui-Chan CW, Chein G. A randomized clinical trial of TENS and exercise for patients with chronic neck pain. *Clin Rehabil* 2005;19:850-60.
- Christensen FB, Laurberg I, Bunge CE. Importance of the back-café concept to rehabilitation after lumbar spinal fusion: a randomized clinical study with a 2-year follow-up. *Spine* 2003;28:2561-9.
- Critchley DJ, Ratcliffe J, Noonan S, Jones RH, Hurley MV. Effectiveness and cost-effectiveness of three types of physiotherapy used to reduce chronic low back pain disability: a pragmatic randomized trial with economic evaluation. *Spine* 2007;32:1474-81.
- Dalichau S, Stein B, Schafer K, Buhlmann JJ, Menken P. [Ultrasonic Quantification of spinal configuration and postural capacity for evaluation of different muscle strengthening programs in the therapy of back pain]. *Z Orthop Ihre Grenzgeb* 2005;143:79-85.
- de Buck PD, Breedveld J, van der Giesen FJ, Vlieland TP. A multidisciplinary job retention vocational rehabilitation programme for patients with chronic rheumatic diseases: patients' and occupational physicians' satisfaction. *Ann Rheum Dis* 2004;63:562-8.
- de Buck PD, le Cessie S, van den Hout WB, Peeters AJ, Runday HK, Westedt ML, et al. Randomized comparison of a multidisciplinary job-retention vocational rehabilitation program with usual outpatient care in patients with chronic arthritis at risk for job loss. *Arthritis Rheum* 2005;53:682-90.
- Djavid GE, Mehrdad R, Ghasemi M, Hasan-Zadeh H, Sotoodeh-Manesh A, Pouryaghoub G. In chronic low back pain, low level laser therapy combined with exercise is more beneficial than exercise alone in the long term: a randomised trial. *Aust J Physiother* 2007;53:155-60.
- Donmez A, Karagulle MZ, Tercan N, Dinler M, Issever H, Karagulle M, et al. SPA therapy in fibromyalgia: a randomised controlled clinic study. *Rheumatol Int* 2005;26:168-72.
- Donta ST, Clauw DJ, Engel CC, Jr, Guarino P, Peduzzi P, Williams DA, et al. Cognitive behavioral therapy and aerobic exercise for Gulf War veterans' illnesses:

- a randomized controlled trial. *JAMA* 2003;289:1396-404.
- Donzelli S, Di Domenica E, Cova AM, Galletti R, Giunta N. Two different techniques in the rehabilitation treatment of low back pain: a randomized controlled trial. *Eura Medicophys* 2006;42:205-10.
- Dziedzic K, Hill J, Lewis M, Sim J, Daniels J, Hay EM. Effectiveness of manual therapy or pulsed shortwave diathermy in addition to advice and exercise for neck disorders: a pragmatic randomized controlled trial in physical therapy clinics. *Arthritis Rheum* 2005; 53:214-22.
- Ektor-Andersen J, Ingvarsson E, Kullendorff M, Orbaek P. High cost-benefit of early team-based biomedical and cognitive-behaviour intervention for long-term pain-related sickness absence. *J Rehabil Med* 2008;40:1-8.
- Erdogmus CB, Resch KL, Sabitzer R, Muller H, Nuhr M, Schoggl A, et al. Physiotherapy-based rehabilitation following disc herniation operation: results of a randomized clinical trial. *Spine* 2007; 32:2041-9.
- Fairbank J, Frost H, Wilson-MacDonald J, Yu LM, Barker K, Collins R. Randomised controlled trial to compare surgical stabilisation of the lumbar spine with an intensive rehabilitation programme for patients with chronic low back pain: the MRC spine stabilisation trial. *BMJ* 2005;330:1233.
- Ferreira ML, Ferreira PH, Latimer J, Herbert RD, Hodges PW, Jennings MD, et al. Comparison of general exercise, motor control exercise and spinal manipulative therapy for chronic low back pain: A randomized trial. *Pain* 2007;131:31-7.
- Fioravanti A, Perpignano G, Tirri G, Cardinale G, Gianniti C, Lanza CE, et al. Effects of mud-bath treatment on fibromyalgia patients: a randomized clinical trial. *Rheumatol Int* 2007;27: 1157-61.
- Frey Law LA, Evans S, Knudtson J, Nus S, Scholl K, Sluka KA. Massage reduces pain perception and hyperalgesia in experimental muscle pain: a randomized, controlled trial. *J Pain* 2008;9:714-21.
- Friedrich M, Gittler G, Arendasy M, Friedrich KM. Long-term effect of a combined exercise and motivational program on the level of disability of patients with chronic low back pain. *Spine* 2005;30: 995-1000.
- Frost H, Lamb SE, Doll HA, Carver PT, Stewart-Brown S. Randomised controlled trial of physiotherapy compared with advice for low back pain. *BMJ* 2004;329:708.
- Frost P, Haahr JP, Andersen JH. Reduction of pain-related disability in working populations: a randomized intervention study of the effects of an educational booklet addressing psychosocial risk factors and screening workplaces for physical health hazards. *Spine* 2007; 32:1949-54.
- Galantino ML, Bzdewka TM, Eissler-Russo JL, Holbrook ML, Mogck EP, Geigle P, et al. The impact of modified Hatha yoga on chronic low back pain: a pilot study. *Altern Ther Health Med* 2004;10:56-9.
- Geisser ME, Wiggert EA, Haig AJ, Colwell MO. A randomized, controlled trial of manual therapy and specific adjuvant exercise for chronic low back pain. *Clin J Pain* 2005;21:463-70.

- George SZ, Fritz JM, Bialosky JE, Donald DA. The effect of a fear-avoidance-based physical therapy intervention for patients with acute low back pain: results of a randomized clinical trial. *Spine* 2003;28:2551-60.
- George SZ, Fritz JM, Childs JD. Investigation of elevated fear-avoidance beliefs for patients with low back pain: a secondary analysis involving patients enrolled in physical therapy clinical trials. *J Orthop Sports Phys Ther* 2008;38:50-8.
- Geraets JJ, Goossens ME, de Groot IJ, de Bruijn CP, de Bie RA, Dinant GJ, et al. Effectiveness of a graded exercise therapy program for patients with chronic shoulder complaints. *Aust J Physiother* 2005;51:87-94.
- Ginn KA, Cohen ML. Exercise therapy for shoulder pain aimed at restoring neuromuscular control: a randomized comparative clinical trial. *J Rehabil Med* 2005;37:115-22.
- Godges JJ, Anger MA, Zimmerman G, Delitto A. Effects of education on return-to-work status for people with fear-avoidance beliefs and acute low back pain. *Phys Ther* 2008;88:231-9.
- Goldby LJ, Moore AP, Doust J, Trew ME. A randomized controlled trial investigating the efficiency of musculo-skeletal physiotherapy on chronic low back disorder. *Spine* 2006;31:1083-93.
- Graham N, Gross AR, Goldsmith C. Mechanical traction for mechanical neck disorders: a systematic review. *J Rehabil Med* 2006;38:145-52.
- Gross AR, Hoving JL, Haines TA, Goldsmith CH, Kay T, Aker P, et al. A Cochrane review of manipulation and mobilization for mechanical neck disorders. *Spine* 2004;29:1541-8.
- Gur A, Karakoc M, Cevik R, Nas K, Sarac AJ. Efficacy of low power laser therapy and exercise on pain and functions in chronic low back pain. *Lasers Surg Med* 2003;32:233-8.
- Gusi N, Tomas-Carus P, Hakkinen A, Hakkinen K, Ortega-Alonso A. Exercise in waist-high warm water decreases pain and improves health-related quality of life and strength in the lower extremities in women with fibromyalgia. *Arthritis Rheum* 2006;55:66-73.
- Gustavsson C, von Koch L. Applied relaxation in the treatment of long-lasting neck pain: a randomized controlled pilot study. *J Rehabil Med* 2006;38:100-7.
- Haas M, Grouppe E, Kraemer DF. Dose-response for chiropractic care of chronic low back pain. *Spine J* 2004;4:574-83.
- Hakkinen A, Salo P, Tarvainen U, Wiren K, Ylinen J. Effect of manual therapy and stretching on neck muscle strength and mobility in chronic neck pain. *J Rehabil Med* 2007;39:575-9.
- Harts CC, Helmhout PH, de Bie RA, Staal JB. A high-intensity lumbar extensor strengthening program is little better than a low-intensity program or a waiting list control group for chronic low back pain: a randomised clinical trial. *Aust J Physiother* 2008;54:23-31.
- Hasson D, Arnetz B, Jelveus L, Edelstam B. A randomized clinical trial of the treatment effects of massage compared to relaxation tape recordings on diffuse long-

- term pain. *Psychother Psychosom* 2004; 73:17-24.
- Hayden JA, van Tulder MW, Malmivaara AV, Koes BW. Meta-analysis: exercise therapy for nonspecific low back pain. *Ann Intern Med* 2005;142:765-75.
- Hayden JA, van Tulder MW, Tomlinson G. Systematic review: strategies for using exercise therapy to improve outcomes in chronic low back pain. *Ann Intern Med* 2005;142:776-85.
- Helewa A, Goldsmith CH, Smythe HA, Lee P, Obright K, Stitt L. Effect of therapeutic exercise and sleeping neck support on patients with chronic neck pain: a randomized clinical trial. *J Rheumatol* 2007;34:151-8.
- Helmhout PH, Harts CC, Staal JB, Candel MJ, de Bie RA. Comparison of a high-intensity and a low-intensity lumbar extensor training program as minimal intervention treatment in low back pain: a randomized trial. *Eur Spine J* 2004;13:537-47.
- Hemmila HM. Bone setting for prolonged neck pain: a randomized clinical trial. *J Manipulative Physiol Ther* 2005; 28:508-15.
- Heymans MW, de Vet HC, Bongers PM, Knol DL, Koes BW, van Mechelen W. The effectiveness of high-intensity versus low-intensity back schools in an occupational setting: a pragmatic randomized controlled trial. *Spine* 2006;31:1075-82.
- Hoving JL, de Vet HC, Koes BW, Mameren H, Deville WL, van der Windt DA, et al. Manual therapy, physical therapy, or continued care by the general practitioner for patients with neck pain: long-term results from a pragmatic randomized clinical trial. *Clin J Pain* 2006;22:370-7.
- Hsieh LL, Kuo CH, Lee LH, Yen AM, Chien KL, Chen TH. Treatment of low back pain by acupressure and physical therapy: randomised controlled trial. *BMJ* 2006;332:696-700.
- Hsieh LL, Kuo CH, Yen MF, Chen TH. A randomized controlled clinical trial for low back pain treated by acupressure and physical therapy. *Prev Med* 2004; 39:168-76.
- Hurwitz EL, Morgenstern H, Chiao C. Effects of recreational physical activity and back exercises on low back pain and psychological distress: findings from the UCLA Low Back Pain Study. *Am J Public Health* 2005;95:1817-24.
- Hurwitz EL, Morgenstern H, Kominski GF, Yu F, Chiang LM. A randomized trial of chiropractic and medical care for patients with low back pain: eighteen-month follow-up outcomes from the UCLA low back pain study. *Spine* 2006;31:611-21; discussion 622.
- Johnson RE, Jones GT, Wiles NJ, Chaddock C, Potter RG, Roberts C, et al. Active exercise, education, and cognitive behavioral therapy for persistent disabling low back pain: a randomized controlled trial. *Spine* 2007; 32:1578-85.
- Jones TA. Rolfing. *Phys Med Rehabil Clin N Am* 2004;15:799-809, vi.
- Jousset N, Fanello S, Bontoux L, Dubus V, Billabert C, Vielle B, et al. Effects of functional restoration versus 3 hours per

- week physical therapy: a randomized controlled study. *Spine* 2004;29:487-93; discussion 494.
- Kaapa EH, Frantsi K, Sarna S, Malmivaara A. Multidisciplinary group rehabilitation versus individual physiotherapy for chronic nonspecific low back pain: a randomized trial. *Spine* 2006;31:371-6.
- Karjalainen K, Malmivaara A, van Tulder M, Roine R, Jauhiainen M, Hurri H, et al. Multidisciplinary rehabilitation for fibromyalgia and musculoskeletal pain in working age adults. *Cochrane Database Syst Rev* 2000:CD001984.
- Karjalainen K, Malmivaara A, van Tulder M, Roine R, Jauhiainen M, Hurri H, et al. Biopsychosocial rehabilitation for upper limb repetitive strain injuries in working age adults. *Cochrane Database Syst Rev* 2000:CD002269.
- Karjalainen K, Malmivaara A, van Tulder M, Roine R, Jauhiainen M, Hurri H, et al. Multidisciplinary biopsychosocial rehabilitation for subacute low back pain among working age adults. *Cochrane Database Syst Rev* 2003:CD002193.
- Karjalainen K, Malmivaara A, van Tulder M, Roine R, Jauhiainen M, Hurri H, et al. Multidisciplinary biopsychosocial rehabilitation for neck and shoulder pain among working age adults. *Cochrane Database Syst Rev* 2003:CD002194.
- Keller A, Brox JI, Gunderson R, Holm I, Friis A, Reikeras O. Trunk muscle strength, cross-sectional area, and density in patients with chronic low back pain randomized to lumbar fusion or cognitive intervention and exercises. *Spine* 2004;29:3-8.
- Klüber Moffett JA, Carr J, Howarth E. High fear-avoiders of physical activity benefit from an exercise program for patients with back pain. *Spine* 2004; 29:1167-72; discussion 1173.
- Klüber Moffett JA, Jackson DA, Richmond S, Hahn S, Coulton S, Farrin A, et al. Randomised trial of a brief physiotherapy intervention compared with usual physiotherapy for neck pain patients: outcomes and patients' preference. *BMJ* 2005; 330:75.
- Kladny B, Fischer FC, Haase I. [Evaluation of specific stabilizing exercise in the treatment of low back pain and lumbar disk disease in outpatient rehabilitation]. *Z Orthop Ihre Grenzgeb* 2003;141:401-5.
- Knoop H, Stulemeijer M, Prins JB, van der Meer JW, Bleijenberg G. Is cognitive behaviour therapy for chronic fatigue syndrome also effective for pain symptoms? *Behav Res Ther* 2007;45:2034-43.
- Kofotolis N, Kellis E. Effects of two 4-week proprioceptive neuromuscular facilitation programs on muscle endurance, flexibility, and functional performance in women with chronic low back pain. *Phys Ther* 2006;86:1001-12.
- Kofotolis ND, Vlachopoulos SP, Kellis E. Sequentially allocated clinical trial of rhythmic stabilization exercises and TENS in women with chronic low back pain. *Clin Rehabil* 2008;22:99-111.
- Koldas Dogan S, Sonel Tur B, Kurtais Y, Atay MB. Comparison of three different approaches in the treatment of chronic low back pain. *Clin Rheumatol* 2008; 27:873-81.

- Kominski GF, Heslin KC, Morgenstern H, Hurwitz EL, Harber PI. Economic evaluation of four treatments for low-back pain: results from a randomized controlled trial. *Med Care* 2005;43:428-35.
- Kool J, Bachmann S, Oesch P, Knuesel O, Ambergen T, de Bie R, et al. Function-centered rehabilitation increases work days in patients with nonacute nonspecific low back pain: 1-year results from a randomized controlled trial. *Arch Phys Med Rehabil* 2007;88:1089-94.
- Kool JP, Oesch PR, Bachmann S, Knuesel O, Dierkes JG, Russo M, et al. Increasing days at work using function-centered rehabilitation in nonacute nonspecific low back pain: a randomized controlled trial. *Arch Phys Med Rehabil* 2005;86:857-64.
- Korthals-de Bos IB, Hoving JL, van Tulder MW, Rutten-van Molken MP, Ader HJ, de Vet HC, et al. Cost effectiveness of physiotherapy, manual therapy, and general practitioner care for neck pain: economic evaluation alongside a randomised controlled trial. *BMJ* 2003;326:911.
- Koumantakis GA, Watson PJ, Oldham JA. Supplementation of general endurance exercise with stabilisation training versus general exercise only. Physiological and functional outcomes of a randomised controlled trial of patients with recurrent low back pain. *Clin Biomech (Bristol, Avon)* 2005;20:474-82.
- Koumantakis GA, Watson PJ, Oldham JA. Trunk muscle stabilization training plus general exercise versus general exercise only: randomized controlled trial of patients with recurrent low back pain. *Phys Ther* 2005;85:209-25.
- Kuukkanen T, Malkia E, Kautiainen H, Pohjolainen T. Effectiveness of a home exercise programme in low back pain: a randomized five-year follow-up study. *Physiother Res Int* 2007;12:213-24.
- Kwekkeboom KL, Gretarsdottir E. Systematic review of relaxation interventions for pain. *J Nurs Scholarsh* 2006;38:269-77.
- Lamb SE, Lall R, Hansen Z, Withers EJ, Griffiths FE, Szczepura A, et al. Design considerations in a clinical trial of a cognitive behavioural intervention for the management of low back pain in primary care: Back Skills Training Trial. *BMC Musculoskelet Disord* 2007;8:14.
- Lansinger B, Larsson E, Persson LC, Carlsson JY. Qigong and exercise therapy in patients with long-term neck pain: a prospective randomized trial. *Spine* 2007;32:2415-22.
- Lemstra M, Olszynski WP. The effectiveness of multidisciplinary rehabilitation in the treatment of fibromyalgia: a randomized controlled trial. *Clin J Pain* 2005;21:166-74.
- Leonhardt C, Keller S, Chenot JF, Luckmann J, Basler HD, Wegscheider K, et al. TTM-based motivational counselling does not increase physical activity of low back pain patients in a primary care setting – A cluster-randomized controlled trial. *Patient Educ Couns* 2008;70:50-60.
- Lewis JS, Hewitt JS, Billington L, Cole S, Byng J, Karayiannis S. A randomized clinical trial comparing two physiotherapy interventions for chronic low back pain. *Spine* 2005;30:711-21.

- Lewis M, James M, Stokes E, Hill J, Sim J, Hay E, et al. An economic evaluation of three physiotherapy treatments for non-specific neck disorders alongside a randomized trial. *Rheumatology (Oxford)* 2007;46:1701-8.
- Linton SJ, Boersma K, Jansson M, Svard L, Botvalde M. The effects of cognitive-behavioral and physical therapy preventive interventions on pain-related sick leave: a randomized controlled trial. *Clin J Pain* 2005;21:109-19.
- Linton SJ, Nordin E. A 5-year follow-up evaluation of the health and economic consequences of an early cognitive behavioral intervention for back pain: a randomized, controlled trial. *Spine* 2006;31:853-8.
- Little P, Lewith G, Webley F, Evans M, Beattie A, Middleton K, et al. Randomised controlled trial of Alexander technique lessons, exercise, and massage (ATEAM) for chronic and recurrent back pain. *BMJ* 2008;337:a884.
- Machado LA, Azevedo DC, Capanema MB, Neto TN, Cerceau DM. Client-centered therapy vs exercise therapy for chronic low back pain: a pilot randomized controlled trial in Brazil. *Pain Med* 2007;8:251-8.
- Magnussen L, Strand LI, Skouen JS, Eriksen HR. Motivating disability pensioners with back pain to return to work – a randomized controlled trial. *J Rehabil Med* 2007;39:81-7.
- Manca A, Dumville JC, Torgerson DJ, Klaber Moffett JA, Mooney MP, Jackson DA, et al. Randomized trial of two physiotherapy interventions for primary care back and neck pain patients: cost effectiveness analysis. *Rheumatology (Oxford)* 2007;46:1495-501.
- Marshall P, Murphy B. Self-report measures best explain changes in disability compared with physical measures after exercise rehabilitation for chronic low back pain. *Spine* 2008;33:326-38.
- Matsutani LA, Marques AP, Ferreira EA, Assumpcao A, Lage LV, Casarotto RA, et al. Effectiveness of muscle stretching exercises with and without laser therapy at tender points for patients with fibromyalgia. *Clin Exp Rheumatol* 2007;25:410-5.
- Mattila R, Malmivaara A, Kastarinen M, Kivela SL, Nissinen A. Effects of lifestyle intervention on neck, shoulder, elbow and wrist symptoms. *Scand J Work Environ Health* 2004;30:191-8.
- Mattila R, Malmivaara A, Kastarinen M, Kivela SL, Nissinen A. The effects of lifestyle intervention for hypertension on low back pain: a randomized controlled trial. *Spine* 2007;32:2943-7.
- Maul I, Laubli T, Oliveri M, Krueger H. Long-term effects of supervised physical training in secondary prevention of low back pain. *Eur Spine J* 2005;14:599-611.
- Melancon B, Miller LH. Massage therapy versus traditional therapy for low back pain relief: implications for holistic nursing practice. *Holist Nurs Pract* 2005;19:116-21.
- Menzel NN, Lilley S, Robinson ME. Interventions to reduce back pain in rehabilitation hospital nursing staff. *Rehabil Nurs* 2006;31:138-47; discussion 148.

- Menzel NN, Robinson ME. Back pain in direct patient care providers: early intervention with cognitive behavioral therapy. *Pain Manag Nurs* 2006;7:53-63.
- Menzies V, Taylor AG, Bourguignon C. Effects of guided imagery on outcomes of pain, functional status, and self-efficacy in persons diagnosed with fibromyalgia. *J Altern Complement Med* 2006;12:23-30.
- Michaelson P, Sjolander P, Johansson H. Factors predicting pain reduction in chronic back and neck pain after multimodal treatment. *Clin J Pain* 2004;20:447-54.
- Milling LS, Shores JS, Coursen EL, Menario DJ, Farris CD. Response expectancies, treatment credibility, and hypnotic suggestibility: mediator and moderator effects in hypnotic and cognitive-behavioral pain interventions. *Ann Behav Med* 2007;33:167-78.
- Mirza SK, Deyo RA. Systematic review of randomized trials comparing lumbar fusion surgery to nonoperative care for treatment of chronic back pain. *Spine* 2007;32:816-23.
- Moffett JK, Jackson DA, Gardiner ED, Torgerson DJ, Coulton S, Eaton S, et al. Randomized trial of two physiotherapy interventions for primary care neck and back pain patients: 'McKenzie' vs brief physiotherapy pain management. *Rheumatology (Oxford)* 2006;45:1514-21.
- Mongini F, Ciccone G, Rota E, Ferrero L, Ugolini A, Evangelista A, et al. Effectiveness of an educational and physical programme in reducing headache, neck and shoulder pain: a workplace controlled trial. *Cephalalgia* 2008;28:541-52.
- Morone NE, Greco CM, Weiner DK. Mindfulness meditation for the treatment of chronic low back pain in older adults: a randomized controlled pilot study. *Pain* 2008;134:310-9.
- Moseley GL, Nicholas MK, Hodges PW. A randomized controlled trial of intensive neurophysiology education in chronic low back pain. *Clin J Pain* 2004;20:324-30.
- Munguia-Izquierdo D, Legaz-Arrese A. Exercise in warm water decreases pain and improves cognitive function in middle-aged women with fibromyalgia. *Clin Exp Rheumatol* 2007;25:823-30.
- Naylor MR, Keefe FJ, Brigidi B, Naud S, Helzer JE. Therapeutic Interactive Voice Response for chronic pain reduction and relapse prevention. *Pain* 2008;134:335-45.
- Neubauer E, Zahlten-Hinguranage A, Schiltenswolf M, Buchner M. [Multimodal therapy patients with chronic cervical and lumbar pain. Results of a comparative prospective study]. *Schmerz* 2006;20:210-8.
- Nickel MK, Lahmann C, Muehlbacher M, Nickel C, Pedrosa Gil F, Buschmann W, et al. Change in instrumental activities of daily living disability in female senior patients with musculoskeletal pain: a prospective, randomized, controlled trial. *Arch Gerontol Geriatr* 2006;42:247-55.
- Niemisto L, Sarna S, Lahtinen-Suopanki T, Lindgren KA, Hurri H. Predictive factors for 1-year outcome of chronic low back pain following manipulation, stabilizing exercises, and physician consultation or physician consultation alone. *J Rehabil Med* 2004;36:104-9.
- Nikander R, Malkia E, Parkkari J, Heinonen A, Starck H, Ylinen J. Dose-

- response relationship of specific training to reduce chronic neck pain and disability. *Med Sci Sports Exerc* 2006;38:2068-74.
- Nystuen P, Hagen KB. Solution-focused intervention for sick listed employees with psychological problems or muscle skeletal pain: a randomised controlled trial [ISRCTN39140363]. *BMC Public Health* 2006;6:69.
- Ostelo RW, de Vet HC, Berfelo MW, Kerckhoffs MR, Vlaeyen JW, Wolters PM, et al. Effectiveness of behavioral graded activity after first-time lumbar disc surgery: short term results of a randomized controlled trial. *Eur Spine J* 2003;12:637-44.
- Ostelo RW, Goossens ME, de Vet HC, van den Brandt PA. Economic evaluation of a behavioral-graded activity program compared to physical therapy for patients following lumbar disc surgery. *Spine* 2004;29:615-22.
- Paez-Blarrina M, Luciano C, Gutierrez-Martinez O, Valdivia S, Rodriguez-Valverde M, Ortega J. Coping with pain in the motivational context of values: comparison between an acceptance-based and a cognitive control-based protocol. *Behav Modif* 2008;32:403-22.
- Petersen T, Larsen K, Jacobsen S. One-year follow-up comparison of the effectiveness of McKenzie treatment and strengthening training for patients with chronic low back pain: outcome and prognostic factors. *Spine* 2007;32:2948-56.
- Plews-Ogan M, Owens JE, Goodman M, Wolfe P, Schorling J. A pilot study evaluating mindfulness-based stress reduction and massage for the management of chronic pain. *J Gen Intern Med* 2005;20:1136-8.
- Rakel B, Barr JO. Physical modalities in chronic pain management. *Nurs Clin North Am* 2003;38:477-94.
- Rasmussen-Barr E, Nilsson-Wikmar L, Arvidsson I. Stabilizing training compared with manual treatment in sub-acute and chronic low-back pain. *Man Ther* 2003;8:233-41.
- Rempel DM, Krause N, Goldberg R, Benner D, Hudes M, Goldner GU. A randomised controlled trial evaluating the effects of two workstation interventions on upper body pain and incident musculoskeletal disorders among computer operators. *Occup Environ Med* 2006;63:300-6.
- Riipinen M, Niemisto L, Lindgren KA, Hurri H. Psychosocial differences as predictors for recovery from chronic low back pain following manipulation, stabilizing exercises and physician consultation or physician consultation alone. *J Rehabil Med* 2005;37:152-8.
- Rivero-Arias O, Campbell H, Gray A, Fairbank J, Frost H, Wilson-MacDonald J. Surgical stabilisation of the spine compared with a programme of intensive rehabilitation for the management of patients with chronic low back pain: cost utility analysis based on a randomised controlled trial. *BMJ* 2005;330:1239.
- Rivero-Arias O, Gray A, Frost H, Lamb SE, Stewart-Brown S. Cost-utility analysis of physiotherapy treatment compared with physiotherapy advice in low back pain. *Spine* 2006;31:1381-7.
- Roche G, Ponthieux A, Parot-Shinkel E, Jousset N, Bontoux L, Dubus V, et al. Comparison of a functional restoration program with active individual physical therapy for patients with chronic low back

- pain: a randomized controlled trial. *Arch Phys Med Rehabil* 2007;88:1229-35.
- Rydeard R, Leger A, Smith D. Pilates-based therapeutic exercise: effect on subjects with nonspecific chronic low back pain and functional disability: a randomized controlled trial. *J Orthop Sports Phys Ther* 2006;36:472-84.
- Sarig-Bahat H. Evidence for exercise therapy in mechanical neck disorders. *Man Ther* 2003;8:10-20.
- Savolainen A, Ahlberg J, Nummila H, Nissinen M. Active or passive treatment for neck-shoulder pain in occupational health care? A randomized controlled trial. *Occup Med (Lond)* 2004;54:422-4.
- Schaffer SD, Yucha CB. Relaxation & pain management: the relaxation response can play a role in managing chronic and acute pain. *Am J Nurs* 2004;104:75-6, 78-9, 81-2.
- Schonstein E, Kenny DT, Keating J, Koes BW. Work conditioning, work hardening and functional restoration for workers with back and neck pain. *Cochrane Database Syst Rev* 2003:CD001822.
- Shaughnessy M, Caulfield B. A pilot study to investigate the effect of lumbar stabilisation exercise training on functional ability and quality of life in patients with chronic low back pain. *Int J Rehabil Res* 2004;27:297-301.
- Sherman KJ, Cherkin DC, Erro J, Miglioretti DL, Deyo RA. Comparing yoga, exercise, and a self-care book for chronic low back pain: a randomized, controlled trial. *Ann Intern Med* 2005;143:849-56.
- Sjogren T, Nissinen KJ, Jarvenpaa SK, Ojanen MT, Vanharanta H, Malkia EA. Effects of a workplace physical exercise intervention on the intensity of headache and neck and shoulder symptoms and upper extremity muscular strength of office workers: a cluster randomized controlled cross-over trial. *Pain* 2005;116:119-28.
- Skillgate E, Vingard E, Alfredsson L. Naprapathic manual therapy or evidence-based care for back and neck pain: a randomized, controlled trial. *Clin J Pain* 2007;23:431-9.
- Skouen JS, Grasdal A, Haldorsen EM. Return to work after comparing outpatient multidisciplinary treatment programs versus treatment in general practice for patients with chronic widespread pain. *Eur J Pain* 2006;10:145-52.
- Smeets RJ, Beelen S, Goossens ME, Schouten EG, Knottnerus JA, Vlaeyen JW. Treatment expectancy and credibility are associated with the outcome of both physical and cognitive-behavioral treatment in chronic low back pain. *Clin J Pain* 2008;24:305-15.
- Smeets RJ, Vlaeyen JW, Hidding A, Kester AD, van der Heijden GJ, van Geel AC, et al. Active rehabilitation for chronic low back pain: cognitive-behavioral, physical, or both? First direct post-treatment results from a randomized controlled trial [ISRCTN22714229]. *BMC Musculoskelet Disord* 2006;7:5.
- Soderlund A, Lindberg P. Cognitive behavioural components in physiotherapy management of chronic whiplash associated disorders (WAD) – a randomised group study. *G Ital Med Lav Ergon* 2007;29:A5-11.

- Spinhoven P, Ter Kuile M, Kole-Snijders AM, Hutten Mansfeld M, Den Ouden DJ, Vlaeyen JW. Catastrophizing and internal pain control as mediators of outcome in the multidisciplinary treatment of chronic low back pain. *Eur J Pain* 2004;8:211-9.
- Sran MM. To treat or not to treat: new evidence for the effectiveness of manual therapy. *Br J Sports Med* 2004;38:521-5.
- Staal JB, Hlobil H, Koke AJ, Twisk JW, Smid T, van Mechelen W. Graded activity for workers with low back pain: who benefits most and how does it work? *Arthritis Rheum* 2008;59:642-9.
- Steenstra IA, Anema JR, Bongers PM, de Vet HC, Knol DL, van Mechelen W. The effectiveness of graded activity for low back pain in occupational healthcare. *Occup Environ Med* 2006;63:718-25.
- Stener-Victorin E, Kruse-Smidje C, Jung K. Comparison between electro-acupuncture and hydrotherapy, both in combination with patient education and patient education alone, on the symptomatic treatment of osteoarthritis of the hip. *Clin J Pain* 2004;20:179-85.
- Stewart MJ, Maher CG, Refshauge KM, Herbert RD, Bogduk N, Nicholas M. Randomized controlled trial of exercise for chronic whiplash-associated disorders. *Pain* 2007;128:59-68.
- Suni J, Rinne M, Natri A, Statistisian MP, Parkkari J, Alaranta H. Control of the lumbar neutral zone decreases low back pain and improves self-evaluated work ability: a 12-month randomized controlled study. *Spine* 2006;31:E611-20.
- Tavafian SS, Jamshidi A, Mohammad K, Montazeri A. Low back pain education and short term quality of life: a randomized trial. *BMC Musculoskelet Disord* 2007;8:21.
- Tavafian SS, Jamshidi AR, Montazeri A. A randomized study of back school in women with chronic low back pain: quality of life at three, six, and twelve months follow-up. *Spine* 2008;33:1617-21.
- Teys P, Bisset L, Vicenzino B. The initial effects of a Mulligan's mobilization with movement technique on range of movement and pressure pain threshold in pain-limited shoulders. *Man Ther* 2008;13:37-42.
- Thieme K, Gromnica-Ihle E, Flor H. Operant behavioral treatment of fibromyalgia: a controlled study. *Arthritis Rheum* 2003;49:314-20.
- Thomas E, Croft PR, Paterson SM, Dziedzic K, Hay EM. What influences participants' treatment preference and can it influence outcome? Results from a primary care-based randomised trial for shoulder pain. *Br J Gen Pract* 2004;54:93-6.
- Tomas-Carus P, Hakkinen A, Gusi N, Leal A, Hakkinen K, Ortega-Alonso A. Aquatic training and detraining on fitness and quality of life in fibromyalgia. *Med Sci Sports Exerc* 2007;39:1044-50.
- Tornkvist L, Gardulf A, Strender LE. Effects of 'pain-advisers': district nurses' opinions regarding their own knowledge, management and documentation of patients in chronic pain. *Scand J Caring Sci* 2003;17:332-8.
- Underwood MR, Morton V, Farrin A. Do baseline characteristics predict response to treatment for low back pain?

- Secondary analysis of the UK BEAM dataset [ISRCTN32683578]. *Rheumatology (Oxford)* 2007;46:1297-302.
- Vad VB, Bhat AL, Tarabichi Y. The role of the Back Rx exercise program in diskogenic low back pain: a prospective randomized trial. *Arch Phys Med Rehabil* 2007;88:577-82.
- Walach H, Guthlin C, Konig M. Efficacy of massage therapy in chronic pain: a pragmatic randomized trial. *J Altern Complement Med* 2003;9:837-46.
- Valim V, Oliveira L, Suda A, Silva L, de Assis M, Barros Neto T, et al. Aerobic fitness effects in fibromyalgia. *J Rheumatol* 2003;30:1060-9.
- van den Hout JH, Vlaeyen JW, Heuts PH, Zijlema JH, Wijnen JA. Secondary prevention of work-related disability in nonspecific low back pain: does problem-solving therapy help? A randomized clinical trial. *Clin J Pain* 2003;19:87-96.
- van der Hulst M, Vollenbroek-Hutten MM, Groothuis-Oudshoorn KG, Hermens HJ. Multidisciplinary rehabilitation treatment of patients with chronic low back pain: a prognostic model for its outcome. *Clin J Pain* 2008;24:421-30.
- van der Roer N, van Tulder M, van Mechelen W, de Vet H. Economic evaluation of an intensive group training protocol compared with usual care physiotherapy in patients with chronic low back pain. *Spine* 2008;33:445-51.
- Veiersted KB, Gould KS, Osteras N, Hansson GA. Effect of an intervention addressing working technique on the biomechanical load of the neck and shoulders among hairdressers. *Appl Ergon* 2008;39:183-90.
- Veihelmann A, Devens C, Trouillier H, Birkenmaier C, Gerdesmeyer L, Refior HJ. Epidural neuroplasty versus physiotherapy to relieve pain in patients with sciatica: a prospective randomized blinded clinical trial. *J Orthop Sci* 2006;11:365-9.
- Weinstein JN, Tosteson TD, Lurie JD, Tosteson AN, Blood E, Hanscom B, et al. Surgical versus nonsurgical therapy for lumbar spinal stenosis. *N Engl J Med* 2008;358:794-810.
- Wessels T, Ewert T, Limm H, Rackwitz B, Stucki G. Change factors explaining reductions of "interference" in a multidisciplinary and an exercise prevention program for low back pain. *Clin J Pain* 2007;23:629-34.
- Viljanen M, Malmivaara A, Uitti J, Rinne M, Palmroos P, Laippala P. Effectiveness of dynamic muscle training, relaxation training, or ordinary activity for chronic neck pain: randomised controlled trial. *BMJ* 2003;327:475.
- Voerman GE, Sandsjo L, Vollenbroek-Hutten MM, Larsman P, Kadefors R, Hermens HJ. Effects of ambulant myo-feedback training and ergonomic counselling in female computer workers with work-related neck-shoulder complaints: a randomized controlled trial. *J Occup Rehabil* 2007;17:137-52.
- Voerman GE, Sandsjo L, Vollenbroek-Hutten MM, Larsman P, Kadefors R, Hermens HJ. Changes in cognitive-behavioral factors and muscle activation patterns after interventions for work-related neck-shoulder complaints: rela-

- tions with discomfort and disability. *J Occup Rehabil* 2007;17:593-609.
- Vollenbroek-Hutten MM, Hermens HJ, Wever D, Gorter M, Rinkel J, Ijzerman MJ. Differences in outcome of a multidisciplinary treatment between subgroups of chronic low back pain patients defined using two multiaxial assessment instruments: the multidimensional pain inventory and lumbar dynamometry. *Clin Rehabil* 2004;18:566-79.
- Von Korff M, Balderson BH, Saunders K, Miglioretti DL, Lin EH, Berry S, et al. A trial of an activating intervention for chronic back pain in primary care and physical therapy settings. *Pain* 2005; 113:323-30.
- Yeung CK, Leung MC, Chow DH. The use of electro-acupuncture in conjunction with exercise for the treatment of chronic low-back pain. *J Altern Complement Med* 2003;9:479-90.
- Ylinen J, Hakkinen A, Nykanen M, Kautiainen H, Takala EP. Neck muscle training in the treatment of chronic neck pain: a three-year follow-up study. *Eura Medicophys* 2007;43:161-9.
- Ylinen J, Kautiainen H, Wiren K, Hakkinen A. Stretching exercises vs manual therapy in treatment of chronic neck pain: a randomized, controlled cross-over trial. *J Rehabil Med* 2007; 39:126-32.
- Ylinen J, Takala EP, Nykanen M, Hakkinen A, Malkia E, Pohjolainen T, et al. Active neck muscle training in the treatment of chronic neck pain in women: a randomized controlled trial. *JAMA* 2003;289:2509-16.
- Ylinen JJ, Hakkinen AH, Takala EP, Nykanen MJ, Kautiainen HJ, Malkia EA, et al. Effects of neck muscle training in women with chronic neck pain: one-year follow-up study. *J Strength Cond Res* 2006;20:6-13.
- Ylinen JJ, Takala EP, Nykanen MJ, Kautiainen HJ, Hakkinen AH, Airaksinen OV. Effects of twelve-month strength training subsequent to twelve-month stretching exercise in treatment of chronic neck pain. *J Strength Cond Res* 2006;20:304-8.
- Zambito A, Bianchini D, Gatti D, Rossini M, Adami S, Viapiana O. Interferential and horizontal therapies in chronic low back pain due to multiple vertebral fractures: a randomized, double blind, clinical study. *Osteoporos Int* 2007;18:1541-5.
- Zaproudina N, Hanninen OO, Airaksinen O. Effectiveness of traditional bone setting in chronic neck pain: randomized clinical trial. *J Manipulative Physiol Ther* 2007; 30:432-7.

3.2 Fysisk aktivitet/träning, manuella och fysikaliska metoder samt beteendemedicinska behandlingar

Studier som granskats men inte uppfyllde inklusionskriterierna eller kravet på minst medelhög kvalitet och relevans

United Kingdom back pain exercise and manipulation (UK BEAM) randomised trial: effectiveness of physical treatments for back pain in primary care. *BMJ* 2004;329:1377.

Almeida TF, Roizenblatt S, Benedito-Silva AA, Tufik S. The effect of combined therapy (ultrasound and interferential current) on pain and sleep in fibromyalgia. *Pain* 2003;104:665-72.

Altan L, Bingol U, Aykac M, Koc Z, Yurtkuran M. Investigation of the effects of pool-based exercise on fibromyalgia syndrome. *Rheumatol Int* 2004;24:272-7.

Andersen LL, Kjaer M, Sogaard K, Hansen L, Kryger AI, Sjogaard G. Effect of two contrasting types of physical exercise on chronic neck muscle pain. *Arthritis Rheum* 2008;59:84-91.

Ansari NN, Ebadi S, Talebian S, Naghdi S, Mazaheri H, Olyaei G, et al. A randomized, single blind placebo controlled clinical trial on the effect of continuous ultrasound on low back pain. *Electromyogr Clin Neurophysiol* 2006;46:329-36.

Aure OF, Nilsen JH, Vasseljen O. Manual therapy and exercise therapy in patients with chronic low back pain: a randomized, controlled trial with 1-year follow-up. *Spine* 2003;28:525-31; discussion 531-2.

Bergman GJ, Winters JC, Groenier KH, Pool JJ, Meyboom-de Jong B, Postema K, et al. Manipulative therapy in addition to usual medical care for patients with shoulder dysfunction and pain: a randomized, controlled trial. *Ann Intern Med* 2004;141:432-9.

Bernaards CM, Ariens GA, Hildebrandt VH. The (cost-)effectiveness of a lifestyle physical activity intervention in addition to a work style intervention on the recovery from neck and upper limb symptoms in computer workers. *BMC Musculoskelet Disord* 2006;7:80.

Blangsted AK, Sogaard K, Hansen EA, Hannerz H, Sjogaard G. One-year randomized controlled trial with different physical-activity programs to reduce musculoskeletal symptoms in the neck and shoulders among office workers. *Scand J Work Environ Health* 2008;34:55-65.

Browder DA, Childs JD, Cleland JA, Fritz JM. Effectiveness of an extension-oriented treatment approach in a subgroup of subjects with low back pain: a randomized clinical trial. *Phys Ther* 2007;87:1608-18; discussion 1577-9.

Brox JI, Reikeras O, Nygaard O, Sorensen R, Indahl A, Holm I, et al. Lumbar instrumented fusion compared with cognitive intervention and exercises in patients with chronic back pain after previous surgery for disc herniation: a prospective randomized controlled study. *Pain* 2006;122:145-55.

Brox JI, Sorensen R, Friis A, Nygaard O, Indahl A, Keller A, et al. Randomized clinical trial of lumbar instrumented fusion and cognitive intervention and

- exercises in patients with chronic low back pain and disc degeneration. *Spine* 2003;28:1913-21.
- Buhrman M, Faltenhag S, Strom L, Andersson G. Controlled trial of Internet-based treatment with telephone support for chronic back pain. *Pain* 2004;111:368-77.
- Cambron JA, Gudavalli MR, Hedeker D, McGregor M, Jedlicka J, Keenum M, et al. One-year follow-up of a randomized clinical trial comparing flexion distraction with an exercise program for chronic low-back pain. *J Altern Complement Med* 2006;12:659-68.
- Carr JL, Klaber Moffett JA, Howarth E, Richmond SJ, Torgerson DJ, Jackson DA, et al. A randomized trial comparing a group exercise programme for back pain patients with individual physiotherapy in a severely deprived area. *Disabil Rehabil* 2005;27:929-37.
- Celestini M, Marchese A, Serenelli A, Graziani G. A randomized controlled trial on the efficacy of physical exercise in patients braced for instability of the lumbar spine. *Eura Medicophys* 2005;41:223-31.
- Chatzitheodorou D, Kabitsis C, Malliou P, Mougios V. A pilot study of the effects of high-intensity aerobic exercise versus passive interventions on pain, disability, psychological strain, and serum cortisol concentrations in people with chronic low back pain. *Phys Ther* 2007;87:304-12.
- Chatzitheodorou D, Mavromoustakos S, Milioti S. The effect of exercise on adrenocortical responsiveness of patients with chronic low back pain, controlled for psychological strain. *Clin Rehabil* 2008;22:319-28.
- Childs JD, Fritz JM, Flynn TW, Irrgang JJ, Johnson KK, Majkowski GR, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med* 2004;141:920-8.
- Christensen FB, Laurberg I, Bunge CE. Importance of the back-cafe concept to rehabilitation after lumbar spinal fusion: a randomized clinical study with a 2-year follow-up. *Spine* 2003;28:2561-9.
- Dalichau S, Stein B, Schafer K, Buhlmann JJ, Menken P. [Ultrasonic Quantification of spinal configuration and postural capacity for evaluation of different muscle strengthening programs in the therapy of back pain]. *Z Orthop Ihre Grenzgeb* 2005;143:79-85.
- Dehghani M, Sharpe L, Nicholas MK. Modification of attentional biases in chronic pain patients: a preliminary study. *Eur J Pain* 2004;8:585-94.
- Djavid GE, Mehrdad R, Ghasemi M, Hasan-Zadeh H, Sotoodeh-Manesh A, Pouryaghoob G. In chronic low back pain, low level laser therapy combined with exercise is more beneficial than exercise alone in the long term: a randomised trial. *Aust J Physiother* 2007;53:155-60.
- Donmez A, Karagulle MZ, Tercan N, Dinler M, Issever H, Karagulle M, et al. SPA therapy in fibromyalgia: a randomised controlled clinic study. *Rheumatol Int* 2005;26:168-72.

- Donta ST, Clauw DJ, Engel CC, Jr, Guarino P, Peduzzi P, Williams DA, et al. Cognitive behavioral therapy and aerobic exercise for Gulf War veterans' illnesses: a randomized controlled trial. *JAMA* 2003;289:1396-404.
- Donzelli S, Di Domenica E, Cova AM, Galletti R, Giunta N. Two different techniques in the rehabilitation treatment of low back pain: a randomized controlled trial. *Eura Medicophys* 2006;42:205-10.
- Ektor-Andersen J, Ingvarsson E, Kullendorff M, Orbaek P. High cost-benefit of early team-based biomedical and cognitive-behaviour intervention for long-term pain-related sickness absence. *J Rehabil Med* 2008;40:1-8.
- Erdogmus CB, Resch KL, Sabitzer R, Muller H, Nuhr M, Schoggl A, et al. Physiotherapy-based rehabilitation following disc herniation operation: results of a randomized clinical trial. *Spine* 2007;32:2041-9.
- Galantino ML, Bzdewka TM, Eissler-Russo JL, Holbrook ML, Mogck EP, Geigle P, et al. The impact of modified Hatha yoga on chronic low back pain: a pilot study. *Altern Ther Health Med* 2004;10:56-9.
- Geisser ME, Wiggert EA, Haig AJ, Colwell MO. A randomized, controlled trial of manual therapy and specific adjuvant exercise for chronic low back pain. *Clin J Pain* 2005;21:463-70.
- George SZ, Fritz JM, Childs JD. Investigation of elevated fear-avoidance beliefs for patients with low back pain: a secondary analysis involving patients enrolled in physical therapy clinical trials. *J Orthop Sports Phys Ther* 2008;38:50-8.
- Geraets JJ, Goossens ME, de Groot IJ, de Bruijn CP, de Bie RA, Dinant GJ, et al. Effectiveness of a graded exercise therapy program for patients with chronic shoulder complaints. *Aust J Physiother* 2005;51:87-94.
- Graham N, Gross AR, Goldsmith C. Mechanical traction for mechanical neck disorders: a systematic review. *J Rehabil Med* 2006;38:145-52.
- Gross AR, Hoving JL, Haines TA, Goldsmith CH, Kay T, Aker P, et al. A Cochrane review of manipulation and mobilization for mechanical neck disorders. *Spine* 2004;29:1541-8.
- Gur A, Karakoc M, Cevik R, Nas K, Sarac AJ. Efficacy of low power laser therapy and exercise on pain and functions in chronic low back pain. *Lasers Surg Med* 2003;32:233-8.
- Gusi N, Tomas-Carus P, Hakkinen A, Hakkinen K, Ortega-Alonso A. Exercise in waist-high warm water decreases pain and improves health-related quality of life and strength in the lower extremities in women with fibromyalgia. *Arthritis Rheum* 2006;55:66-73.
- Gustavsson C, von Koch L. Applied relaxation in the treatment of long-lasting neck pain: a randomized controlled pilot study. *J Rehabil Med* 2006;38:100-7.
- Haas M, Grouppe E, Kraemer DF. Dose-response for chiropractic care of chronic low back pain. *Spine J* 2004;4:574-83.
- Hakkinen A, Salo P, Tarvainen U, Wiren K, Ylinen J. Effect of manual therapy and stretching on neck muscle strength and mobility in chronic neck pain. *J Rehabil Med* 2007;39:575-9.

- Harts CC, Helmhout PH, de Bie RA, Staal JB. A high-intensity lumbar extensor strengthening program is little better than a low-intensity program or a waiting list control group for chronic low back pain: a randomised clinical trial. *Aust J Physiother* 2008;54:23-31.
- Hasson D, Arnetz B, Jelveus L, Edelstam B. A randomized clinical trial of the treatment effects of massage compared to relaxation tape recordings on diffuse long-term pain. *Psychother Psychosom* 2004;73:17-24.
- Hayden JA, van Tulder MW, Malmivaara AV, Koes BW. Meta-analysis: exercise therapy for nonspecific low back pain. *Ann Intern Med* 2005;142:765-75.
- Hayden JA, van Tulder MW, Tomlinson G. Systematic review: strategies for using exercise therapy to improve outcomes in chronic low back pain. *Ann Intern Med* 2005;142:776-85.
- Helewa A, Goldsmith CH, Smythe HA, Lee P, Obright K, Stitt L. Effect of therapeutic exercise and sleeping neck support on patients with chronic neck pain: a randomized clinical trial. *J Rheumatol* 2007;34:151-8.
- Helmhout PH, Harts CC, Staal JB, Candel MJ, de Bie RA. Comparison of a high-intensity and a low-intensity lumbar extensor training program as minimal intervention treatment in low back pain: a randomized trial. *Eur Spine J* 2004;13:537-47.
- Hemmila HM. Bone setting for prolonged neck pain: a randomized clinical trial. *J Manipulative Physiol Ther* 2005;28:508-15.
- Heymans MW, de Vet HC, Bongers PM, Knol DL, Koes BW, van Mechelen W. The effectiveness of high-intensity versus low-intensity back schools in an occupational setting: a pragmatic randomized controlled trial. *Spine* 2006;31:1075-82.
- Hoving JL, de Vet HC, Koes BW, Mameren H, Deville WL, van der Windt DA, et al. Manual therapy, physical therapy, or continued care by the general practitioner for patients with neck pain: long-term results from a pragmatic randomized clinical trial. *Clin J Pain* 2006;22:370-7.
- Hsieh LL, Kuo CH, Lee LH, Yen AM, Chien KL, Chen TH. Treatment of low back pain by acupuncture and physical therapy: randomised controlled trial. *BMJ* 2006;332:696-700.
- Hsieh LL, Kuo CH, Yen MF, Chen TH. A randomized controlled clinical trial for low back pain treated by acupuncture and physical therapy. *Prev Med* 2004;39:168-76.
- Hurwitz EL, Morgenstern H, Chiao C. Effects of recreational physical activity and back exercises on low back pain and psychological distress: findings from the UCLA Low Back Pain Study. *Am J Public Health* 2005;95:1817-24.
- Hurwitz EL, Morgenstern H, Kominski GF, Yu F, Chiang LM. A randomized trial of chiropractic and medical care for patients with low back pain: eighteen-month follow-up outcomes from the UCLA low back pain study. *Spine* 2006;31:611-21; discussion 622.
- Johnson RE, Jones GT, Wiles NJ, Chaddock C, Potter RG, Roberts C,

- et al. Active exercise, education, and cognitive behavioral therapy for persistent disabling low back pain: a randomized controlled trial. *Spine* 2007;32:1578-85.
- Jones TA. Rolfing. *Phys Med Rehabil Clin N Am* 2004;15:799-809, vi.
- Jousset N, Fanello S, Bontoux L, Dubus V, Billabert C, Vielle B, et al. Effects of functional restoration versus 3 hours per week physical therapy: a randomized controlled study. *Spine* 2004;29:487-93; discussion 494.
- Kaapa EH, Frantsi K, Sarna S, Malmivaara A. Multidisciplinary group rehabilitation versus individual physiotherapy for chronic nonspecific low back pain: a randomized trial. *Spine* 2006;31:371-6.
- Keller A, Brox JI, Gunderson R, Holm I, Friis A, Reikeras O. Trunk muscle strength, cross-sectional area, and density in patients with chronic low back pain randomized to lumbar fusion or cognitive intervention and exercises. *Spine* 2004;29:3-8.
- Klaber Moffett JA, Carr J, Howarth E. High fear-avoiders of physical activity benefit from an exercise program for patients with back pain. *Spine* 2004;29:1167-72; discussion 1173.
- Klaber Moffett JA, Jackson DA, Richmond S, Hahn S, Coulton S, Farrin A, et al. Randomised trial of a brief physiotherapy intervention compared with usual physiotherapy for neck pain patients: outcomes and patients' preference. *BMJ* 2005;330:75.
- Kladny B, Fischer FC, Haase I. [Evaluation of specific stabilizing exercise in the treatment of low back pain and lumbar disk disease in outpatient rehabilitation]. *Z Orthop Ihre Grenzgeb* 2003;141:401-5.
- Kofotolis N, Kellis E. Effects of two 4-week proprioceptive neuromuscular facilitation programs on muscle endurance, flexibility, and functional performance in women with chronic low back pain. *Phys Ther* 2006;86:1001-12.
- Kofotolis ND, Vlachopoulos SP, Kellis E. Sequentially allocated clinical trial of rhythmic stabilization exercises and TENS in women with chronic low back pain. *Clin Rehabil* 2008;22:99-111.
- Koldas Dogan S, Sonel Tur B, Kurtais Y, Atay MB. Comparison of three different approaches in the treatment of chronic low back pain. *Clin Rheumatol* 2008;27:873-81.
- Kominski GF, Heslin KC, Morgenstern H, Hurwitz EL, Harber PI. Economic evaluation of four treatments for low-back pain: results from a randomized controlled trial. *Med Care* 2005;43:428-35.
- Kool J, Bachmann S, Oesch P, Knuesel O, Ambergen T, de Bie R, et al. Function-centered rehabilitation increases work days in patients with nonacute nonspecific low back pain: 1-year results from a randomized controlled trial. *Arch Phys Med Rehabil* 2007;88:1089-94.
- Kool JP, Oesch PR, Bachmann S, Knuesel O, Dierkes JG, Russo M, et al. Increasing days at work using function-centered rehabilitation in nonacute nonspecific low back pain: a randomized controlled trial. *Arch Phys Med Rehabil* 2005;86:857-64.
- Korthals-de Bos IB, Hoving JL, van Tulder MW, Rutten-van Molken MP, Ader HJ, de Vet HC, et al. Cost effectiveness of physiotherapy, manual therapy, and general practitioner care for neck pain: economic evaluation alongside a random-

- mised controlled trial. *BMJ* 2003; 326:911.
- Koumantakis GA, Watson PJ, Oldham *JA. Trunk muscle stabilization training plus general exercise versus general exercise only: randomized controlled trial of patients with recurrent low back pain. *Phys Ther* 2005;85:209-25.
- Kuukkanen T, Malkia E, Kautiainen H, Pohjolainen T. Effectiveness of a home exercise programme in low back pain: a randomized five-year follow-up study. *Physiother Res Int* 2007;12:213-24.
- Kwekkeboom KL, Gretarsdottir E. Systematic review of relaxation interventions for pain. *J Nurs Scholarsh* 2006;38:269-77.
- Lamb SE, Lall R, Hansen Z, Withers EJ, Griffiths FE, Szczepura A, et al. Design considerations in a clinical trial of a cognitive behavioural intervention for the management of low back pain in primary care: Back Skills Training Trial. *BMC Musculoskelet Disord* 2007;8:14.
- Lansinger B, Larsson E, Persson LC, Carlsson JY. Qigong and exercise therapy in patients with long-term neck pain: a prospective randomized trial. *Spine* 2007;32:2415-22.
- Leonhardt C, Keller S, Chenot JF, Luckmann J, Basler HD, Wegscheider K, et al. TTM-based motivational counselling does not increase physical activity of low back pain patients in a primary care setting – A cluster-randomized controlled trial. *Patient Educ Couns* 2008;70:50-60.
- Lewis JS, Hewitt JS, Billington L, Cole S, Byng J, Karayiannis S. A randomized clinical trial comparing two physiotherapy interventions for chronic low back pain. *Spine* 2005;30:711-21.
- Linton SJ, Boersma K, Jansson M, Svard L, Botvalde M. The effects of cognitive-behavioral and physical therapy preventive interventions on pain-related sick leave: a randomized controlled trial. *Clin J Pain* 2005;21:109-19.
- Linton SJ, Nordin E. A 5-year follow-up evaluation of the health and economic consequences of an early cognitive behavioral intervention for back pain: a randomized, controlled trial. *Spine* 2006;31:853-8.
- Little P, Lewith G, Webley F, Evans M, Beattie A, Middleton K, et al. Randomised controlled trial of Alexander technique lessons, exercise, and massage (ATEAM) for chronic and recurrent back pain. *BMJ* 2008;337:a884.
- Machado LA, Azevedo DC, Capanema MB, Neto TN, Cerceau DM. Client-centered therapy vs exercise therapy for chronic low back pain: a pilot randomized controlled trial in Brazil. *Pain Med* 2007;8:251-8.
- Magnussen L, Rognsvag T, Tveito TH, Eriksen HR. Effect of a brief cognitive training programme in patients with long-lasting back pain evaluated as unfit for surgery. *J Health Psychol* 2005;10: 233-43.
- Manca A, Dumville JC, Torgerson DJ, Klaber Moffett JA, Mooney MP, Jackson DA, et al. Randomized trial of two physiotherapy interventions for primary care back and neck pain patients: cost effectiveness analysis. *Rheumatology (Oxford)* 2007; 46:1495-501.
- Marshall P, Murphy B. Self-report measures best explain changes in disability

compared with physical measures after exercise rehabilitation for chronic low back pain. *Spine* 2008;33:326-38.

Matsutani LA, Marques AP, Ferreira EA, Assumpcao A, Lage LV, Casarotto RA, et al. Effectiveness of muscle stretching exercises with and without laser therapy at tender points for patients with fibromyalgia. *Clin Exp Rheumatol* 2007;25:410-5.

Mattila R, Malmivaara A, Kastarinen M, Kivela SL, Nissinen A. Effects of lifestyle intervention on neck, shoulder, elbow and wrist symptoms. *Scand J Work Environ Health* 2004;30:191-8.

Maul I, Laubli T, Oliveri M, Krueger H. Long-term effects of supervised physical training in secondary prevention of low back pain. *Eur Spine J* 2005;14:599-611.

Melancon B, Miller LH. Massage therapy versus traditional therapy for low back pain relief: implications for holistic nursing practice. *Holist Nurs Pract* 2005;19:116-21.

Menzel NN, Lilley S, Robinson ME. Interventions to reduce back pain in rehabilitation hospital nursing staff. *Rehabil Nurs* 2006;31:138-47; discussion 148.

Menzel NN, Robinson ME. Back pain in direct patient care providers: early intervention with cognitive behavioral therapy. *Pain Manag Nurs* 2006;7:53-63.

Milling LS, Levine MR, Meunier SA. Hypnotic enhancement of cognitive-behavioral interventions for pain: an analogue treatment study. *Health Psychol* 2003;22:406-13.

Milling LS, Shores JS, Coursen EL, Menario DJ, Farris CD. Response expectancies, treatment credibility, and hypnotic suggestibility: mediator and moderator effects in hypnotic and cognitive-behavioral pain interventions. *Ann Behav Med* 2007;33:167-78.

Mirza SK, Deyo RA. Systematic review of randomized trials comparing lumbar fusion surgery to nonoperative care for treatment of chronic back pain. *Spine* 2007;32:816-23.

Moffett JK, Jackson DA, Gardiner ED, Torgerson DJ, Coulton S, Eaton S, et al. Randomized trial of two physiotherapy interventions for primary care neck and back pain patients: 'McKenzie' vs brief physiotherapy pain management. *Rheumatology (Oxford)* 2006;45:1514-21.

Mongini F, Ciccone G, Rota E, Ferrero L, Ugolini A, Evangelista A, et al. Effectiveness of an educational and physical programme in reducing headache, neck and shoulder pain: a workplace controlled trial. *Cephalalgia* 2008;28:541-52.

Moseley GL, Nicholas MK, Hodges PW. A randomized controlled trial of intensive neurophysiology education in chronic low back pain. *Clin J Pain* 2004;20:324-30.

Munguia-Izquierdo D, Legaz-Arrese A. Exercise in warm water decreases pain and improves cognitive function in middle-aged women with fibromyalgia. *Clin Exp Rheumatol* 2007;25:823-30.

Naylor MR, Keefe FJ, Brigidi B, Naud S, Helzer JE. Therapeutic Interactive Voice Response for chronic pain reduction and relapse prevention. *Pain* 2008;134:335-45.

- Nikander R, Malkia E, Parkkari J, Heinonen A, Starck H, Ylinen J. Dose-response relationship of specific training to reduce chronic neck pain and disability. *Med Sci Sports Exerc* 2006;38:2068-74.
- Ostelo RW, Goossens ME, de Vet HC, van den Brandt PA. Economic evaluation of a behavioral-graded activity program compared to physical therapy for patients following lumbar disc surgery. *Spine* 2004;29:615-22.
- Paez-Blarrina M, Luciano C, Gutierrez-Martinez O, Valdivia S, Rodriguez-Valverde M, Ortega J. Coping with pain in the motivational context of values: comparison between an acceptance-based and a cognitive control – based protocol. *Behav Modif* 2008;32:403-22.
- Plews-Ogan M, Owens JE, Goodman M, Wolfe P, Schorling J. A pilot study evaluating mindfulness-based stress reduction and massage for the management of chronic pain. *J Gen Intern Med* 2005;20:1136-8.
- Rakel B, Barr JO. Physical modalities in chronic pain management. *Nurs Clin North Am* 2003;38:477-94.
- Rasmussen-Barr E, Nilsson-Wikmar L, Arvidsson I. Stabilizing training compared with manual treatment in sub-acute and chronic low-back pain. *Man Ther* 2003;8:233-41.
- Rydeard R, Leger A, Smith D. Pilates-based therapeutic exercise: effect on subjects with nonspecific chronic low back pain and functional disability: a randomized controlled trial. *J Orthop Sports Phys Ther* 2006;36:472-84.
- Sarig-Bahat H. Evidence for exercise therapy in mechanical neck disorders. *Man Ther* 2003;8:10-20.
- Savolainen A, Ahlberg J, Nummila H, Nissinen M. Active or passive treatment for neck-shoulder pain in occupational health care? A randomized controlled trial. *Occup Med (Lond)* 2004;54:422-4.
- Schaffer SD, Yucha CB. Relaxation & pain management: the relaxation response can play a role in managing chronic and acute pain. *Am J Nurs* 2004;104:75-6, 78-9, 81-2.
- Schonstein E, Kenny DT, Keating J, Koes BW. Work conditioning, work hardening and functional restoration for workers with back and neck pain. *Cochrane Database Syst Rev* 2003: CD001822.
- Shaughnessy M, Caulfield B. A pilot study to investigate the effect of lumbar stabilisation exercise training on functional ability and quality of life in patients with chronic low back pain. *Int J Rehabil Res* 2004;27:297-301.
- Sjogren T, Nissinen KJ, Jarvenpaa SK, Ojanen MT, Vanharanta H, Malkia EA. Effects of a workplace physical exercise intervention on the intensity of headache and neck and shoulder symptoms and upper extremity muscular strength of office workers: a cluster randomized controlled cross-over trial. *Pain* 2005;116:119-28.
- Skillgate E, Vingard E, Alfredsson L. Naprapathic manual therapy or evidence-based care for back and neck pain: a randomized, controlled trial. *Clin J Pain* 2007;23:431-9.

- Smeets RJ, Beelen S, Goossens ME, Schouten EG, Knottnerus JA, Vlaeyen JW. Treatment expectancy and credibility are associated with the outcome of both physical and cognitive-behavioral treatment in chronic low back pain. *Clin J Pain* 2008;24:305-15.
- Soderlund A, Lindberg P. Cognitive behavioural components in physiotherapy management of chronic whiplash associated disorders (WAD) – a randomised group study. *G Ital Med Lav Ergon* 2007;29:A5-11.
- Sran MM. To treat or not to treat: new evidence for the effectiveness of manual therapy. *Br J Sports Med* 2004;38:521-5.
- Staal JB, Hlobil H, Koke AJ, Twisk JW, Smid T, van Mechelen W. Graded activity for workers with low back pain: who benefits most and how does it work? *Arthritis Rheum* 2008;59:642-9.
- Steenstra IA, Anema JR, Bongers PM, de Vet HC, Knol DL, van Mechelen W. The effectiveness of graded activity for low back pain in occupational healthcare. *Occup Environ Med* 2006;63:718-25.
- Suni J, Rinne M, Natri A, Statistisian MP, Parkkari J, Alaranta H. Control of the lumbar neutral zone decreases low back pain and improves self-evaluated work ability: a 12-month randomized controlled study. *Spine* 2006;31:E611-20.
- Tavafian SS, Jamshidi A, Mohammad K, Montazeri A. Low back pain education and short term quality of life: a randomized trial. *BMC Musculoskelet Disord* 2007;8:21.
- Thieme K, Flor H, Turk DC. Psychological pain treatment in fibromyalgia syndrome: efficacy of operant behavioural and cognitive behavioural treatments. *Arthritis Res Ther* 2006;8:R121.
- Thieme K, Gromnica-Ihle E, Flor H. Operant behavioral treatment of fibromyalgia: a controlled study. *Arthritis Rheum* 2003;49:314-20.
- Thieme K, Turk DC, Flor H. Responder criteria for operant and cognitive-behavioral treatment of fibromyalgia syndrome. *Arthritis Rheum* 2007;57:830-6.
- Tomas-Carus P, Hakkinen A, Gusi N, Leal A, Hakkinen K, Ortega-Alonso A. Aquatic training and detraining on fitness and quality of life in fibromyalgia. *Med Sci Sports Exerc* 2007;39:1044-50.
- Tornkvist L, Gardulf A, Strender LE. Effects of 'pain-advisers': district nurses' opinions regarding their own knowledge, management and documentation of patients in chronic pain. *Scand J Caring Sci* 2003;17:332-8.
- Turner-Stokes L, Erkeller-Yuksel F, Miles A, Pincus T, Shipley M, Pearce S. Out-patient cognitive behavioral pain management programs: a randomized comparison of a group-based multidisciplinary versus an individual therapy model. *Arch Phys Med Rehabil* 2003;84:781-8.
- Vad VB, Bhat AL, Tarabichi Y. The role of the Back Rx exercise program in diskogenic low back pain: a prospective randomized trial. *Arch Phys Med Rehabil* 2007;88:577-82.
- Walach H, Guthlin C, Konig M. Efficacy of massage therapy in chronic pain: a pragmatic randomized trial. *J Altern Complement Med* 2003;9:837-46.

- Valim V, Oliveira L, Suda A, Silva L, de Assis M, Barros Neto T, et al. Aerobic fitness effects in fibromyalgia. *J Rheumatol* 2003;30:1060-9.
- van den Hout JH, Vlaeyen JW, Heuts PH, Zijlema JH, Wijnen JA. Secondary prevention of work-related disability in nonspecific low back pain: does problem-solving therapy help? A randomized clinical trial. *Clin J Pain* 2003;19:87-96.
- van der Roer N, van Tulder M, van Mechelen W, de Vet H. Economic evaluation of an intensive group training protocol compared with usual care physiotherapy in patients with chronic low back pain. *Spine* 2008;33:445-51.
- Vas J, Ortega C, Olmo V, Perez-Fernandez F, Hernandez L, Medina I, et al. Single-point acupuncture and physiotherapy for the treatment of painful shoulder: a multicentre randomized controlled trial. *Rheumatology (Oxford)* 2008;47:887-93.
- Veihelmann A, Devens C, Trouillier H, Birkenmaier C, Gerdesmeyer L, Refior HJ. Epidural neuroplasty versus physiotherapy to relieve pain in patients with sciatica: a prospective randomized blinded clinical trial. *J Orthop Sci* 2006;11:365-9.
- Weinstein JN, Tosteson TD, Lurie JD, Tosteson AN, Blood E, Hanscom B, et al. Surgical versus nonsurgical therapy for lumbar spinal stenosis. *N Engl J Med* 2008;358:794-810.
- Wessels T, Ewert T, Limm H, Rackwitz B, Stucki G. Change factors explaining reductions of "interference" in a multidisciplinary and an exercise prevention program for low back pain. *Clin J Pain* 2007;23:629-34.
- Viljanen M, Malmivaara A, Uitti J, Rinne M, Palmroos P, Laippala P. Effectiveness of dynamic muscle training, relaxation training, or ordinary activity for chronic neck pain: randomised controlled trial. *BMJ* 2003;327:475.
- Voerman GE, Sandsjo L, Vollenbroek-Hutten MM, Larsman P, Kadefors R, Hermens HJ. Effects of ambulant myofeed-back training and ergonomic counselling in female computer workers with work-related neck-shoulder complaints: a randomized controlled trial. *J Occup Rehabil* 2007;17:137-52.
- Ylinen J, Hakkinen A, Nykanen M, Kautiainen H, Takala EP. Neck muscle training in the treatment of chronic neck pain: a three-year follow-up study. *Eura Medicophys* 2007;43:161-9.
- Ylinen JJ, Takala EP, Nykanen MJ, Kautiainen HJ, Hakkinen AH, Airaksinen OV. Effects of twelve-month strength training subsequent to twelve-month stretching exercise in treatment of chronic neck pain. *J Strength Cond Res* 2006;20:304-8.
- Zambito A, Bianchini D, Gatti D, Rossini M, Adami S, Viapiana O. Interferential and horizontal therapies in chronic low back pain due to multiple vertebral fractures: a randomized, double blind, clinical study. *Osteoporos Int* 2007;18:1541-5.
- Zaproudina N, Hanninen OO, Airaksinen O. Effectiveness of traditional bone setting in chronic neck pain: randomized clinical trial. *J Manipulative Physiol Ther* 2007;30:432-7.

3.3 Psykologiska behandlingsmetoder givna av en behandlare

Studier som granskats men inte uppfyllde inklusionskriterierna

Alvarez-Nemegyei J, Negreros-Castillo A, Nuño-Gutiérrez BL, Alvarez-Berzunza J, Alcocer-Martínez LM. [Ericksonian hypnosis in women with fibromyalgia syndrome]. In: Revista médica del Instituto Mexicano del Seguro Social; 2007. p 395-401.

Asenlof P, Denison E, Lindberg P. Idiographic outcome analyses of the clinical significance of two interventions for patients with musculoskeletal pain. *Behav Res Ther* 2006;44:947-65.

Astin JA. Mind-body therapies for the management of pain. *Clin J Pain* 2004; 20:27-32.

Astin JA, Berman BM, Bausell B, Lee WL, Hochberg M, Forsys KL. The efficacy of mindfulness meditation plus Qigong movement therapy in the treatment of fibromyalgia: a randomized controlled trial. *J Rheumatol* 2003;30:2257-62.

Astin JA, Shapiro SL, Eisenberg DM, Forsys KL. Mind-body medicine: state of the science, implications for practice. *J Am Board Fam Pract* 2003;16:131-47.

Babu AS, Mathew E, Danda D, Prakash H. Management of patients with fibromyalgia using biofeedback: a randomized control trial. *Indian J Med Sci* 2007;61:455-61.

Becker A, Leonhardt C, Kochen MM, Keller S, Wegscheider K, Baum E, et al. Effects of two guideline implementation strategies on patient outcomes in primary

care: a cluster randomized controlled trial. In: *Spine*; 2008. p 473-80.

Bernaards CM, Ariens GA, Knol DL, Hildebrandt VH. The effectiveness of a work style intervention and a lifestyle physical activity intervention on the recovery from neck and upper limb symptoms in computer workers. *Pain* 2007;132:142-53.

Bieber C, Muller KG, Blumenstiel K, Schneider A, Richter A, Wilke S, et al. Long-term effects of a shared decision-making intervention on physician-patient interaction and outcome in fibromyalgia. A qualitative and quantitative 1 year follow-up of a randomized controlled trial. *Patient Educ Couns* 2006;63: 357-66.

Brattberg G. Internet-based rehabilitation for individuals with chronic pain and burnout II: a long-term follow-up. *Int J Rehabil Res* 2007;30:231-4.

Buhrman M, Faltenhag S, Strom L, Andersson G. Controlled trial of Internet-based treatment with telephone support for chronic back pain. *Pain* 2004;111:368-77.

Carson JW, Keefe FJ, Lynch TR, Carson KM, Goli V, Fras AM, et al. Loving-kindness meditation for chronic low back pain: results from a pilot trial. *J Holist Nurs* 2005;23:287-304.

Castel A, Perez M, Sala J, Padrol A, Rull M. Effect of hypnotic suggestion on fibromyalgic pain: comparison between hypnosis and relaxation. *Eur J Pain* 2007;11: 463-8.

Chou R, Huffman LH. Nonpharmacologic therapies for acute and chronic low

- back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Ann Intern Med* 2007;147:492-504.
- Coudeyre E, Givron P, Vanbiervliet W, Benaïm C, Hérisson C, Pelissier J, et al. [The role of an information booklet or oral information about back pain in reducing disability and fear-avoidance beliefs among patients with subacute and chronic low back pain. A randomized controlled trial in a rehabilitation unit]. In: *Annales de réadaptation et de médecine physique: revue scientifique de la Société française de rééducation fonctionnelle de réadaptation et de médecine physique*; 2006. p 600-8.
- Dahl J, Wilson KG, Nilsson A. Acceptance and commitment therapy and the treatment of persons at risk for long-term disability resulting from stress and pain symptoms: A preliminary randomized trial. *Behav Ther* 2004;35:785-801.
- Dobscha SK, Corson K, Leibowitz RQ, Sullivan MD, Gerrity MS. Rationale, design, and baseline findings from a randomized trial of collaborative care for chronic musculoskeletal pain in primary care. In: *Pain medicine (Malden, Mass.)*; 2008. p 1050-64.
- Dobscha SK, Corson K, Perrin NA, Hanson GC, Leibowitz RQ, Doak MN, et al. Collaborative care for chronic pain in primary care: a cluster randomized trial. *JAMA* 2009;301:1242-52.
- Douglas W, Graham C, Anderson D, Rogerson K. Managing chronic pain through cognitive change and multidisciplinary treatment program. *Australian Psychologist* 2004;39:201-207.
- Edinger JD, Wohlgenuth WK, Krystal AD, Rice JR. Behavioral insomnia therapy for fibromyalgia patients: a randomized clinical trial. *Arch Intern Med* 2005;165:2527-35.
- Elkins G, Jensen MP, Patterson DR. Hypnototherapy for the management of chronic pain. *International Journal of Clinical and Experimental Hypnosis* 2007;55:275-287.
- Elkins G, Jensen MP, Patterson DR. Hypnototherapy for the management of chronic pain. *Int J Clin Exp Hypn* 2007;55:275-87.
- Engers A, Jellema P, Wensing M, van der Windt DA, Grol R, van Tulder MW. Individual patient education for low back pain. *Cochrane Database Syst Rev* 2008: CD004057.
- Field T, Hernandez-Reif M, Diego M, Fraser M. Lower back pain and sleep disturbance are reduced following massage therapy. In: *Journal of Bodywork and Movement Therapies*; 2007. p 141-5.
- Fors EA, Sexton H, Gotestam KG. The effect of guided imagery and amitriptyline on daily fibromyalgia pain: a prospective, randomized, controlled trial. *J Psychiatr Res* 2002;36:179-87.
- Galantino ML, Bzdewka TM, Eissler-Russo JL, Holbrook ML, Mogck EP, Geigle P, et al. The impact of modified Hatha yoga on chronic low back pain: a pilot study. *Altern Ther Health Med* 2004;10:56-9.
- Gardner-Nix J, Backman S, Barbati J, Grummitt J. Evaluating distance education of a mindfulness-based meditation

- programme for chronic pain management. *J Telemed Telecare* 2008;14:88-92.
- Gerard S, Smith BH, Simpson JA. A randomized controlled trial of spiritual healing in restricted neck movement. *J Altern Complement Med* 2003;9:467-77.
- Grondahl JR, Rosvold EO. Hypnosis as a treatment of chronic widespread pain in general practice: a randomized controlled pilot trial. *BMC Musculoskelet Disord* 2008;9:124.
- Gross AR, Goldsmith C, Hoving JL, Haines T, Peloso P, Aker P, et al. Conservative management of mechanical neck disorders: a systematic review. *J Rheumatol* 2007;34:1083-102.
- Gustavsson C, von Koch L. Applied relaxation in the treatment of long-lasting neck pain: a randomized controlled pilot study. *J Rehabil Med* 2006;38:100-7.
- Hadhazy VA, Ezzo J, Creamer P, Berman BM. Mind-body therapies for the treatment of fibromyalgia. A systematic review. *J Rheumatol* 2000;27:2911-8.
- Haines T, Gross A, Goldsmith CH, Perry L. Patient education for neck pain with or without radiculopathy. *Cochrane Database Syst Rev* 2008:CD005106.
- Hammond A, Freeman K. Community patient education and exercise for people with fibromyalgia: a parallel group randomized controlled trial. *Clin Rehabil* 2006;20:835-46.
- Hasson D, Arnetz B, Jelveus L, Edelstam B. A randomized clinical trial of the treatment effects of massage compared to relaxation tape recordings on diffuse long-term pain. *Psychother Psychosom* 2004;73:17-24.
- Haugli L, Steen E, Laerum E, Nygard R, Finset A. Learning to have less pain – is it possible? A one-year follow-up study of the effects of a personal construct group learning programme on patients with chronic musculoskeletal pain. *Patient Educ Couns* 2001;45:111-8.
- Haythornthwaite JA. Clinical trials studying pharmacotherapy and psychological treatments alone and together. *Neurology* 2005;65:S20-31.
- Henningsen P, Zipfel S, Herzog W. Management of functional somatic syndromes. *Lancet* 2007;369:946-55.
- Hochlehnert A, Richter A, Bludau HB, Bieber C, Blumenstiel K, Mueller K, et al. A computer-based information-tool for chronic pain patients. Computerized information to support the process of shared decision-making. *Patient Educ Couns* 2006;61:92-8.
- Hoffman BM, Papas RK, Chatkoff DK, Kerns RD. Meta-analysis of psychological interventions for chronic low back pain. *Health Psychol* 2007;26:1-9.
- Hurwitz EL, Carragee EJ, van der Velde G, Carroll LJ, Nordin M, Guzman J, et al. Treatment of neck pain: noninvasive interventions: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine* 2008;33:S123-52.
- Jellema P, van der Roer N, van der Windt DA, van Tulder MW, van der Horst HE, Stalman WA, et al. Low back pain in general practice: cost-effectiveness of a

- minimal psychosocial intervention versus usual care. *Eur Spine J* 2007;16:1812-21.
- Jensen M, Patterson DR. Hypnotic treatment of chronic pain. *J Behav Med* 2006; 29:95-124.
- Johnson RE, Jones GT, Wiles NJ, Chad-dock C, Potter RG, Roberts C, et al. Active exercise, education, and cognitive behavioral therapy for persistent disabling low back pain: a randomized controlled trial. *Spine* 2007;32:1578-85.
- Kainz B, Gülich M, Engel EM, Jäckel WH. [Comparison of three outpatient therapy forms for treatment of chronic low back pain – findings of a multi-centre, cluster randomized study]. In: *Die Rehabilitation*; 2006. p 65-77.
- Kerns JW, White A, Nashelsky J, Sherman S. Does psychiatric treatment help patients with intractable chronic pain? *J Fam Pract* 2006;55:235-6.
- King SJ, Wessel J, Bhambhani Y, Sholter D, Maksymowych W. The effects of exercise and education, individually or combined, in women with fibromyalgia. *J Rheumatol* 2002;29:2620-7.
- Kitahara M, Kojima KK, Ohmura A. Efficacy of interdisciplinary treatment for chronic nonmalignant pain patients in Japan. In: *The Clinical journal of pain*; 2006. p 647-55.
- Kjellgren A, Sundequist U, Norlander T, Archer T. Effects of flotation-REST on muscle tension pain. *Pain Res Manag* 2001;6:181-9.
- Kluber Moffett JA, Carr J, Howarth E. High fear-avoiders of physical activity benefit from an exercise program for patients with back pain. *Spine* 2004; 29:1167-72; discussion 1173.
- Kole-Snijders AM, Vlaeyen JW, Goossens ME, Rutten-van Molken MP, Heuts PH, van Breukelen G, et al. Chronic low-back pain: what does cognitive coping skills training add to operant behavioral treatment? Results of a randomized clinical trial. *J Consult Clin Psychol* 1999;67: 931-44.
- Kullich W, Bernatzky G, Hesse HP, Wendtner F, Likar R, Klein G. Music therapy - impact on pain, sleep, and quality of life in low back pain. In: *Wiener Medizinische Wochenschrift*; 2003. p 217-221.
- Kurtais Y, Kutlay S, Ergin S. Exercise and cognitive-behavioural treatment in fibromyalgia syndrome. *Curr Pharm Des* 2006;12:37-45.
- Lewandowski WA. Patterning of pain and power with guided imagery. *Nurs Sci Q* 2004;17:233-41.
- Liddle SD, Gracey JH, Baxter GD. Advice for the management of low back pain: a systematic review of randomised controlled trials. *Man Ther* 2007;12:310-27.
- Linton SJ, Boersma K, Jansson M, Overmeer T, Lindblom K, Vlaeyen JW. A randomized controlled trial of exposure in vivo for patients with spinal pain reporting fear of work-related activities. *Eur J Pain* 2008;12:722-30.
- Linton SJ, Nordin E. A 5-year follow-up evaluation of the health and economic consequences of an early cognitive behavioral intervention for back pain: a ran-

- domized, controlled trial. *Spine* 2006; 31:853-8.
- Linton SJ, Ryberg M. A cognitive-behavioral group intervention as prevention for persistent neck and back pain in a non-patient population: a randomized controlled trial. *Pain* 2001;90:83-90.
- Little P, Lewith G, Webley F, Evans M, Beattie A, Middleton K, et al. Randomised controlled trial of Alexander technique lessons, exercise, and massage (ATEAM) for chronic and recurrent back pain. *BMJ* 2008;337:a884.
- Lohnberg JA. A review of outcome studies on cognitive-behavioral therapy for reducing fear-avoidance beliefs among individuals with chronic pain. *Journal of Clinical Psychology in Medical Settings* 2007;14:113-122.
- Mannerkorpi K, Arndorw M. Efficacy and feasibility of a combination of body awareness therapy and qigong in patients with fibromyalgia: a pilot study. *J Rehabil Med* 2004;36:279-81.
- McCracken LM, Turk DC. Behavioral and cognitive-behavioral treatment for chronic pain: outcome, predictors of outcome, and treatment process. *Spine* 2002;27:2564-73.
- Mehling WE, Hamel KA, Acree M, Byl N, Hecht FM. Randomized, controlled trial of breath therapy for patients with chronic low-back pain. *Altern Ther Health Med* 2005;11:44-52.
- Molton IR, Graham C, Stoelb BL, Jensen MP. Current psychological approaches to the management of chronic pain. *Curr Opin Anaesthesiol* 2007;20:485-9.
- Moore JE, Von Korff M, Cherkin D, Saunders K, Lorig K. A randomized trial of a cognitive-behavioral program for enhancing back pain self care in a primary care setting. *Pain* 2000;88: 145-53.
- Morley S, Eccleston C, Williams A. Systematic review and meta-analysis of randomized controlled trials of cognitive behaviour therapy and behaviour therapy for chronic pain in adults, excluding headache. *Pain* 1999;80:1-13.
- Moseley GL. Joining forces – combining cognition-targeted motor control training with group or individual pain physiology education: a successful treatment for chronic low back pain. In: *Journal of Manual & Manipulative Therapy*; 2003. p 88-94.
- Moseley GL, Nicholas MK, Hodges PW. A randomized controlled trial of intensive neurophysiology education in chronic low back pain. *Clin J Pain* 2004;20:324-30.
- Naylor MR, Keefe FJ, Brigidi B, Naud S, Helzer JE. Therapeutic Interactive Voice Response for chronic pain reduction and relapse prevention. *Pain* 2008;134:335-45.
- Nielson WR, Weir R. Biopsychosocial approaches to the treatment of chronic pain. *Clin J Pain* 2001;17:S114-27.
- Nordin M, Balague F, Cedraschi C. Non-specific lower-back pain: surgical versus nonsurgical treatment. *Clin Orthop Relat Res* 2006;443:156-67.
- Norman SA, Lumley MA, Dooley JA, Diamond MP. For whom does it work? Moderators of the effects of written

- emotional disclosure in a randomized trial among women with chronic pelvic pain. In: *Psychosomatic medicine*; 2004. p 174-83.
- Ostelo RW, van Tulder MW, Vlaeyen JW, Linton SJ, Morley SJ, Assendelft WJ. Behavioural treatment for chronic low-back pain. *Cochrane Database Syst Rev* 2005;CD002014.
- Patterson DR, Jensen MP. Hypnosis and clinical pain. *Psychol Bull* 2003; 129:495-521.
- Plews-Ogan M, Owens JE, Goodman M, Wolfe P, Schorling J. A pilot study evaluating mindfulness-based stress reduction and massage for the management of chronic pain. *J Gen Intern Med* 2005;20:1136-8.
- Raine R, Haines A, Sensky T, Hutchings A, Larkin K, Black N. Systematic review of mental health interventions for patients with common somatic symptoms: can research evidence from secondary care be extrapolated to primary care? *BMJ* 2002;325:1082.
- Rooks DS, Gautam S, Romeling M, Cross ML, Stratigakis D, Evans B, et al. Group exercise, education, and combination self-management in women with fibromyalgia: a randomized trial. *Arch Intern Med* 2007; 167:2192-200.
- Rossy LA, Buckelew SP, Dorr N, Hagglund KJ, Thayer JF, McIntosh MJ, et al. A meta-analysis of fibromyalgia treatment interventions. *Ann Behav Med* 1999;21:180-91.
- Schofield P. The effects of Snoezelen on chronic pain. *Nurs Stand* 2000;15:33-4.
- Schofield P. Evaluating Snoezelen for relaxation within chronic pain management. *Br J Nurs* 2002;11:812-21.
- Schweikert B, Jacobi E, Seitz R, Cziske R, Ehlert A, Knab J, et al. Effectiveness and cost-effectiveness of adding a cognitive behavioral treatment to the rehabilitation of chronic low back pain. *J Rheumatol* 2006;33:2519-26.
- Sephton SE, Salmon P, Weissbecker I, Ulmer C, Floyd A, Hoover K, et al. Mindfulness meditation alleviates depressive symptoms in women with fibromyalgia: results of a randomized clinical trial. *Arthritis Rheum* 2007;57:77-85.
- Sherman KJ, Cherkin DC, Erro J, Miglioretti DL, Deyo RA. Comparing yoga, exercise, and a self-care book for chronic low back pain: a randomized, controlled trial. *Ann Intern Med* 2005; 143:849-56.
- Siedliecki SL, Good M. Effect of music on power, pain, depression and disability. *J Adv Nurs* 2006;54:553-62.
- Sim J, Adams N. Physical and other non-pharmacological interventions for fibromyalgia. *Baillieres Best Pract Res Clin Rheumatol* 1999;13:507-23.
- Sim J, Adams N. Systematic review of randomized controlled trials of non-pharmacological interventions for fibromyalgia. *Clin J Pain* 2002;18:324-36.
- Smeets RJ, Beelen S, Goossens ME, Schouten EG, Knottnerus JA, Vlaeyen JW. Treatment expectancy and credibility are associated with the outcome of both physical and cognitive-behavioral treat-

- ment in chronic low back pain. *Clin J Pain* 2008;24:305-15.
- Smeets RJ, Vlaeyen JW, Hidding A, Kester AD, van der Heijden GJ, Knottnerus JA. Chronic low back pain: physical training, graded activity with problem solving training, or both? The one-year post-treatment results of a randomized controlled trial. *Pain* 2008;134:263-76.
- Smeets RJ, Vlaeyen JW, Hidding A, Kester AD, van der Heijden GJ, van Geel AC, et al. Active rehabilitation for chronic low back pain: cognitive-behavioral, physical, or both? First direct post-treatment results from a randomized controlled trial [ISRCTN22714229]. *BMC Musculoskeletal Disord* 2006;7:5.
- Smeets RJ, Vlaeyen JW, Kester AD, Knottnerus JA. Reduction of pain catastrophizing mediates the outcome of both physical and cognitive-behavioral treatment in chronic low back pain. *J Pain* 2006;7:261-71.
- Smith DW, Arnstein P, Rosa KC, Wells-Federman C. Effects of integrating therapeutic touch into a cognitive behavioral pain treatment program. Report of a pilot clinical trial. *J Holist Nurs* 2002;20:367-87.
- Soderlund A, Lindberg P. Cognitive behavioural components in physiotherapy management of chronic whiplash associated disorders (WAD) – a randomised group study. *G Ital Med Lav Ergon* 2007;29:A5-11.
- Sogaard R, Bunger CE, Laurberg I, Christensen FB. Cost-effectiveness evaluation of an RCT in rehabilitation after lumbar spinal fusion: a low-cost, behavioural approach is cost-effective over individual exercise therapy. *Eur Spine J* 2008;17:262-71.
- Thieme K, Flor H, Turk DC. Psychological pain treatment in fibromyalgia syndrome: efficacy of operant behavioural and cognitive behavioural treatments. *Arthritis Res Ther* 2006;8:R121.
- Thieme K, Gromnica-Ihle E, Flor H. Operant behavioral treatment of fibromyalgia: a controlled study. *Arthritis Rheum* 2003;49:314-20.
- Thieme K, Turk DC, Flor H. Responder criteria for operant and cognitive-behavioral treatment of fibromyalgia syndrome. *Arthritis Rheum* 2007;57:830-6.
- Thorn BE, Cross TH, Walker BB. Meta-analyses and systematic reviews of psychological treatments for chronic pain: relevance to an evidence-based practice. *Health Psychol* 2007;26:10-2.
- Walach H, Lewith G, Bosch H, Utts J. Spiritual healing as a therapy for chronic pain: a randomized, clinical trial. *Pain* 2002;96:403-5; author reply 405-6.
- van den Hout JH, Vlaeyen JW, Heuts PH, Zijlema JH, Wijnen JA. Secondary prevention of work-related disability in nonspecific low back pain: does problem-solving therapy help? A randomized clinical trial. *Clin J Pain* 2003;19:87-96.
- van der Windt D, Hay E, Jellema P, Main C. Psychosocial interventions for low back pain in primary care: lessons learned from recent trials. *Spine* 2008;33:81-9.
- van Tulder MW, Esmail R, Bombardier C, Koes BW. Back schools for non-specific

- low back pain. *Cochrane Database Syst Rev* 2000;CD000261.
- van Tulder MW, Koes B, Malmivaara A. Outcome of non-invasive treatment modalities on back pain: an evidence-based review. *Eur Spine J* 2006;15 Suppl 1:S64-81.
- van Tulder MW, Ostelo R, Vlaeyen JW, Linton SJ, Morley SJ, Assendelft WJ. Behavioral treatment for chronic low back pain: a systematic review within the framework of the Cochrane Back Review Group. *Spine* 2000;25:2688-99.
- van Tulder MW, Ostelo R, Vlaeyen JW, Linton SJ, Morley SJ, Assendelft WJ. Behavioral treatment for chronic low back pain: a systematic review within the framework of the Cochrane Back Review Group. *Spine* 2001;26:270-81.
- Wicksell RK, Ahlqvist J, Bring A, Melin L, Olsson GL. Can exposure and acceptance strategies improve functioning and life satisfaction in people with chronic pain and whiplash-associated disorders (WAD)? A randomized controlled trial. *Cogn Behav Ther* 2008;37:169-82.
- Williams DA, Cary MA, Groner KH, Chaplin W, Glazer LJ, Rodriguez AM, et al. Improving physical functional status in patients with fibromyalgia: a brief cognitive behavioral intervention. *J Rheumatol* 2002;29:1280-6.
- Williams KA, Petronis J, Smith D, Goodrich D, Wu J, Ravi N, et al. Effect of Iyengar yoga therapy for chronic low back pain. In: *Pain*; 2005. p 107-17.
- Vlaeyen JWS, de Jong JR, Onghena P, Kerckhoffs-Hanssen M, Kole-Snijders AMJ. Can pain-related fear be reduced? The application of cognitive-behavioural exposure in vivo. *Pain Research & Management* 2002;7:144-153.
- Woods MP, Asmundson GJ. Evaluating the efficacy of graded in vivo exposure for the treatment of fear in patients with chronic back pain: a randomized controlled clinical trial. *Pain* 2008;136:271-80.

3.4 Akupunktur, TENS, triggerpunktsbehandling

Studier som granskats men inte uppfyllde inklusionskriterierna eller kravet på minst medelhög kvalitet och relevans

Barron MC, Rubin BR. Managing osteoarthritic knee pain. *J Am Osteopath Assoc* 2007;107:ES21-7.

Chou R, Huffman LH. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Ann Intern Med* 2007;147:492-504.

Clauw DJ. Does acupuncture help reduce pain in patients with fibromyalgia? *Nat Clin Pract Rheumatol* 2005;1:76-7.

Ga H, Choi JH, Park CH, Yoon HJ. Acupuncture needling versus lidocaine injection of trigger points in myofascial pain syndrome in elderly patients – a randomised trial. *Acupunct Med* 2007;25:130-6.

Ga H, Choi JH, Park CH, Yoon HJ. Dry needling of trigger points with and without paraspinal needling in myofascial pain syndromes in elderly patients. *J Altern Complement Med* 2007;13:617-24.

Guo X. Comparison of therapeutic effects on fibromyalgia syndrome between dermal-neurological electric stimulation and electric acupuncture. *Chinese Journal of Clinical Rehabilitation* 2005;9:171-73.

Harris RE, Tian X, Williams DA, Tian TX, Cupps TR, Petzke F, et al. Treatment of fibromyalgia with formula acupuncture: investigation of needle placement, needle

stimulation, and treatment frequency. *J Altern Complement Med* 2005;11:663-71.

Hayden JA, van Tulder MW, Malmivaara AV, Koes BW. Meta-analysis: exercise therapy for nonspecific low back pain. *Ann Intern Med* 2005;142:765-75.

Hayden JA, van Tulder MW, Tomlinson G. Systematic review: strategies for using exercise therapy to improve outcomes in chronic low back pain. *Ann Intern Med* 2005;142:776-85.

He D, Hostmark AT, Veiersted KB, Medbo JI. Effect of intensive acupuncture on pain-related social and psychological variables for women with chronic neck and shoulder pain – an RCT with six month and three year follow up. *Acupunct Med* 2005;23:52-61.

Hsieh LL, Kuo CH, Lee LH, Yen AM, Chien KL, Chen TH. Treatment of low back pain by acupressure and physical therapy: randomised controlled trial. *BMJ* 2006;332:696-700.

Hsieh LL, Kuo CH, Yen MF, Chen TH. A randomized controlled clinical trial for low back pain treated by acupressure and physical therapy. *Prev Med* 2004;39:168-76.

Hurwitz EL, Carragee EJ, van der Velde G, Carroll LJ, Nordin M, Guzman J, et al. Treatment of neck pain: noninvasive interventions: results of the Bone and Joint Decade 2000–2010 Task Force on Neck Pain and Its Associated Disorders. *Spine* 2008;33:S123-52.

Inoue M, Kitakoji H, Ishizaki N, Tawa M, Yano T, Katsumi Y, Kawakita K.

- Relief of low back pain immediately after acupuncture treatment – a randomised, placebo controlled trial. *Acupunct Med* 2006;24:103-8.
- Itoh K, Katsumi Y, Hirota S, Kitakoji H. Effects of trigger point acupuncture on chronic low back pain in elderly patients – a sham-controlled randomised trial. *Acupunct Med* 2006;24:5-12.
- Itoh K, Katsumi Y, Kitakoji H. Trigger point acupuncture treatment of chronic low back pain in elderly patients – a blinded RCT. *Acupunct Med* 2004;22:170-7.
- Johnson M, Martinson M. Efficacy of electrical nerve stimulation for chronic musculoskeletal pain: a meta-analysis of randomized controlled trials. *Pain* 2007; 130:157-65.
- Kalauokalani D, Cherkin DC, Sherman KJ. A comparison of physician and non-physician acupuncture treatment for chronic low back pain. *Clin J Pain* 2005; 21:406-11.
- Kavcic NS, Lehman GH, McGill SM. Effect of modulated TENS of muscle activation, oxygenation, and pain: Searching for a physiological mechanism. *Journal of Musculoskeletal Pain* 2005;13:19-30.
- Khadilkar A, Milne S, Brosseau L, Robinson V, Saginur M, Shea B, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic low-back pain. *Cochrane Database Syst Rev* 2005; CD003008.
- Koke AJ, Schouten JS, Lamerichs-Geelen MJ, Lipsch JS, Waltje EM, van Kleef M, Patijn J. Pain reducing effect of three types of transcutaneous electrical nerve stimulation in patients with chronic pain: a randomized crossover trial. *Pain* 2004; 108:36-42.
- Kostopoulos D, Nelson AJ, Ingber RS, Larkin RW. Reduction of spontaneous electrical activity and pain perception of trigger points in the upper trapezius muscle through trigger point compression and passive stretching. *Journal of Musculoskeletal Pain* 2008;16:266-78.
- Lee MS, Pittler MH, Shin BC, Kong JC, Ernst E. Bee venom acupuncture for musculoskeletal pain: a review. *J Pain* 2008; 9:289-97.
- Lucas B. Treatment options for patients with osteoarthritis of the knee. *Br J Nurs* 2005;14:976-81.
- Luijsterburg PA, Verhagen AP, Ostelo RW, van Os TA, Peul WC, Koes BW. Effectiveness of conservative treatments for the lumbosacral radicular syndrome: a systematic review. *Eur Spine J* 2007; 16:881-99.
- MacPherson H, Thorpe L, Thomas K, Campbell M. Acupuncture for low back pain: traditional diagnosis and treatment of 148 patients in a clinical trial. *Complement Ther Med* 2004;12: 38-44.
- Madsen MV, Gotzsche PC, Hrobjartsson A. Acupuncture treatment for pain: systematic review of randomised clinical trials with acupuncture, placebo acupuncture, and no acupuncture groups. *BMJ* 2009;338:a3115.
- Manheimer E, Linde K, Lao L, Bouter LM, Berman BM. Meta-analysis: acupuncture for osteoarthritis of the knee. *Ann Intern Med* 2007;146:868-77.

- Manheimer E, White A, Berman B, Forys K, Ernst E. Meta-analysis: acupuncture for low back pain. *Ann Intern Med* 2005; 142:651-63.
- Mayhew E, Ernst E. Acupuncture for fibromyalgia – a systematic review of randomized clinical trials. *Rheumatology (Oxford)* 2007;46:801-4.
- Mirza SK, Deyo RA. Systematic review of randomized trials comparing lumbar fusion surgery to nonoperative care for treatment of chronic back pain. *Spine* 2007;32:816-23.
- Oosterhof J, De Boo TM, Oostendorp RA, Wilder-Smith OH, Crul BJ. Outcome of transcutaneous electrical nerve stimulation in chronic pain: short-term results of a double-blind, randomised, placebo-controlled trial. *J Headache Pain* 2006;7:196-205.
- Oosterhof J, Samwel HJ, de Boo TM, Wilder-Smith OH, Oostendorp RA, Crul BJ. Predicting outcome of TENS in chronic pain: a prospective, randomized, placebo controlled trial. *Pain* 2008;136: 11-20.
- Poitras S, Brosseau L. Evidence-informed management of chronic low back pain with transcutaneous electrical nerve stimulation, interferential current, electrical muscle stimulation, ultrasound, and thermotherapy. *Spine J* 2008;8: 226-33.
- Salter GC, Roman M, Bland MJ, MacPherson H. Acupuncture for chronic neck pain: a pilot for a randomised controlled trial. *BMC Musculoskeletal Disord* 2006;7:99.
- Scott NA, Guo B, Barton PM, Gerwin RD. Trigger point injections for chronic non-malignant musculoskeletal pain: a systematic review. *Pain Med* 2009; 10:54-69.
- Smania N, Corato E, Fiaschi A, Pietropoli P, Aglioti SM, Tinazzi M. Repetitive magnetic stimulation: a novel therapeutic approach for myofascial pain syndrome. *J Neurol* 2005;252:307-14.
- Thompson JW, Bower S, Tyrer SP. A double blind randomised controlled clinical trial on the effect of transcutaneous spinal electroanalgesia (TSE) on low back pain. *Eur J Pain* 2008;12:371-7.
- Tough EA, White AR, Cummings TM, Richards SH, Campbell JL. Acupuncture and dry needling in the management of myofascial trigger point pain: a systematic review and meta-analysis of randomised controlled trials. *Eur J Pain* 2009;13: 3-10.
- Trinh K, Graham N, Gross A, Goldsmith C, Wang E, Cameron I, Kay T. Acupuncture for neck disorders. *Spine* 2007;32: 236-43.
- Trinh KV, Graham N, Gross AR, Goldsmith CH, Wang E, Cameron ID, Kay T. Acupuncture for neck disorders. *Cochrane Database Syst Rev* 2006;3: CD004870.
- van der Velde G, van Tulder M, Cote P, Hogg-Johnson S, Aker P, Cassidy JD, et al. The sensitivity of review results to

- methods used to appraise and incorporate trial quality into data synthesis. *Spine* 2007;32:796-806.
- Wang C, Xiong Z, Deng C, Yu W, Ma W. Miniscalpel-needle versus triggerpoint injection for cervical myofascial pain syndrome: a randomized comparative trial. *J Altern Complement Med* 2007;13:14-6.
- Vas J, Perea-Milla E, Mendez C, Silva LC, Herrera Galante A, Aranda Regules JM, et al. Efficacy and safety of acupuncture for the treatment of non-specific acute low back pain: a randomised controlled multicentre trial protocol [ISRCTN65814467]. *BMC Complement Altern Med* 2006;6:14.
- Witt C, Brinkhaus B, Jena S, Linde K, Streng A, Wagenpfeil S, et al. Acupuncture in patients with osteoarthritis of the knee: a randomised trial. *Lancet* 2005; 366:136-43.
- Witt CM, Jena S, Brinkhaus B, Liecker B, Wegscheider K, Willich SN. Acupuncture for patients with chronic neck pain. *Pain* 2006;125:98-106.
- Witt CM, Jena S, Selim D, Brinkhaus B, Reinhold T, Wruck K, et al. Pragmatic randomized trial evaluating the clinical and economic effectiveness of acupuncture for chronic low back pain. *Am J Epidemiol* 2006;164:487-96.
- Yokoyama M, Sun X, Oku S, Taga N, Sato K, Mizobuchi S, et al. Comparison of percutaneous electrical nerve stimulation with transcutaneous electrical nerve stimulation for long-term pain relief in patients with chronic low back pain. *Anesth Analg* 2004;98:1552-6, table of contents.
- Yuan J, Kerr D, Park J, Liu XH, McDonough S. Treatment regimens of acupuncture for low back pain – a systematic review. *Complement Ther Med* 2008;16:295-304.
- Yuan J, Purepong N, Kerr DP, Park J, Bradbury I, McDonough S. Effectiveness of acupuncture for low back pain: a systematic review. *Spine* 2008;33: E887-900.
- Zambito A, Bianchini D, Gatti D, Viapiana O, Rossini M, Adami S. Interferential and horizontal therapies in chronic low back pain: a randomized, double blind, clinical study. *Clin Exp Rheumatol* 2006;24:534-9.